

Proof of the Geometric Langlands Conjecture

Dima Arinkin, Justin Campbell, Lin Chen, Ambrus Dhillon, Joakim Fargnason,
Dennis Gaitsgory, Andreas Hayash, Kevin Lin, Sam Raskin, Nick Rosenblyum, Yifei Zhao.

Tutorial 1.1 (Justin Campbell) Introduction to Bun_G .

$\text{char } k = 0$, $k = \bar{k}$. $G = \text{conn'd reductive gp} / k$

pt/G classifying stack of G

$\text{Map}(S, \text{pt}/G) = \left\{ \begin{array}{c} \text{total locally trivial principal } G\text{-bundles on } S \\ \text{affine scheme} \end{array} \right\}$

E.g. pt/GL_n classifies rank n vector bundles.

X - smooth conn'd projective curve $/k$.

$\text{Bun}_G := \text{Map}(X, \text{pt}/G)$.

$\text{Map}(S, \text{Bun}_G) = \{ G\text{-bundles on } S \times X \}$.

Prop Bun_G is an algebraic stack, locally of finite type $/k$.

More precisely,

i) $\Delta: \text{Bun}_G \rightarrow \text{Bun}_G \times \text{Bun}_G$ is affine.

ii) \exists scheme U locally of finite type $/k$ and smooth surj. mor. $U \rightarrow \text{Bun}_G$
(an "atlas")

References: Sorger, notes from Dennis's 2009 seminar

Prop. Bun_G is smooth.

Proof sketch: An alg stack locally of fin. type has a tangent complex which (for sufficiently nice stacks) is a perfect complex.

Such a stack is smooth \Leftrightarrow its tangent complex is connective.

(i.e. concentrated in nonpositive cohomological degrees)

A standard calculation shows the ^{fiber of} tangent complex of Bun_G at a G -bundle P_G is

$$R\Gamma(X, g_X^* P_G)[1].$$

$$R^i \Gamma(X, \mathcal{F}) = 0, \quad \forall i > 1 \quad \text{because } \dim X = 1.$$

|
coherent

$\Rightarrow \text{Bun}_G$ is smooth.

Remark. None of the above used the assumption that G is reductive.

Aside: for G reductive, Bun_G is never quasi-compact.

$$\text{Eg. } \pi_0 \text{Bun}_{G_m} \stackrel{\deg}{\cong} \mathbb{Z}$$

For G nonabelian, the connected compts of Bun_G are not q. cpt.

Eg. in $\text{Bun}_{GL_2}(\mathbb{P}^1)$ we have

$$\mathcal{O}^{\oplus 2} \rightsquigarrow \mathcal{O}(1) \oplus \mathcal{O}(-1) \rightsquigarrow \mathcal{O}(2) \oplus \mathcal{O}(-2) \rightsquigarrow \dots \rightsquigarrow \mathcal{O}(n) \oplus \mathcal{O}(-n) \rightsquigarrow$$

The Hecke action

$x \in X(k)$. Suppose that $k = \mathbb{C}$ and we work in the analytic topology.

Then a G -bundle on the Riemann surface X is determined by

— a G -bundle P'_G on a small disk D_x^{an} ;

— a G -bundle P''_G on $X \setminus \{x\}$;

— an isom. $P'_G|_{D_x^{\circ \text{an}}} \xrightarrow{\alpha} P''_G|_{D_x^{\circ \text{an}}}$

$$D_x^{\circ \text{an}} := D_x^{\text{an}} - \{x\}$$

In alg geom, $\hat{\mathcal{O}}_x =$ completed local ring of X at x

$$\hat{K}_x = \text{Frac}(\hat{\mathcal{O}}_x)$$

$$D_{x,S} = \text{Spec}(\mathcal{O}_S \hat{\otimes} \hat{\mathcal{O}}_x), \quad D_{x,S}^{\circ} := \text{Spec}(\mathcal{O}_S \hat{\otimes} \hat{K}_x) = D_{x,S} \setminus (S \times \{x\}).$$

Theorem (Beauville - Laszlo)

on $D_{x,S}^{\circ}$

$$\text{Map}(S, \text{Bun}_G) \xrightarrow{\sim} \left\{ (P'_G, P''_G, \alpha) \right\}$$

$$\begin{array}{c} \downarrow \quad \downarrow \\ \text{on } D_{x,S} \quad \text{on } S \times (X \setminus \{x\}) \end{array}$$

$\mathcal{H}_{G,x} =$ stack of Hecke modifications at x .

$$\text{Map}(S, \mathcal{H}_{G,x}) = \left\{ (P_G^1, P_G^2, \alpha) : P_G^1, P_G^2, G\text{-bundles on } D_{x,S}, \right.$$

$$\left. P_G^1|_{D_{x,S}^{\circ}} \xrightarrow{\alpha} P_G^2|_{D_{x,S}^{\circ}} \right\}$$

$\mathcal{H}_{G,x}$ is a groupoid over $\text{Bun}_G(D_x)$ via composition of α 's.

$$\text{Beauville - Laszlo} \rightarrow \mathcal{H}_{G,x} \simeq_{\text{Bun}_G(D_x)} \text{Bun}_G$$

$$\mathcal{H}_{G,x} \times_{\text{Bun}_G(D_x)} \text{Bun}_G \longrightarrow \text{Bun}_G$$

Uniformization

$L_x G$ = loop group of G at x

\cup

$L_x^+ G$ = arc group of G at x

$$\text{Map}(S, L_x G) = \text{Map}(\dot{D}_{x,S}, G)$$

$$\text{Map}(S, L_x^+ G) = \text{Map}(D_{x,S}, G)$$

Prop 1) $L_x^+ G$ is an affine group scheme

ii) $L_x G$ is an ind-affine group ind-scheme.

Observation: locally on S , any G -bundle on $D_{x,S}$ descends to S .

$$\Rightarrow \text{Bun}_G(D_x) \simeq \text{pt} / L_x^+ G$$

The same is true for G -bundles on $\dot{D}_{x,S}$ which extend to $D_{x,S}$.

$$\Rightarrow \mathcal{H}_{G,x} = \text{Bun}_G(D_x) \times_{\text{Bun}_G(\dot{D}_x)} \text{Bun}_G(D_x) \simeq \text{pt} / L_x^+ G \times_{\text{pt} / L_x G} \text{pt} / L_x^+ G \simeq L_x^+ G \backslash L_x G / L_x^+ G$$

$\text{Gr}_{G,X} = \text{affine Grassmannian} := \mathbb{A}_X^1 G / \mathbb{A}_X^+ G$

(
ind-proper ind-scheme

$$\mathcal{H}_{G,X} = \mathbb{A}_X^+ G \backslash \text{Gr}_{G,X}$$

Tutorial 1.2 (Dima Arinkin) Local Systems.

$\text{char } \mathbb{C} = 0$, $\bar{\mathbb{C}} = \mathbb{C}$. $X = \text{smooth proj. curve}$, $G = \text{reductive group}$.

Def. $\text{LocSys} = \text{LocSys}_{G,X} = \{ G\text{-local systems on } X \}$
 de Rham local systems

1st approach. Local system:


G -bundle w/ an (integrable) connection

eg: If $G = \text{GL}(n)$,

(E, ∇) : $E = \text{rk } n \text{ v. bundle on } X$

$\nabla: E \rightarrow E \otimes \Omega_X$ \mathbb{C} -linear

$$\nabla(fs) = f \nabla s + s \otimes df$$



$(E, \nabla) \in \text{LocSys}_G$

Bun_G
 \uparrow
 E

For fixed E , ∇ 's form an affine space
 (over $\text{R}^1(X, \text{End}(E) \otimes \Omega_X)$)

Properties. LocSys is a (derived)
 alg stack of finite type, quasi-smooth
 (derived l.c.i.)

Remark Why is LocSys only quasi-smooth
 while Bun_G is smooth?

- because $H^i(X, \text{loc. sys}) = 0$ for $i > 1$.

but $H_{\text{dR}}^i(X, \text{loc. sys}) = 0$ for $i > 2$.

Example $G = GL_2$, $X = \mathbb{P}^1$

Bun_G

$$\begin{aligned} \mathcal{O}^2 &\rightsquigarrow \mathcal{O}(1) \oplus \mathcal{O}(-1) \rightsquigarrow \mathcal{O}(2) \oplus \mathcal{O}(-2) \rightsquigarrow \dots \\ \mathcal{O} \oplus \mathcal{O}(-1) &\rightsquigarrow \mathcal{O}(1) \oplus \mathcal{O}(-2) \rightsquigarrow \dots \end{aligned}$$

So connections on \mathcal{O}^2 ? $d + \begin{pmatrix} a_{11}(x) & a_{12}(x) \\ a_{21}(x) & a_{22}(x) \end{pmatrix} dx$ $x \rightsquigarrow \frac{1}{z}$ polynomials

$$a_{11}(x) = a_{11}^{(d)} x^d + a_{11}^{(d-1)} x^{d-1} + \dots + a_{11}^{(0)}$$

$$a_{11}(x) dx = \left(-\frac{a_{11}^{(d)}}{z^{d+2}} - \frac{a_{11}^{(d-1)}}{z^{d+1}} - \dots - \frac{0}{z} \right) d\zeta$$

Answer. $Loc Sys_{G, \mathbb{P}^1} = (pt \times_{\mathbb{G}} pt) / G$



2nd approach

Local systems = G -bundles on X_{dR} .

Def $X_{dR}(S) = Maps(S^{red}, X)$ [formalize parallel transport]


$Loc Sys = Maps(X_{dR}, pt/G)$, so $Loc Sys(S) = Maps(S \times X_{dR}, pt/G)$.



Ran space of X

$$Ran(X) = \{ \text{finite } \underset{\text{non empty}}{\text{subsets}} \text{ of } X \}.$$

Def. $\text{Maps}(S, \text{Ran } X) = \left\{ \text{finite nonempty subsets of } \text{Maps}(S, X_{dR}) \right\}$
 \parallel
 $\text{Maps}(S^{\text{red}}, X)$



Less formally. $n > 0$,

$$X_{dR}^n \rightarrow \text{Ran } X$$


$$X_{dR}^n \rightarrow X_{dR}^m \quad \text{for any surjective } \{1, \dots, m\} \rightarrow \{1, \dots, n\}.$$

(eg. diagonals, permutations)

Prop. $\varinjlim X_{dR}^n = \text{Ran } X$

What is a q-coh-sheaf on $\text{Ran } X$?


① a sheaf on X_{dR} (a q-coh. sheaf w/ a connection, i.e. D_X -module)

②  D_{X^2} -module
 \mathbb{G}_2 -equivariant

$$X^2: (x_1, x_2) \mapsto (x_2, x_1)$$


Def $\text{Rep}(G)_{\text{Ran}} = \mathcal{O}(\text{pt}/G)_{\text{Ran}}$

$$(\text{pt}/G)_{\text{Ran}} = \left\{ (\underline{x} \in \text{Ran } X, G\text{-local system on } D_{\underline{x}} - (\text{its formal nbhd})) \right\}$$



$$(pt/h)^n$$

Lecture 1 (Sam Raskin) Construction of the (coarse) Langlands functor



Goal: \mathbb{L}_G is an equiv.

Two ingredients in construction:

$$1) \text{ Poinc}_! = \text{Poinc}_!^{\text{vac}} \in D\text{-mod}(Bun_G) \quad \text{compact}$$

$$2) \text{ The spectral action } \text{Qcoh}(LS_G^V) \simeq D\text{-mod}(Bun_G) \quad \left. \begin{array}{l} \text{(related to Hecke action)} \end{array} \right\}$$

$$\text{Hom}(\text{Poinc}_! F) \simeq \text{coeff}(F)$$

commutes w direct sums
(\Leftrightarrow) w all colimits.

most of the
talk is about this.

A variant on Yoneda for $\text{Qcoh}(Y)$, Y alg stack s.t.

$$g \in \text{Qcoh}(Y) \longmapsto F_g : \text{Qcoh}(Y) \rightarrow \text{Vect}$$

$$H \longmapsto \Gamma(Y, g \otimes_Y H)$$

This functor commutes w colimits (assump. on Y)

Claim. $\mathcal{Q}\text{Coh}(\mathcal{Y}) \xrightarrow{\sim} \text{Funct}_{k\text{-lin}}^{\text{cts}}(\mathcal{Q}\text{Coh}(\mathcal{Y}), \text{Vect})$ is an equiv.

Assumptions. \mathcal{Y} is quasi-cpt, affine diag, $\mathcal{Q}\text{Coh}(\mathcal{Y})$ is dualisable (true for quasi-smooth)

True for LS_X^\vee .

To define $\mathbb{L}_{\mathcal{A}, \text{coarse}}(F)$, $F \in D\text{-mod}(\text{Bun}_{\mathcal{A}})$, I'll use dual Yoneda,

$\forall g \in \mathcal{Q}\text{Coh}(LS_X^\vee)$,

$$\Gamma(LS_X^\vee, g \otimes_{\mathcal{O}} \mathbb{L}_{\mathcal{A}, \text{coarse}}(F)) := \text{coet}(g \overset{\text{spectral action}}{\times} F)$$

Axioms of GLC: $\text{Point}! \mapsto \mathcal{O}_{LS_X^\vee}$.

Def. of $\mathbb{L}_{\mathcal{A}, \text{coarse}}$ is forced by this requirement.

Prop $\mathbb{L}_{\mathcal{A}, \text{coarse}}$ is the unique $\mathcal{Q}\text{Coh}(LS_X^\vee)$ -linear functor s.t. TFDC:

$$\begin{array}{ccc} D\text{-mod}(\text{Bun}_{\mathcal{A}}) & \xrightarrow{\mathbb{L}_{\mathcal{A}, \text{coarse}}} & \mathcal{Q}\text{Coh}(LS_X^\vee) \\ & \searrow \text{coet} & \swarrow \Gamma(LS_X^\vee, -) \\ & \text{Vect} & \end{array}$$

\hookrightarrow

Poincaré sheaf, $N \subset B \twoheadrightarrow T$
 unip Bred Cartan
 rad of B

Think.

$$\text{Bun}_N = \text{Bun}_B \times_{\text{Bun}_T} \{p_T^{\text{triv}}\}$$

Example $G = \text{SL}_2$, $B = \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix}$, $N = \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}$, $T = G_m = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$

$$\text{Bun}_B = \{0 \rightarrow \mathcal{L} \rightarrow \mathcal{E} \rightarrow \mathcal{L}^{-1} \rightarrow 0\}$$

$$\text{Bun}_T = \{\mathcal{L}\}$$

$$\text{Bun}_N = \{0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{E} \rightarrow \mathcal{O}_X \rightarrow 0\}$$

$$\text{Bun}_N^{\Omega} := \text{Bun}_B \times_{\text{Bun}_T} \{p(\Omega_X)\},$$

$$2p = \sum_{\substack{\alpha > 0 \\ \text{root}}} \alpha : G_m \rightarrow T$$

Choose $\Omega_X^{1/2}$ square root of Ω_X

$$p(\Omega_X) := 2p(\Omega_X^{1/2})$$

Ex. $G = \text{SL}_2$, $\text{Bun}_N^{\Omega} = \{0 \rightarrow \Omega^{1/2} \rightarrow \mathcal{E} \rightarrow \Omega^{-1/2} \rightarrow 0\}$

$$= \underline{R}\Gamma(\Omega_X)[1] \rightarrow \underline{H}^1(X, \Omega_X) = \mathbb{A}^1$$

For gen'l G , each simple root $\leadsto N \rightarrow G_a$

$$\text{Bun}_N^{\Omega} \rightarrow \text{Bun}_{G_a}^{\Omega} = \underline{R}\Gamma(X, \Omega) \rightarrow \mathbb{A}^1$$

$$\leadsto \text{Bun}_N^{\Omega} \rightarrow \mathbb{A}^{2k(G)} \xrightarrow{\text{sum conds}} \mathbb{A}^1$$

ψ

$$\begin{pmatrix} 1 & \Omega & \Omega^2 \\ & 1 & \Omega \\ & & 1 \end{pmatrix}$$

$\text{Poinc!} := p_{N!}(\psi^*(\text{exp}))$ where $p_N: \text{Bun}_N^{\mathbb{R}} \rightarrow \text{Bun}_G$ is the projection

$$\text{exp} = (\mathcal{O}_{\mathbb{A}^1}, \nabla = d - dt) \in D\text{-mod}(\mathbb{A}^1)$$

$$\text{Coet}(F) = \underset{\substack{\uparrow \\ dR \text{ coh}}}{\text{dR}} \left(\text{Bun}_N^{\mathbb{R}}, p_N^!(F) \otimes \psi^*(\text{exp}) \right) [\text{shift?}]$$

Example. $G = \text{Gm}$, $\text{Poinc!} = \delta_{\text{triv!}} \in D\text{-mod}(\text{Bun}_{\text{Gm}})$

$$\text{pt} \xrightarrow[\substack{\text{fibers} \\ \text{are Gm}}]{\text{triv}} \text{Bun}_{\text{Gm}} \quad ! - \text{pushforward of const. sheaf.}$$

Hecke action:

$x \in X$, Justin said: Hecke groupoid acting on Bun_G .

$$\text{Geometric Satake: } \text{Rep } \check{G} \longrightarrow D\text{-mod}(\mathcal{H}_{G,x})$$


$$V^\lambda \longmapsto \text{IC}_{\mathcal{H}_{G,x}^\lambda} [?]$$

$$V \in \text{Rep } \check{G}, \quad x \in X, \quad H_{V,x}: D\text{-mod}(\text{Bun}_G) \longrightarrow D\text{-mod}(\text{Bun}_G)$$

Hecke functor

pull-back to $\mathcal{H}_x^{\text{global}}$, \otimes w/ Satake sheaf, pushforward

$$\mathcal{H}_x^{\text{glob}} = \{ p_G, \tilde{p}_G \in \text{Bun}_G + d: p_G|_{X \setminus x} \xrightarrow{\sim} \tilde{p}_G|_{X \setminus x} \}$$



Variant 1. Don't fix $x \in X$

$$\begin{array}{ccc}
 V \in \text{Rep } \check{G} & , & H_V: D\text{-mod}(Bun_G) \rightarrow D\text{-mod}(Bun_G \times X) \\
 & \searrow H_{V,x} & \downarrow (id \times x)! \\
 & & D\text{-mod}(Bun_G)
 \end{array}$$

Variant 2. $V, w \in \text{Rep } \check{G}$ or $\{V_i\}_{i \in I}$

$$H_V \circ H_w: D\text{-mod}(Bun_G) \rightarrow D\text{-mod}(Bun_G \times X^2)$$

Relations from geom Satake:

a) $\text{Swap}_{X^2} \circ H_V \circ H_w = H_w \circ H_V$

b) restricting along $X \subset X \times X$, obtain $H_V \circ H_w$

a) & b) compatible.

Variant 3

$\text{Rep } \check{h}_{\text{Ran}}$ acts on $D\text{-mod}(\text{Bun}_h)$

$$V_x \text{ etc.} \quad \begin{array}{|c|c|c|} \hline x^1 & v_1 & \\ \hline y^1 & v_2 & v_3 \\ \hline & z & \\ \hline \end{array} \quad X$$

\leadsto obj. of $\text{Rep } \check{h}_{\text{Ran}}$ acts by $H_{V_1, x} \circ H_{V_2, y} \circ H_{V_3, z}$

$\text{Rep } \check{h}_{\text{Ran}}$

$$\downarrow \text{Loi}^{\text{spec}} = \text{evaluation bundles}$$

$$\mathcal{Q}\text{coh}(LS_h^v)$$

$x \in \text{Ran}$

$$LS_x \rightarrow (\text{pt}/h)_{\text{Ran}} \Big|_x$$

Loi^{spec} is pull back.

Thm (Gaitsgory - Rozenblyum) Loi^{spec} is a quotient functor (Ref: GLC IV)

Thm (Drinfeld - Gaitsgory) action of $\text{Rep } \check{h}_{\text{Ran}}$ on $D\text{-mod}(\text{Bun}_h)$ factors through action of $\mathcal{Q}\text{coh}(LS_h^v)$

$(V \in \text{Ker}(\text{Loi}^{\text{spec}}), \text{ acts by zero on } D\text{-mod}(\text{Bun}_h))$ "generalized vanishing conj."
+ input from GLC 2 & GLC 4


Tutorial 1.3 (Dirac Arinkin) IndCoh & Sing

① Singular support : refinement of support to measure direction of non-vanishing
for coherent sheaves on a quasi-smooth derived scheme (eg. l.c.i. scheme)

Apply to LocSys.

q. smooth: cotangent complex (Tor amplitude)
is in degrees -1 and above.

locally $X = \text{pt} \times_{\text{Smooth}}$



$$\mathcal{F} \in \text{Coh}(X) (= D_{\text{Coh}}^b(X))$$


Zariski closed & conical

$$\begin{array}{c} \mathcal{F} \\ \uparrow \\ \text{Coh}(X) \end{array} \rightsquigarrow \text{Sing Supp } \mathcal{F} \subset \text{Sing}(X)$$

$$\begin{array}{c} \parallel \\ H^{-1} T^* X \\ \parallel \\ \text{Spec}_X \text{Sym}^* H^1(TX) \end{array}$$

$$\text{Ex. } \mathbb{A}^n \supset X = \{x : b_1(x) = \dots = b_k(x) = 0\}$$

$$\text{Sing } X = \left\{ \begin{array}{c} (x, a) \\ \uparrow \quad \uparrow \\ X \quad \mathbb{A}^k \end{array} : a_1 db_1(x) + \dots + a_k db_k(x) = 0 \right\}$$



Properties. ① F has bounded tor amplitude (i.e., $F \in \text{Perf}(X)$)

$$\Leftrightarrow \text{Sing Supp } F \subset \text{zero section}$$

$$\textcircled{2} \quad F_1 \rightarrow F_2 \rightarrow F_3 \quad \Rightarrow \quad \text{Sing Supp } F_2 \subset \text{Sing Supp } F_1 \cup \text{Sing Supp } F_3.$$

③ Behavior w.r.t. maps $f: Y \rightarrow X$:

$$\text{e.g., pullback } f^* \text{ for smooth } f: \quad \text{Sing } Y = \text{Sing } X \times_X Y$$

Def (sketch) $H^1(TX)$ maps to $\text{Ext}^2(F, F)$

Thm (Liljekvist) $\text{Ext}^*(F, F)$ is f.g. as a $\text{Sym}^*(H^1(TX))$ -module.

$\text{Sing Supp } F := \text{support of this module.}$

$\xrightarrow{\quad \Sigma \quad}$

Application to Loc Sys

Let \mathcal{X} be a quasi-smooth derived stack, define $\text{Sing } \mathcal{X}$ and $\text{Sing Supp } F$ for F on \mathcal{X} via smooth covers (using ③)

Example. $\text{Sing}(\text{Loc Sys}_G)$

$$(L \in \text{Loc Sys}_G, \quad \zeta \in H^{-1}(T_L^* \text{Loc Sys}_G))$$

$$H^1 T_L^*(\text{Loc Sys}_G) = H_{dR}^2(X, \mathfrak{g}_L)$$

$$\zeta \in H^1 T_L^*(\text{Loc Sys}_G)^* = H_{dR}^2(X, \mathfrak{g}_L)^* = H_{dR}^0(X, \mathfrak{g}_L) \quad \text{infinitesimal symmetry of } L$$

Def. $\text{Nilp}^{\text{global nilp cone}} \subset \text{Sing}(\text{Loc Sys})$
 $\{ (L, \mathbb{Z}) : \mathbb{Z} \text{ is nilpotent} \}$



Ind coherent sheaves on a (derived) scheme

$$\mathcal{Q}\text{Coh}(X) \subset \text{Ind Coh}(X)$$

X f.type, \mathcal{O}_X bounded

$$\text{"} \mathcal{D}_{qc}(X) \text{"}$$

Compact obj. $\xrightarrow{\text{cocomplete (admits } \varinjlim)}$

$$(\mathcal{Q}\text{Coh}(X))^{\text{compact}} \xleftarrow{\text{---}} \mathcal{Q}\text{Coh}(X) - \text{compactly generated}$$

\parallel

$$\text{"} \text{D-section Perf}(X) \text{"} \xrightarrow{\text{ind-completion}} \text{Ind}(\text{Perf}(X)) \quad \left(\text{adding formal colimits} \right)$$

cat. of functors $\text{Perf}(X)^{\text{op}} \rightarrow \text{Vect}$



$$\mathcal{D}^{\text{Sing } X} \text{Coh}(X) \xrightarrow{\text{ind-completion}} \text{Ind}(\text{Coh}(X))$$

Given conical $\mathcal{N} \subset \text{Sing}(X)$

$$\text{define } \text{Ind Coh}(X)_{\mathcal{N}} = \text{Ind} \left(\{ F \in \text{Coh}(X) : \text{Sing Supp}(F) \subset \mathcal{N} \} \right)$$

Remark Def'n works locally; for q -smooth stacks \mathcal{X} , define $\text{Ind Coh}(\mathcal{X})_{\mathcal{N}}$

$$= \lim_{\substack{S \rightarrow \mathcal{X} \\ S \text{ smooth}}} \text{Ind Coh}(S)_{\mathcal{N} \times S}$$

$$\mathcal{Q}\text{Coh}(\text{Loc Sys}_{\check{a}}^{\text{nilp}}) \subset \text{Ind Coh}(\text{Loc Sys}_{\check{a}}^{\text{nilp}})$$

Lecture 2 (Dennis Gaitsgory)

$$1) \operatorname{Hom}(\mathcal{Y}, \operatorname{pt}/\mathcal{A}) = \text{sym. monoidal functors } \operatorname{Rep}(\mathcal{A}) \longrightarrow \operatorname{Qcoh}(\mathcal{Y})$$

\nwarrow
 right t-exact

$$ii) \operatorname{Hom}(S, LS_{\mathcal{A}}^{\vee}) = \text{sym. mon. right t-exact functors } \operatorname{Rep}(\check{\mathcal{A}}) \longrightarrow \operatorname{Qcoh}(S) \otimes \operatorname{Dmod}(X)$$

$$\operatorname{Qcoh}(S_1) \otimes \operatorname{Qcoh}(S_2) = \operatorname{Qcoh}(S_1 \times S_2)$$

$$2) (\operatorname{pt}/\mathcal{A})_{\operatorname{Ran}} \leftarrow \text{give a def'n}$$

$$\downarrow$$

Ran

$$2') \operatorname{Rep}(\mathcal{A})_{\operatorname{Ran}} = \operatorname{Qcoh}((\operatorname{pt}/\mathcal{A})_{\operatorname{Ran}})$$

$$2-) \operatorname{Qcoh}(\mathcal{Y}) = \varprojlim_{S \rightarrow \mathcal{Y}} \operatorname{Qcoh}(S)$$

$$3) \quad \begin{array}{ccc} LS_{\check{\mathcal{A}}, X} & \xleftarrow{p} & LS_{\check{\mathcal{A}}, X} \times \operatorname{Ran} \xrightarrow{ev} (\operatorname{pt}/\check{\mathcal{A}})_{\operatorname{Ran}} \\ & & \searrow \quad \swarrow \\ & & \operatorname{Ran} \end{array}$$

$$\operatorname{Rep}(\check{\mathcal{A}})_{\operatorname{Ran}} \xrightarrow[p_* \circ ev^*]{\operatorname{Loc}_{\check{\mathcal{A}}}^{\operatorname{spec}}} \operatorname{Qcoh}(LS_{\check{\mathcal{A}}})$$



$$3') \quad x_1, \dots, x_n \in X$$

$$V_1, \dots, V_n \in \operatorname{Rep}(\check{\mathcal{A}}) \rightsquigarrow \bigotimes_{i=1}^n V_{i, x_i} \in \operatorname{Rep}(\check{\mathcal{A}})_{\operatorname{Ran}}$$

$$\left(\operatorname{Loc}_{\check{\mathcal{A}}}^{\operatorname{spec}} \left(\bigotimes_{i=1}^n V_{i, x_i} \right) \right)_{\sigma} = \bigotimes_{i=1}^n \left((V_i)_{\sigma} \right)_{x_i}$$

$$\sigma \in LS_{\check{\mathcal{A}}}$$

3") Thm $\text{Loc}^{\text{Spec}}_{\chi}$ is a quotient.



$$\text{Rep}(\check{h})_{\text{Ran}} \xrightarrow{\text{Sat}_h^{\text{nv}}} \text{Dmod}(U_{\text{Ran}}^{\text{loc}})$$

$$\begin{array}{ccc}
 \text{Rep}(\check{h})_{\text{Ran}} \otimes \text{Dmod}(\text{Bun}_{\chi}) & \xrightarrow{\text{Hecke}} & \text{Dmod}(\text{Bun}_{\chi}) \\
 \downarrow & & \downarrow \\
 V & & M
 \end{array}$$

$$\text{Hecke}(V \otimes M) = \bar{h}_* \left(s^! \underset{\text{Sat}_h^{\text{nv}}(V)}{V} \otimes \bar{h}^!(M) \right)$$

Thm. The above binary operation factors through

$$\begin{array}{ccc}
 \text{Rep}(\check{h})_{\text{Ran}} \otimes \text{Dmod}(\text{Bun}_{\chi}) & & \\
 \downarrow \text{Loc}^{\text{Spec}}_{\chi} \otimes \text{Id} & \searrow \text{Hecke} & \\
 \mathcal{O}\text{coh}(LS_{\chi}^{\vee}) \otimes \text{Dmod}(\text{Bun}_{\chi}) & \longrightarrow & \text{Dmod}(\text{Bun}_{\chi})
 \end{array}$$

$$5) \text{Dmod}(\text{Bun}_{\chi}) \xrightarrow{\mathbb{L}_{h, \text{coarse}}} \mathcal{O}\text{coh}(LS_{\chi}^{\vee})$$

Poinc^{vac}!

$$\begin{array}{ccc}
 \text{Dmod}(\text{Bun}_{\chi}) & \xleftrightarrow{\mathbb{L}_{h, \text{coarse}}^L} & \mathcal{O}\text{coh}(LS_{\chi}^{\vee}) \\
 \text{Poinc}^{\text{vac}}! & & \downarrow \\
 & \xleftrightarrow{\mathbb{L}_{h, \text{coarse}}} & \mathcal{O}_{LS_{\chi}^{\vee}}
 \end{array}$$

$$\begin{array}{ccc}
 & \xrightarrow{\mathbb{L}_G} & \text{IndCoh}_{\text{nilp}}(LS_G^v) \\
 \text{Dmod}(Bun_G) & \xrightarrow{\mathbb{L}_{G, \text{coarse}}} & \text{QCoh}(LS_G^v) \\
 & & \downarrow \psi
 \end{array}$$

$$\text{IndCoh}(Y)^{>-\infty}$$

$\psi_Y \mid \leftarrow$ t-exact, equiv on eventually coconnective

$$\text{QCoh}(Y)^{>-\infty}$$

$$\text{IndCoh}_X(Y)$$

$$\downarrow$$

$$\text{QCoh}(Y)$$

$$\begin{array}{ccc}
 & \xrightarrow{\quad} & \text{IndCoh}_{\text{nilp}}(LS_G^v)^{>-\infty} \\
 \text{Dmod}(Bun_G)^c & \xrightarrow{\mathbb{L}_{G, \text{coarse}}} & \text{QCoh}_{\text{nilp}}(LS_G^v)^{>-\infty} \\
 & & \downarrow \psi
 \end{array}$$

Want to show

Thm 0. $\mathbb{L}_{G, \text{coarse}}$ sends compact objects in $\text{Dmod}(Bun_G)$ to eventually coconnective objects in $\text{QCoh}(LS_G^v)$.

Thm 1. $\text{Dmod}(Bun_G)^c \subset \text{Dmod}(Bun_G)^{>-\infty}$

Thm 2. $\mathbb{L}_{G, \text{coarse}}$ is left exact $[-1000]$

NB \mathbb{L}_G is not left-exact. (eg. apply to constant sheaf)

Thm 2' $\mathbb{L}_{G, \text{coarse}}$ is right t-exact $[1500]$. Cor. \mathbb{L}_G is right t-exact $[1500]$

Proof of Thm 1

\mathcal{Y} qc algebraic stack

$$D_{\text{mod}}(\mathcal{Y})^c \subset D_{\text{mod}}(\mathcal{Y})^{>-\infty}$$

$$\text{Ind Coh}(\mathcal{Y}) \xrightleftharpoons[\text{oblv}^2]{\text{ind}^2} D_{\text{mod}}(\mathcal{Y}) \quad \text{t-exact}$$

$$\begin{array}{c} \text{qc} \\ \downarrow \\ \mathcal{U} \hookrightarrow \text{Bun}_{\mathcal{U}} \end{array}$$

Every compact in $D_{\text{mod}}(\text{Bun}_{\mathcal{U}})$ is of the form $\varinjlim (F_{\mathcal{U}})$, $F_{\mathcal{U}} \in D_{\text{mod}}(\mathcal{U})^c$.
not defined in general

Claim. $\text{Bun}_{\mathcal{U}}$ is a union of \mathcal{U}' 's for which $j_!$ is defined.

$$\mathcal{U} \hookrightarrow \mathcal{U}'$$

Proof of Thm 2

$$M \in D_{\text{mod}}(\text{Bun}_{\mathcal{U}})^{>-\infty}, \quad \mathbb{L}_{\mathcal{U}, \text{coarse}}(M) = F$$

\mathcal{Y} - algebraic stack, $\text{Spa}(k) \xrightarrow{\text{ig}} \mathcal{Y}$

$$i_{\mathcal{Y}}^!(F)$$

BBT $\exists d$ s.t. $\forall \mathcal{Y}, i_{\mathcal{Y}}^!(F) \geq d \Rightarrow F \geq 0$

$\leftarrow \mathcal{Y}$ eventually connective?
allow all pts, not just k -pts.

Counter example. $\mathcal{Y} = \text{Spa } k[\zeta], \deg(\zeta) = -2, F = k[\zeta, \zeta^{-1}]$ i.e. all base change of \mathcal{Y} .

Counter example $\mathcal{Y} = \mathbb{A}^1, F = \bigoplus_n k(t)[n]$

Need to show $\forall \sigma \in LS_G^v(k), \text{ pt } \xrightarrow{i_\sigma} LS_G^v$

$$i_\sigma^! (\mathbb{L}_{G, \text{coarse}}(M)) \geq d.$$

$$\begin{array}{ccc} \text{[AGKRRV]} & LS_G^{\text{restr}} & \xrightarrow{i^{\text{spa}}} LS_G^v \\ & \downarrow i_\sigma & \\ & \text{pt} & \end{array}$$

Enough to show : $(i_\sigma)^! \cdot (i^{\text{spa}})^! \mathbb{L}_{G, \text{coarse}}(M) \geq d.$

$$D_{\text{mod Nilp}}(\text{Bun}_G) \subset D_{\text{mod}}(\text{Bun}_G)$$

$$\text{Thm [AGKRRV]} \quad D_{\text{mod Nilp}}(\text{Bun}_G) \simeq \mathcal{Q}\text{coh}(LS_G^{\text{restr}}) \otimes_{\mathcal{Q}\text{coh}(LS_G^v)} D_{\text{mod}}(\text{Bun}_G)$$

$$D_{\text{mod Nilp}}(\text{Bun}_G) \xrightleftharpoons[i^R]{i^!} D_{\text{mod}}(\text{Bun}_G)$$

$$\begin{array}{ccc} D_{\text{mod}}(\text{Bun}_G) & \xrightarrow{\mathbb{L}_{G, \text{coarse}}} & \mathcal{Q}\text{coh}(LS_G^v) \\ i \uparrow \downarrow i^R & & \cdot i_x^{\text{spa}} \uparrow \downarrow (i^{\text{spa}})^! \\ D_{\text{mod Nilp}}(\text{Bun}_G) & \xrightarrow[\mathbb{L}_{G, \text{coarse}}^{\text{restr}}]{} & \mathcal{Q}\text{coh}(LS_G^{\text{restr}}) \end{array}$$

$$(i^{\text{spa}})^! \cdot \mathbb{L}_{G, \text{coarse}}$$

$$\parallel$$

$$\mathbb{L}_{G, \text{coarse}}^{\text{restr}} \cdot i^R$$

$$(i_\sigma)^! \cdot \left(\mathbb{L}_{G, \text{coarse}}^{\text{restr}} \right) \cdot i^R(M)$$

t-exact [FR]

$D_{\text{mod}}(\text{Bun}_M)$ $\text{IndCoh}(LS_M)$

$$\begin{array}{ccc}
 D_{\text{mod}}(\text{Bun}_M) & & \text{IndCoh}(LS_M) \\
 \downarrow E_{\text{is}} & & \downarrow E_{\text{is}}^{\text{Spec}} \\
 D_{\text{mod}}(\text{Bun}_G) & \xrightarrow{\quad \mathbb{L}_G \quad} & \text{IndCoh}(LS_G)
 \end{array}$$

Bonus Material A (Johann Fargnani)

Plan Introduce other versions of GLC

- 1) Betti 2) Restricted 3) Temporal

§1 Betti GLC (Ben-Zvi - Nadler '16)

Let X be smooth proj. curve / \mathbb{C} , Fix $e = \bar{e}$ char. 0.

Goal: Define a version of GL replacing $LS_G^{\vee} \rightsquigarrow LS_G^{\text{Betti}} = \text{Hom}(\pi_1(X), \check{G})^{\vee} / G$

Spectral side

$$LS_G^{\text{Betti}} := \text{Maps}_{\text{Spec}}(X, B_{\check{G}}^{\vee})$$

$$LS_G^{\text{Betti}}(S) = \text{Maps}_{\text{Spec}}(X, \text{Maps}_{\text{Stk}}(S, B_{\check{G}}^{\vee}))$$

Alternatively, $g = g(X)$ genus of X

$$\pi_1(X) = \langle a_i, b_i, i=1, \dots, g: \prod_i [a_i, b_i] = 1 \rangle$$

$$LS_G^{\text{Betti}} = \check{G}^{2g} \times_{\check{G}} \{e\} / \check{G}$$

$$\underline{\text{Ex}} \quad X = \mathbb{P}^1, \quad LS_{\check{A}}^{\text{Betti}} = p_!^X \check{A}^p / \check{A}^v = \Omega \check{g} / \check{A}$$

$$\underline{\text{Ex}} \quad G = T, \quad \text{then} \quad LS_T^{\text{Betti}} \simeq \check{T}^{2g} \times \check{T}^X \{e\} / \check{T} \simeq \check{T}^{2g} \times \Omega \check{T}^X \times B\check{T}$$

Automorphic side.

For \mathcal{Y}/\mathbb{C} an alg stack. let $\text{Shv}^{\text{all}}(\mathcal{Y})$ to be the cat. of sheaves of \mathbb{C} -vector spaces on top. space underlying $\mathcal{Y}(\mathbb{C})^{\text{an}}$.

$$\text{i.e. } \text{Shv}^{\text{all}}(\mathcal{Y}) = \lim_{S \rightarrow \mathcal{Y}} \text{Shv}^{\text{all}}(S)$$

If $\Lambda \subset T^*\mathcal{Y}$ closed conical subset, we may consider

$$\text{Shv}_{\Lambda}^{\text{all}}(\mathcal{Y}) = \{F \in \text{Shv}^{\text{all}}(\mathcal{Y}) : \text{ss}(\mathcal{H}^i(F)) \subset \Lambda\}$$

$$\underline{\mathcal{Y}} = \text{Bun}_G. \quad \text{Recall} \quad T^*\text{Bun}_G = \{(P_G, \varphi) \in \Gamma(X, \mathfrak{g}_{P_G}^* \otimes \Omega_X)\}$$

$$\underline{\Lambda} = \text{Nilp} \subset T^*\text{Bun}_G \quad \text{subset of } (P_G, \varphi) : \varphi \text{ is nilpotent}$$

$$\underline{\text{Betti GLC}} : \quad \text{Shv}_{\text{Nilp}}^{\text{all}}(\text{Bun}_G) \rightrightarrows \text{IndCoh}_{\text{Nilp}}(LS_{\check{A}}^{\text{Betti}})$$

Motivation

1) Laumon '87 conjectured (proved in [AGKRRV'20]) that Hecke eigensheaves have nilp sing support.

2) Eis_! : $\text{Shv}^{\text{all}}(\text{Bun}_G) \rightarrow \text{Shv}^{\text{all}}(\text{Bun}_G)$ preserves nilp. sing. supp.

$$\underline{G=T} \quad \text{Nilp} = 0$$

$$\text{Spectral} \quad LS_{\check{T}}^{\text{Betti}} = \check{T}^{2g} \times \Omega \check{T}^* \times B\check{T}$$

$$\mathcal{Q}\text{coh}(LS_{\check{T}}^{\text{Betti}}) \simeq \mathcal{Q}\text{coh}(\check{T}^{2g}) \otimes \mathcal{Q}\text{coh}(\Omega \check{T}^*) \otimes \mathcal{Q}\text{coh}(B\check{T})$$

$$\simeq k[H_1(X) \otimes \check{\Lambda}]_{\text{-mod}} \otimes \text{Sym}(\check{\Lambda}[1])_{\text{-mod}} \otimes \text{Vect}^{\check{\Lambda}}$$

Automorphic side

$$\text{Bun}_T = \text{Pic}_T^\circ(X) \times BT \times \check{\Lambda}$$

$$\text{Shv}_0^{\text{all}}(\text{Bun}_T) \simeq k[\pi_1(\text{Pic}_T^\circ(X))]_{\text{-mod}} \otimes H_*(T)_{\text{-mod}} \otimes \text{Vect}^{\check{\Lambda}}$$

$$\pi_1(\text{Pic}_T^\circ(X)) = H_1(X) \otimes \check{\Lambda}$$

$$H_*(T) \simeq \text{Sym}(\check{\Lambda}[1])$$



§2. Restricted GLC.

Original motivation: define stack of étale local systems on X/\mathbb{F}_q .

For us: (Restricted GLC)

$$\begin{array}{ccc} & \nwarrow & \nearrow \\ (\text{Betti GLC}) & & (\text{dR GLC}) \end{array}$$

Return to dR-case

$$X/k = \bar{k} \quad \text{char } 0 \quad \text{Maps}(S, LS_{\check{A}}) = \left\{ \begin{array}{l} \text{right exact } \otimes\text{-functors} \\ \text{Rep}(\check{A}) \rightarrow \mathcal{Q}\text{coh}(S) \otimes D(X) \end{array} \right\}$$

Let $\mathcal{OLisse}(X) \subset D(X)$ subcat. of $F \in D(X) : H^i(F)$ union of $(v.b + \nabla)$

Def'n Maps $(S, LS_{\check{A}}^{rest}) = \left\{ \begin{array}{l} \text{right exact } \otimes\text{-functors :} \\ \text{Rep}(\check{A}) \rightarrow \mathcal{O}h(S) \otimes \mathcal{OLisse}(X) \end{array} \right\}$

$$L: LS_{\check{A}}^{rest} \rightarrow LS_{\check{A}}$$

Facts 1) L (bijection) surjective on k -ptr: $\text{Rep}(\check{A}) \rightarrow D(X) \xrightarrow{\quad} \mathcal{OLisse}(X)$

Moreover, for $\sigma \in LS_{\check{A}}$, $LS_{\check{A}}^{rest} \wedge_{\sigma} \simeq LS_{\check{A}} \wedge_{\sigma}$

2) Two $\sigma_1, \sigma_2 \in LS_{\check{A}}$ lie in the same conn'd comp. of $LS_{\check{A}}^{rest}$ iff they have isom. semisimplifications.

The conn'd comp. containing irred. σ is isom. to $LS_{\check{A}} \wedge_{\text{pt}/\text{Stab}_{\check{A}}(\sigma)}$ $\text{pt}/\text{Stab}_{\check{A}}(\sigma) \rightarrow LS_{\check{A}}$

Rank: σ semisimple if whenever it admits a reduction $\sigma_P^v \in LS_P^v$, it further admits reduction to $LS_{\check{M}} \rightarrow LS_P^v$.

σ local system, $\tilde{\sigma}$ semisimple local system, $\tilde{\sigma}$ is SS'n of σ if have reductions $\sigma_P^v, \tilde{\sigma}_P^v$ and they become iso. after $LS_{\check{M}}$.

Restricted hLC:

$$D_{\text{nilp}}(\text{Bun}_{\check{A}}) \simeq \text{Ind Coh}_{\text{nilp}}(LS_{\check{A}}^{rest})$$

Motivations: 1) Any obj. of $D_{\text{nilp}}(\text{Bun}_{\check{A}})$ is regular holonomic.

There is analogous Beilinson restricted hLC and they are equiv. by Riemann-Hilbert.
p. 25

2) Given cat. \mathcal{C}

$$\text{action } \mathcal{Q}\text{Coh}(LS_{\check{G}}^{\text{rest}}) \curvearrowright \mathcal{C}$$

$$\Leftrightarrow \text{Rep}(\check{G})^{\otimes I} \otimes \mathcal{C} \longrightarrow \mathcal{C} \otimes \mathcal{Q}\text{Lisse}(X)^{\otimes I}$$

\forall finite sets I + compatibilities

$$\text{Thm (Maddox-Yun)} \quad \text{Rep}(\check{G}) \otimes D(\text{Bun}_G) \longrightarrow D(\text{Bun}_G) \otimes D(X)$$

\cup

$$\text{Rep}(\check{G}) \otimes D_{\text{nilp}}(\text{Bun}_G) \longrightarrow D(\text{Bun}_G) \otimes \mathcal{Q}\text{Lisse}(X)$$

$$\leadsto \mathcal{Q}\text{Coh}(LS_{\check{G}}^{\text{rest}}) \curvearrowright D_{\text{nilp}}(\text{Bun}_G).$$

$$\text{In fact, } D_{\text{nilp}}(\text{Bun}_G) \simeq D(\text{Bun}_G) \otimes_{\mathcal{Q}\text{Coh}(LS_{\check{G}}^{\vee})} \mathcal{Q}\text{Coh}(LS_{\check{G}}^{\text{rest}})$$

$$\begin{aligned} \text{dR GLC} &\Rightarrow \text{Restricted GLC.} & \text{IndCoh}_{\text{nilp}}(LS_{\check{G}}^{\text{rest}}) &\simeq \text{IndCoh}_{\text{nilp}}(LS_{\check{G}}^{\vee}) \otimes_{\mathcal{Q}\text{Coh}(LS_{\check{G}}^{\vee})} \mathcal{Q}\text{Coh}(LS_{\check{G}}^{\text{rest}}) \\ \text{Restricted GLC} &\Rightarrow \text{dR GLC.} \end{aligned}$$

$$LS_{G_m}^{\text{rest}} = \coprod_{\sigma \in LS_{G_m}} LS_{G_m}^{\wedge} B G_m$$

$$LS_{G_m} \simeq B G_m \times \text{Loc} \times \Omega k,$$

$\text{Loc} =$ coarse moduli space of $2k+1$ v.b. on $X + \nabla$

Tempered GLC

$$D(\text{Bun}_G) \cong \text{Ind Coh}_{\text{nilp}}(LS_{\check{G}}^{\vee})$$

$$\begin{array}{c} \downarrow \qquad \qquad \qquad \uparrow \\ D(\text{Bun}_G)^{\text{temp}} \cong \text{QCoh}(LS_{\check{G}}^{\vee}) \end{array}$$

Q. How to char. $\text{QCoh}(LS_{\check{G}}^{\vee}) \subset \text{Ind Coh}_{\text{nilp}}(LS_{\check{G}}^{\vee})$?

A. Choose $x \in X$: $\text{QCoh}(LS_{\check{G}}^{\vee}) \cong \text{Ind Coh}_{\text{nilp}}(LS_{\check{G}}^{\vee}) \otimes_{\text{Ind Coh}_{\text{nilp}}(\mathcal{R}\check{G}/\check{G})} \text{QCoh}(\mathcal{R}\check{G}/\check{G}^{\vee})$

$$\text{Ind Coh}_{\text{nilp}}(\mathcal{R}\check{G}/\check{G}^{\vee}) \rightsquigarrow \text{Ind Coh}_{\text{nilp}}(LS_{\check{G}}^{\vee})$$

$$\begin{array}{c} \uparrow \\ \text{Rep}(\check{G}) \end{array}$$

Derived Satake

$$D(L_{x^+}^+ G \setminus L_x G / L_{x^+}^+ G) \cong \text{Ind Coh}_{\text{nilp}}(\mathcal{R}\check{G}/\check{G}^{\vee})$$

$$\text{So: } D(\text{Bun}_G)^{\text{temp}} := D(\text{Bun}_G) \otimes_{\text{Ind Coh}_{\text{nilp}}(\mathcal{R}\check{G}/\check{G}^{\vee})} \text{QCoh}(\mathcal{R}\check{G}/\check{G}^{\vee})$$

Tutorial 2.1 (Andreas Hayash)

Goal: Discuss functors coet + left adjoint Poinc !

~ Introduce Whittaker category on the affine Grassmannian.

Recall $\omega_{G, \underline{x}}$, $\forall \underline{x} \in \text{Ran}$, also $\omega_{G, \text{Ran}}$

$$\parallel$$

$$(L\mathfrak{h})_{\underline{x}} / (L^+\mathfrak{h})_{\underline{x}}$$

$$(L\mathfrak{h})_{\underline{x}} \simeq \omega_{G, \underline{x}} \rightsquigarrow (LN)_{\underline{x}} \simeq \omega_{G, \underline{x}}, \quad N \text{ unip. rad. of a Borel } B \subset G.$$

The group $(LN)_{\underline{x}}$ admits a character χ .

χ factors as

$$LN \longrightarrow L(N/[N, N]) \simeq \prod_I L(\mathfrak{a}_\alpha) \xrightarrow{\text{res}} \prod_I \mathfrak{a}_\alpha \xrightarrow{\chi_\alpha} \mathfrak{a}_\alpha$$

\downarrow
 vertices of the
 Dynkin diagram

χ_α
 character,
 restricted on
 each coord.

Rank Suppressing twists,

Def. $\text{Whit}^!(\mathfrak{h})_{\underline{x}} = \text{Dmod}(\omega_{G, \underline{x}})^{LN, \chi}$

ie. Objects of $\text{Whit}^!(\mathfrak{h})_{\underline{x}}$ are \mathbb{D} -modules on $\omega_{G, \underline{x}}$ + an isom.

$$\text{act}: LN \times \omega_{G, \underline{x}} \rightarrow \omega_{G, \underline{x}} \quad \text{act}^!(C) \simeq \chi^!(\exp) \boxtimes C.$$

$$m: \mathbb{H} \times \mathbb{H} \rightarrow \mathbb{H}, \quad m^!(F) \simeq F \boxtimes F + \text{cocycle cond.}$$

$$LN = \bigcup N_\alpha, \quad \text{can consider } D(\omega_{G, \underline{x}})^{N_\alpha, \chi} \xrightarrow{\text{oblv}_{\alpha, \beta}} D(\omega_{G, \underline{x}})^{N_\beta, \chi}$$

\downarrow b.f.

$$\text{Whit}^!(\mathfrak{h})_{\underline{x}} = \varinjlim_{\alpha} D(\omega_{G, \underline{x}})^{N_\alpha, \chi} = \bigcap_{\alpha} D(\omega_{G, \underline{x}})^{N_\alpha, \chi} \subset D(\omega_{G, \underline{x}})$$

$$\exists \text{ obj: } \text{Whit}^1(G)_{\underline{x}} \rightarrow D(\text{hr}_{G, \underline{x}}) \quad \text{fully faithful}$$

It admits a (non-cts) right adjoint $Av_{*}^{L_{N,X}} : D(W_{G,X}) \rightarrow Whitt^i(G)_X$

Rmk The cat. $\text{Whit}^!(G)_{\underline{x}}$ does not depend on the choice of x .

Variant of Whit, $\text{Whit}_*(a)_{\underline{x}} = D(a_{q, \underline{x}})_{I_N, \underline{x}}$

which is equipped w a projection $D(\omega_{G, \underline{x}}) \rightarrow \text{Whit}_*(G)_{\underline{x}}$.

However, $\text{Whit}^!(G)_x \xleftarrow[\sim]{\exists \theta} \text{Whit}_*(G)_x$

Structure of $\text{Whit}^1(G)_\mathbb{Z}$

Recall Naïve geometric Satake

$$D(H_{g, \underline{x}}^{\text{loc}}) = \text{Sph}_{g, \underline{x}} \quad (\text{technically should renormalize})$$
$$D(\omega_{h, \underline{x}})^{(L^+h)_{\underline{x}}} = D((L^+h)_{\underline{x}} \setminus (Lh)_{\underline{x}} \setminus (L^+h)_{\underline{x}})$$

Naive Satake functor is a monoidal functor $\text{Rep}(\check{G})_{\underline{x}} \rightarrow \text{Sph}_{G, \underline{x}}$

eg. V^{\dagger} h.w. $\lambda \mapsto I_C \overline{\alpha_{\lambda}^{\dagger}}$, $\overline{\alpha_{\lambda}^{\dagger}}$ = closure of $L^{\dagger}G \cdot t^{\dagger}$


Whenever, $\text{Sph}_{\mathfrak{g}, \underline{x}} \leadsto e \rightsquigarrow \text{Rep}(\check{\mathfrak{g}})_{\underline{x}} \leadsto e$

In particular, $\text{Rep}(\check{G})_{\underline{x}} \leadsto \text{Whit}^!(\check{h})_{\underline{x}} = (\text{Whit}^!(\text{Ig}_{\underline{x}}))^{\text{I}^+ \check{h}}_{\underline{x}}$

$$\text{Vac}_{\text{whit}} \in \text{Whit}^!(G)_{\underline{x}}, \quad \omega_{N, \underline{x}} \subset \omega_{G, \underline{x}}$$

$$\text{Vac}_{\text{whit}} = \text{*}-\text{extend } \omega_{\omega_{N, \underline{x}}} \otimes \chi^!(\exp) \text{ to } \text{Whit}^!(G)_{\underline{x}}.$$

Rmk. Extension is clean, $! \Rightarrow *$.

$$\text{Rep}(\check{G})_{\underline{x}} \xrightarrow{\text{Sat}^{\text{nv}}} \text{Sph}_{G, \underline{x}} \xrightarrow{* \text{Vac}_{\text{whit}}} \text{Whit}^!(G)_{\underline{x}}$$


Thm (Frenkel - Gaitsgory - Wiltonen, Raskin)

Composition is an equiv., inverse is called CS_G .

$$\text{Coet}_{\underline{x}}: \omega_{G, \underline{x}} \xrightarrow{\pi_{\underline{x}}} \text{Bun}_G$$

$$\sim D(\text{Bun}_G) \xrightarrow{\pi_{\underline{x}}^!} D(\omega_{G, \underline{x}}) \xrightarrow{\text{Av}_*} \text{Whit}^!(G)_{\underline{x}}$$

composition is $\text{coet}_{\underline{x}}$.

unitar factorization cat.

The functor $\text{coet}_{\underline{x}}$ admits a left adjoint $\text{Poinc}^!, \underline{x}$.

$$D(\omega_{G, \underline{x}}) \xrightarrow[\pi_!]{\quad} D(\text{Bun}_G)$$

partially defined, but defined on $\text{Whit}^!(G)_{\underline{x}}$.

Obs 1 If $\pi_!(F)$ is defined, then so is $\pi_!(S \star F) = S \star \pi_!(F)$, $S \in \text{Sph}_G$

Obs 2 $\pi_!(\text{Vac}_{\text{whit}})$ is defined

Tutorial 2.2 (Laurie Dhillon) Kac-Moody modules

basic idea: to construct objects of $D(\text{Ban}_h)$ there are 3 basic moves:

- ① Eisenstein series ② Poincaré series ③ KM localization

classical picture: $H = \{x+iy : x, y \in \mathbb{R}, y > 0\}$ $\frac{dx dy}{y^2}$

beautiful differential operator $\Delta y^2 (\partial_x^2 + \partial_y^2)$

Δ plays well w/ modular functions.

$$\begin{array}{ccc} \Delta E_s & = & s(1-s) E_s \\ \uparrow & & \\ \text{Eisenstein series} & & \Delta = s(1-s) \end{array}$$

\Rightarrow bounds microsupport of special functions (nilpotent)

$$G = \text{SL}_2(\mathbb{R})$$

$$H/\Gamma = \text{SO}_2(\mathbb{R}) \backslash \text{SL}_2(\mathbb{R}) / \Gamma \quad , \quad \Gamma = \text{SL}_2(\mathbb{Z})$$

$$\text{Diff}(G)^{G \times G} \simeq Z(g) = \mathbb{R} \left[\frac{1}{2} h^2 + e f + f e \right]$$

$$\Delta f = \lambda f$$

$$(\Delta - \lambda) f = 0 \quad \text{on symbols: } \text{symbol}(\Delta) = 0$$

$$\begin{array}{ccccc} T^*G & \simeq & G \times g^* & \longleftarrow & G \times N \\ \downarrow \mu & & \downarrow \mu & & \downarrow \\ g^* & = & g^* & \longleftarrow & N \\ & & \downarrow & & \downarrow \\ & & g^* // G & \longleftarrow & 0 \end{array}$$

$$x \in X$$



$$k_x \approx \mathbb{C}((t))$$

$$\mathcal{O}_x \approx \mathbb{C}[[t]]$$

$$SL_2(\mathbb{R})$$

$$\downarrow$$

$$\mathbb{H}/\Gamma$$

$$h(k_x)$$

$$\downarrow$$

$$\text{Bun}_h = \pi_1 h(\mathcal{O}_x) \backslash \pi_1 h(k_x) / h(F)$$

\uparrow
func. field of X

Def $g(k_x) := \text{Lie}(I_x h)$

\simeq

$$\mathfrak{g} \otimes \mathbb{C}((t))$$

pointwise Lie bracket

Want: $\mathbb{Z}(U(g(k_x)))$ should be BIG $\approx LA^{rk h}$

example

h f.d. red. gp

$$\mathbb{Z}(g) \simeq \text{Fun}(t^* // w)$$

$$\simeq \text{Fun}(t^\vee // w)$$

$$\simeq \text{Fun}(\check{\mathfrak{o}} // \check{a})$$

ex $h = T$ torus, $t(k_x)$ still commutative, so

$$U(t(k_x)) \simeq \text{Sym}(t(k_x)) \simeq \text{Fun}((t(k_x))^*)$$

$$\simeq \text{Fun}(t^\vee \otimes \Omega^1)$$

$$\simeq \text{Fun}(\text{Conn}_T^\vee(\mathcal{O}))$$

t^n dual to $t^{-n-1} dt$

$$\begin{array}{ccc} k_x \otimes \Omega^1 & & \\ \downarrow w & & \\ \mathbb{C} & \int dw & \end{array}$$

G_m itself

$\mathbb{Z} G_m$

$$\mathbb{Z} \simeq \mathbb{C}[t]$$

\uparrow
 $t \partial_t$

$$\mathbb{Z} \simeq \mathbb{C}[t_i : i \in \mathbb{Z}]$$

ex SL_2

$$\mathbb{Z}(SL_2) = \mathbb{C}[\Delta]$$

$$\mathbb{Z}(\mathbb{Z}_2(K_X)) \simeq \mathbb{C}[\Delta_n : n \in \mathbb{Z}]$$

$$\Delta = \frac{1}{2} h^2 + e f + f e$$

$$x_n = x \otimes t^n$$

$$\Delta_n = \sum_{i+j=n} \frac{1}{2} h_i h_j + e_i f_j + f_j e_i$$

$$= \sum_{i+j=n} \frac{1}{2} : h_i h_j : + : e_i f_j : + : f_j e_i :$$

$$: x_n y_m : = \begin{cases} x_n y_m & n < 0 \\ y_m x_n & n \geq 0 \end{cases}$$

next: which modules to use?

$$\begin{array}{c} K \backslash SL_2(\mathbb{R}) / \Gamma \\ \parallel \\ SO_2(\mathbb{R}) \\ \text{max. cpt} \end{array}$$

$$G(\mathcal{O}_X) \backslash G(K_X) / \overline{\Gamma}$$

Def $KL := (g(K_X), G(\mathcal{O}_X))$ -modules.

modules for $g(K_X)$ on which $g(\mathcal{O}_X)$ is integrated.

Examples

$$V \hookrightarrow \mathfrak{h}(\mathcal{O}_X)$$

$$\text{ind}_{\mathfrak{g}(\mathcal{O}_X)}^{\mathfrak{g}(K_X)} V \in \text{KL}$$

V f.d., ~~no~~ ind system of compact generators

$\lambda \in \Lambda^+$, V^λ irred. \mathfrak{h} -module

$$\begin{array}{ccc} & \mathfrak{h}(\mathcal{O}_X) & \\ \swarrow & & \searrow \\ \mathfrak{h} & & \mathfrak{h}(K_X) \end{array} \quad \begin{array}{c} V^\lambda := \text{ind}_{\mathfrak{g}(\mathcal{O}_X)}^{\mathfrak{g}(K_X)} V \\ \uparrow \\ \text{Weyl module} \end{array}$$

V^0 called vacuum rep'n

Example $\mathfrak{h}_m \hookrightarrow \mathfrak{h}$ 1-dim v.s.

$$\leadsto \text{diff } t \partial_t = n \cdot \text{id} \in \mathbb{Z}$$

$t \partial_t \sim \lambda \in \mathbb{C}$ defines a rep of Lie algebra, doesn't integrate unless $\lambda \in \mathbb{Z}$

Lemma $V \in \mathfrak{g}(K_X)\text{-mod}$ \mathcal{D} lifts to KL iff

① $X_n, n > 0$ act locally nilpotently

② eg. \mathfrak{h} semisimple s.c., $\dim U(g)v$ finite, $\forall v \in V$.

\mathfrak{h} torus \rightarrow int. eigenvalues from example.

Beautiful fact. $V, W \geq$ f.d. $\mathfrak{h}(\mathcal{O}_X)$ -modules

$$\textcircled{1} \text{ Hom}(V, W) \cong \text{Hom}(W^*, V^*)$$

$$\textcircled{2} \text{ Hom}(\text{ind } V, \text{ind } W) \cong \text{Hom}(\text{ind}(W^*), \text{ind}(V^*))$$

comes from a self duality $KL \cong KL^\vee$.

$$KL \otimes KL \longrightarrow \text{Vect}$$

$$M \boxtimes N \longmapsto C^{\frac{\infty}{2}+}(g(K_x), g(\mathcal{O}_x), M \otimes N) \leftarrow \begin{array}{l} \text{relative semi-infinite} \\ \text{cohomology} \end{array}$$

$$C^{\frac{\infty}{2}+}(g(K_x), g(\mathcal{O}_x), M \otimes N)$$

informal def'n

① group cohomology along $g(\mathcal{O}_x)$

② Lie alg. cohomology along $g(K_x)/g(\mathcal{O}_x)$.

\simeq Hamiltonian
reduction

w.r.t. $g(\mathcal{O}_x)$



3. DS reduction

idea

$$\mathcal{Z}(g) \hookrightarrow \mathcal{U}(g)$$

subalgebra but also "quotient".

(Hamiltonian reduction)

ex

$$\mathfrak{g} = \mathfrak{gl}_n$$


$$\begin{array}{c} \mathfrak{g}^* \\ \downarrow S \\ \mathfrak{g}^*/\mathfrak{h} \end{array}$$

$$S = \begin{pmatrix} * & & & \\ & * & & \\ & & * & \\ & & & \ddots \\ & & & & * \end{pmatrix}$$

f principal \mathfrak{sl}_2

$$\begin{pmatrix} * & & & \\ & * & & \\ & & \ddots & \\ & & & * \end{pmatrix} \longrightarrow \mathfrak{f} + \mathfrak{b} \longrightarrow \mathfrak{f} + \mathfrak{b}/N$$

definition



def'n DS reduction is the functor

$$C^{\frac{10}{2}+*}(n(k_x), - \otimes C_\psi) : \mathfrak{g}(k_x)\text{-mod} \longrightarrow \text{Vect}$$

\uparrow
 derivative of
 char from
 Andreus' talk

Lecture 3 (Dima Arinkin)

$$\text{Whit}(\omega_{G, \underline{x}}) \quad \underline{x} \in |\text{Ran}$$

$$\text{Poinc}_{G, \underline{x}} \downarrow \uparrow \text{coet}^*$$

$$\text{Dmod}(\text{Bun}_G)$$

$$\text{KL}(G)_{\underline{x}}$$

$$\text{Loc}_{G, \underline{x}} \downarrow \uparrow \Gamma_{G, \underline{x}}$$

$$\text{Dmod}(\text{Bun}_G)$$

1) $\text{KL}(G)_{\underline{x}}$ - was defined

2) $\text{KL}(G)_{\underline{x}}$ is self dual \leftarrow will be recapped

3) $C^{\frac{10}{2}}(L\mathfrak{g}) : \text{KL}(G)_{\underline{x}} \longrightarrow \text{Vect}$

$$C^{\frac{10}{2}}(L\mathfrak{n}) : \text{KL}(N)_{\underline{x}} \longrightarrow \text{Vect}$$

$$C^{\frac{10}{2}}(L\mathfrak{n}, x) : \text{KL}(N)_{\underline{x}} \longrightarrow \text{Vect}$$

G-oper

G-bundles w connection on a curve X, G reductive gp
 \cup
 B - Borel

(Beilinson-Drinfeld)

"Def" Oper = $(F_B : B\text{-bundle on } X, \nabla\text{-connection on } F_G = G^B \times F_B)$

discrepancy between F_B and ∇ sits in the "next term".

Details

$F_B \rightsquigarrow \underline{\text{Conn}}(F_B)$ - torsors over $(b)_{F_B} \otimes \Omega$


$F_G \rightsquigarrow \underline{\text{Conn}}(F_G)$ - torsor over $(g)_{F_B} \otimes \Omega$

$\underline{\text{Conn}}(F_B) \rightarrow \underline{\text{Conn}}(F_G) \xrightarrow{\text{discrepancy}} (g/b)_{F_B} \otimes \Omega$

$$\frac{d}{dx} + \begin{pmatrix} * & * & * \\ * & * & * \\ * & * & * \end{pmatrix} \quad \bigoplus_{\alpha} (g-\alpha)_{F_B} \otimes \Omega = (g-1/b)_{F_B} \otimes \Omega$$

$$S(\nabla) + (g-1/b)_{F_B} \otimes \Omega$$

and each component in $(g-\alpha)_{F_B} \otimes \Omega$ is nowhere vanishing.



$$\begin{pmatrix} * & * & * \\ * & * & * \\ * & * & * \end{pmatrix} \begin{pmatrix} * & * & * \\ * & * & * \\ * & * & * \end{pmatrix} \begin{pmatrix} 0 & * & * \\ 0 & 0 & * \\ 0 & 0 & 0 \end{pmatrix}$$

$$g-1 > b > n$$

$$1 \quad [b, b]$$

$$\{x+g: [x, n] \subset b\}$$

$$g-1/b = \bigoplus_{\alpha \text{ simple}} g-\alpha$$

Variation make F_B a point of Bun_N^Ω .

$$\begin{matrix} \uparrow \\ Bun_B \times \{ \check{P}(\Omega) \} \\ \uparrow \\ Bun_T \end{matrix}$$

Example $\mathcal{E} = \text{rk } 3 \text{ local system}$

$$\mathcal{E}_0 \subset \mathcal{E}_1 \subset \mathcal{E}_2 \subset \mathcal{E}_3$$

$$\mathcal{E}_1/\mathcal{E}_0 = \Omega, \quad \mathcal{E}_2/\mathcal{E}_1 = 0, \quad \mathcal{E}_3/\mathcal{E}_2 = \Omega^{-1}$$

$$\nabla: \mathcal{E} \rightarrow \mathcal{E} \otimes \Omega$$

$$\nabla(\mathcal{E}_i) \subset \mathcal{E}_{i+1} \otimes \Omega \hookrightarrow (\mathcal{E}_i/\mathcal{E}_{i-1}) \rightarrow (\mathcal{E}_{i+1}/\mathcal{E}_i) \otimes \Omega$$

is not just nonzero, but the given iso.

Informally:


Oper ① $\frac{d}{dt} + \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix} dt$

rat'l canonical form for connections

② Opers are t , connections

n -th order ODEs --- Systems of n


1st order ODEs.



Global opers \rightsquigarrow D-modules on Bun_A^\sim

local opers \rightsquigarrow reps of $K-M$

Ideally on Ran . Now at a single $x \in X$.



Several spaces

$\mathcal{O}_P^{\text{reg}}(D_x)$

$$\frac{d}{dx} + \begin{pmatrix} a_1(x) & a_2(x) & a_3(x) \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, a_1, a_2, a_3 \in \mathbb{C}[[x]]$$

space is \mathbb{A}^∞
scheme

$$\left[\frac{d^3}{dx^3} \pm a_1(x) \frac{d^2}{dx^2} \pm a_2(x) \frac{d}{dx} \pm a_3(x) \right] y(x) = 0$$

(extends to \mathfrak{g} , consider centralizer
of $b \in \mathfrak{g}$)

$\mathcal{O}_P^{\text{mer}}(D_x)$

opers on punctured
disk

$$\left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right)$$

$$a_i \in \mathbb{C}((x))$$

$$\bigcup_{n \rightarrow \infty} x^{-n} \mathbb{C}[[x]]$$

Ind-scheme:

$$\bigcup_n \mathcal{O}_P^{\text{mer}, \leq n} \cong \mathbb{A}^\infty$$

$\mathcal{O}_P^{\text{mon-free}}(D_x)$

monodromy-free
ii)

$a_1(x), a_2(x), a_3(x)$ s.t. there are 3 linearly
indep solutions in $\mathbb{C}[[x]]$.

Ex ① Formally
smooth

② \bigcup closed
subschemes

$Y_n \times \mathbb{A}^\infty$
finite type

$\mathcal{O}_P^{\text{mon}}(D_x) \times \text{LocSys}(D_x)$
 $\text{LocSys}(D_x^\circ)$

What kind of coherent sheaves live on such spaces $(Y \times \mathbb{A}^\infty)$

We need $\text{IndCoh}^*(Y \times \mathbb{A}^\infty)$

ii

$\lim_{\rightarrow} \text{IndCoh}(Y \times \mathbb{A}^n)$

under
* pullbacks

$$Y \times \mathbb{A}^\infty$$

$$\downarrow$$

$$Y \times \mathbb{A}^n$$

$$\downarrow p$$

$$Y \times \mathbb{A}^m$$

Tutorial 2.3 (Ken Lin) Introduction to factorization

A factorisation space is a prestack $\mathcal{Y}_{\text{Ran}} \rightarrow \text{Ran}$ equipped with isoms

$$\mathcal{Y}_{x \sqcup x'} \simeq \mathcal{Y}_x \times \mathcal{Y}_{x'} \quad \text{for } x, x' \in \text{Ran} \text{ disjoint.}$$

$$[\mathcal{Y}_{x_1 \sqcup \dots \sqcup x_n} \simeq \mathcal{Y}_{x_1} \times \dots \times \mathcal{Y}_{x_n}]$$

A factorisation cat. is a sheaf of cat. \mathcal{C}_{Ran} on Ran w

$$\mathcal{C}_{x \sqcup x'} \simeq \mathcal{C}_x \otimes \mathcal{C}_{x'}.$$

eg. $D(\mathcal{Y}_{\text{Ran}}) = D(\mathcal{Y})_{\text{Ran}}$ for \mathcal{Y} a fact space

$$\text{IndCoh}^*(\mathcal{Y}_{\text{Ran}}) = \text{IndCoh}^*(\mathcal{Y})_{\text{Ran}}$$

for $\mathcal{Y}_{\text{Ran}} = \text{Ran}$, this cat is denoted Vect_{Ran} .

$$\text{Ran} = \text{colim } X_{dR}^I$$

sheaf of cats on $(X_{dR})^I$ is a module cat. over $D(X^I)$.

A factorisation alg. (in a fact. cat. \mathcal{C}_{Ran}) is a D -module \mathcal{A}_{Ran} on Ran space (global section of \mathcal{C}_{Ran}) w $\mathcal{A}_{x \sqcup x'} \simeq \mathcal{A}_x \otimes \mathcal{A}_{x'}$ (in $\mathcal{C}_{x \sqcup x'} \simeq \mathcal{C}_x \otimes \mathcal{C}_{x'}$).

eg. w_{Ran} is a factorisation algebra.

A functor $F_{\text{Ran}} : \mathcal{C}_{\text{Ran}} \rightarrow D_{\text{Ran}}$ is a factorization functor if

$$\mathcal{O}_{X \sqcup X'} \xrightarrow{\sim} \mathcal{O}_X \otimes \mathcal{O}_{X'}$$


$$\begin{array}{ccc} \mathcal{F}_{X \sqcup X'} & \downarrow & \mathcal{F}_X \otimes \mathcal{F}_{X'} \\ \mathcal{D}_{X \sqcup X'} & \xrightarrow{\sim} & \mathcal{D}_X \otimes \mathcal{D}_{X'} \end{array} \quad \text{commutes.}$$

$$\mathcal{D}_{X \sqcup X'} \xrightarrow{\sim} \mathcal{D}_X \otimes \mathcal{D}_{X'}$$

Prop A fact. functor sends fact. algebras to fact. algebras.

—————
Σ

Let x be an S -point of Ran



$\Gamma_x = \text{union of graphs}$

$\mathcal{D}_x = \text{formal completion at } \Gamma_x$
(affine)

$$\mathcal{D}_x^0 = \mathcal{D}_x \setminus \Gamma_x.$$

eg. Let $\mathcal{Y} \rightarrow X$ be a prestack, equipped w/ a connection (i.e. descent to X_{dR})

$$(\mathcal{L}_{\mathcal{D}}^+ \mathcal{Y})_{\text{Ran}} = \{x \in \text{Ran}, \text{ horizontal section of } \mathcal{Y} \text{ over } \mathcal{D}_x\}$$

$$\downarrow$$

Ran

For $\mathcal{Y} = BG \times X$, this produces $(BG)_{\text{Ran}}$.

$\text{Ind Coh}_* (\mathcal{L}_{\mathcal{D}}^+ \mathcal{Y})_{\text{Ran}}$ is a fact. cat.

$\mathcal{O}_{\mathcal{L}_{\mathcal{D}}^+ \mathcal{Y}, \text{Ran}}$ is a fact. alg in the fact. cat.

Rmk if x_1, \dots, x_n are k -points of X ,

$$(L_{\nabla}^+ Y)_{x_1 \sqcup \dots \sqcup x_n} = \prod_{i=1}^n Y_{x_i}.$$

Assume Y is affine over X

$$(L_{\nabla}^+ Y)_{\text{Ran}} = \left\{ x \in \text{Ran}, \text{ horizontal section of } Y \text{ over } \bar{D}_x \right\}$$

\downarrow
Ran

There is a constant map $z: (L_{\nabla}^+ Y)_{\text{Ran}} \rightarrow (L_{\nabla}^+ Y)_{\text{Ran}}$

So $z_*: \text{IndCoh}_* (L_{\nabla}^+ Y)_{\text{Ran}} \rightarrow \text{IndCoh}_* (L_{\nabla}^+ Y)_{\text{Ran}}$ is a fact. functor.

Hence $z_* \mathcal{O}_{L_{\nabla}^+ Y, \text{Ran}}$ is a fact. alg. (in the latter cat.)

In general, $(L_{\nabla}^+ Y)_x$ is an ind-scheme of infinite type.

Let x be a point of X .

$$\text{Ran}_x = \{ y \in \text{Ran} : x \in y \}$$

A factorization module for a fact. alg A_{Ran} (in a fact. cat. \mathcal{C}_{Ran})

is a D-module (a global section of $\mathcal{C}_{\text{Ran}_x}$) $M_{\text{Ran}_x} \rightsquigarrow$

$$M_{y \sqcup y'} \simeq M_y \otimes A_{y'} \quad \text{for } y \in \text{Ran}_x, \quad y' \in \text{Ran} \text{ disjoint}$$

$$M_{x \sqcup y_1 \sqcup \dots \sqcup y_n} \simeq M_x \otimes A_{y_1} \otimes \dots \otimes A_{y_n}$$

eg. A_{Ran_x} is a fact. module for A_{Ran} .

Claim

$$\begin{array}{ccc}
 & \mathcal{O}_{L^+_{\nabla} y, \text{Ran} - \text{mod } x}^{\text{fact}} & \\
 & \downarrow \text{fiber at } x & \\
 \text{Ind Coh}_* (L_{\nabla} y)_x & \xrightarrow{\quad} & \text{Vect}
 \end{array}
 \quad \begin{array}{l}
 \text{(conflating } \mathcal{O}_{L^+_{\nabla} y, \text{Ran}} \\
 \leadsto \Gamma(L^+_{\nabla} y, \mathcal{O}_{L^+_{\nabla} y}(\text{Ran}))
 \end{array}$$

Suffices to define a functor

$$\text{ins. var} : \text{Ind Coh}_* (L_{\nabla} y)_x \longrightarrow \text{Ind Coh}_* (L_{\nabla} y)_{\text{Ran}_x}$$

which fiberwise performs $F \mapsto F \boxtimes \mathcal{O}_{L^+_{\nabla} y, y_1} \boxtimes \dots \boxtimes \mathcal{O}_{L^+_{\nabla} y, y_n}$

Key input: if $y \in \text{Ran}_x$, we have

$$\begin{array}{ccc}
 \tilde{D}_x & \hookrightarrow D_y \setminus x \hookleftarrow \tilde{D}_y \\
 & \text{Sect}_{\nabla}(D_y \setminus x, y) \\
 \leadsto (L_{\nabla} y)_x & \swarrow \quad \searrow & (L_{\nabla} y)_y
 \end{array}$$

$$\leadsto \text{Ind Coh}_* (L_{\nabla} y)_x \longrightarrow \text{Ind Coh}_* (L_{\nabla} y)_y$$

then allow y to vary.

At $x \sqcup y_1 \sqcup \dots \sqcup y_n$

$$\tilde{D}_x \hookrightarrow \tilde{D}_x \sqcup D_{y_1} \sqcup \dots \sqcup D_{y_n} \hookleftarrow \tilde{D}_x \sqcup \tilde{D}_{y_1} \sqcup \dots \sqcup \tilde{D}_{y_n}$$

$$(L_{\nabla} y)_x \times (L^+_{\nabla} y)_{y_1} \times \dots \times (L^+_{\nabla} y)_{y_n}$$

$$\swarrow \\ (L_{\nabla} y)_x$$

$$\searrow \\ (L_{\nabla} y)_x \times (L_{\nabla} y)_{y_1} \times \dots \times (L_{\nabla} y)_{y_n}$$

Thm the functor $\text{Ind Coh}_*(\mathbb{A}^1 Y)_x \longrightarrow \mathcal{O}_{\mathbb{A}^1 Y - \text{mod}}^{\text{fact}}_x$

is an "equivalence" of categories on $> -\infty$ part.

A unital str. on a fact. cat. \mathcal{C}_{Ran} is:

• a fact. alg. unit e_{Ran} (called factorization unit)

• and the ability to assemble

$$\mathcal{C}_x \longrightarrow \mathcal{C}_x \boxtimes \mathcal{C}_{y_1} \boxtimes \cdots \boxtimes \mathcal{C}_{y_n}, \quad F \mapsto F \boxtimes \text{unit } e_{y_1} \boxtimes \cdots \boxtimes \text{unit } e_{y_n}$$

into a functor $\text{insrac}: \mathcal{C}_x \longrightarrow \mathcal{C}_{\text{Ran}_x}$,

eg. $\hat{\mathcal{G}}_K - \text{mod}$ Kai-Moody modules

is a factorization cat, w/ unit $\text{Var}(G)_K$

$\hat{\mathcal{G}}_K - \text{mod} \xrightarrow{\text{oblv}} \text{Vect}$ is a fact. functor

$\leadsto \forall g, k = \text{oblv}(\text{Var}(G)_K)$ is a fact. alg.

$$\begin{array}{ccc} (*) & \xrightarrow{\quad} & \mathcal{G}_{g,k} - \text{mod}_x^{\text{fact}} \\ & \searrow & \downarrow \\ \hat{\mathcal{G}}_K - \text{mod}_x & \xrightarrow{\text{oblv}} & \text{Vect} \end{array}$$

Thm (*) is an equiv. on $> -\infty$.

Lecture 4 (Yifei Zhao)

Summary \exists fact. alg. def'n

pretty wild

Pretty complex notion

carries some weight
Page 44

Factorization algs \longleftrightarrow Vertex algebras

\exists notions of unital fact. alg

modules (at $x \in X$)

unital modules

Local theory of fact. algs:

1) y affine scheme $/X \hookrightarrow \Delta$

\leadsto Fact. alg \mathcal{O}_y

$\mathcal{O}_y\text{-mod}_x^{\text{fact}} \approx \left. \begin{array}{l} \text{coherent} \\ \text{sheaves} \end{array} \right\} \text{ on } L_{\Delta} y_x \Bigg\} \begin{array}{l} \text{space of horizontal sections} \\ \text{on } \tilde{D}_x \end{array}$

corrections: work w/ ev. coconn. modules,

Ind Coh^* on RHS.

2) $\exists \text{ Vac}(g)_k \leftarrow \text{"Kac-Moody factorization algebra"}$

fiber of $\text{Vac}(g)_k$ at $x \in X$ is $\mathcal{V}_k^0 = \mathcal{V}_k \Bigg\} \begin{array}{l} \text{vacuum rep'n of } \hat{g}_k \\ \text{ind } \hat{g}_k \oplus k \text{ (triv)} \end{array}$

$\text{Vac}(g)\text{-mod}_x^{\text{fact}} \approx g(t)\text{-mod}$

\uparrow

really on coconn. objects

w/ k : get modules for central ext'n



Goal. Prove

Thm There is a canonical equiv. of unital factorization categories

$$KL(\mathfrak{g})_{\text{crit}} \simeq \text{IndCoh}_* (\mathcal{O}_P^{\text{mon-free}}_{\check{\mathfrak{h}}}) \quad [\text{FLE}_{\text{crit}}]$$

$$KL(\mathfrak{g})_{\text{crit}} := (\hat{\mathfrak{g}}\text{-mod}_{\text{crit}})^{L^+\mathfrak{g}}$$

level: κ is a \mathfrak{g} -inv't sym. bilinear form $\mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathbb{C}$.

$$0 \rightarrow \mathbb{C} \rightarrow \mathfrak{g}^{\kappa} \rightarrow \mathfrak{g}(\kappa_{\mathbb{C}}) \rightarrow 0 \quad \text{defined by the cocycle}$$

$$\xi \otimes f, \xi' \otimes f' \mapsto \kappa(\xi, \xi') \text{Res}(\text{id}_f) f'$$

$$\text{crit} = -\frac{1}{2} \text{Killing}$$

← uses $\omega^{1/2}$
Pfaffian

Rank $\text{Loc} : KL(\mathfrak{g})_{\text{crit}} \rightarrow D_{\text{mod crit}}(\text{Bun}_{\mathfrak{g}}) \simeq D_{\text{mod}}(\text{Bun}_{\mathfrak{g}})$

$$\text{Sat}^{\text{nu}} : \text{Rep}(\check{\mathfrak{h}})^{\vee} \simeq_{\text{Sym.}} D_{\text{mod crit}}(\mathcal{H}^{\text{loc}})^{\vee} \simeq D_{\text{mod}}(\mathcal{H}^{\text{loc}})^{\vee}, \text{ modified conn.}$$

$$\begin{array}{ccccc} & & \xrightarrow{2^+} & & \\ & \nearrow^{2^+, \text{mon-free}} & & \searrow & \\ \mathcal{O}_P^{\text{reg}}_{\check{\mathfrak{h}}} & \longrightarrow & \mathcal{O}_P^{\text{mon-free}}_{\check{\mathfrak{h}}} & \longrightarrow & \mathcal{O}_P^{\text{mer}}_{\check{\mathfrak{h}}} \\ \downarrow & & & & \downarrow \\ \text{LS}^{\text{reg}}_{\check{\mathfrak{h}}} & \longrightarrow & \text{LS}^{\text{mer}}_{\check{\mathfrak{h}}} & & \end{array}$$

units match under FLE_{crit}: $\text{Var}(\mathfrak{g})_{\text{crit}} \mapsto \left(2^+, \text{mon-free}\right)_*^{\text{IndCoh}} \left(\mathcal{O}_{\mathcal{O}_P^{\text{reg}}_{\check{\mathfrak{h}}}}\right)$

At $x \in X$,

$$H^0 \text{End}(\text{Vac}(h)_{\text{out}}) \simeq H^0 \text{End}(z_*^{+, \text{mf}} \mathcal{O}) \simeq \Gamma(\mathcal{O}_{P_{\check{h}, x}^{\text{reg}}}, \mathcal{O})$$

$$\text{ind}_{\mathcal{G}(\mathcal{O}_x)}^{\hat{\mathcal{G}}_{\text{out}}(k)^{\mathcal{G}(\mathcal{O}_x)}} \quad (\text{Feigin - Frenkel at } x \in X)$$

Step 1 Establish FF.

$H^0 \text{End}(A^{(n)})$
 A classical
 fact. alg $\left\{ \begin{array}{l} \mathcal{G}_{\text{out}} := \text{centre of the fact. algebra } \mathbb{K} \mathcal{G}_{\text{out}} \quad (:= \text{image of } \text{Vac}(h)_{\text{out}} \\ \text{under the forgetful functor}) \end{array} \right.$

Construct an isom. of \mathcal{D}_X -algebras $\mathcal{G}_{\text{out}} \simeq \mathcal{O}_{P_{\check{h}}^{\vee}}$

$$\mathcal{O}_{P_{\check{h}}^{\vee}} \xrightarrow{\text{affine}} X_{\text{dR}}$$

$$L_{\mathcal{D}}^+(\mathcal{O}_{P_{\check{h}}^{\vee}}) \simeq \mathcal{O}_{P_{\check{h}}^{\vee}}^{\text{reg}}$$

$$L_{\mathcal{D}}(\mathcal{O}_{P_{\check{h}}^{\vee}}) \simeq \mathcal{O}_{P_{\check{h}}^{\vee}}^{\text{mer}}$$

Step 2 Construct a functor (enhanced Drinfeld-Sokolov)

$$\text{KL}(h)_{\text{out}} \longrightarrow \text{IndCoh}_*(\mathcal{O}_{P_{\check{h}}^{\vee}}^{\text{mer}}) \quad (\mathcal{D}S^{\text{enh}})$$

Step 3. $(\mathcal{D}S^{\text{enh}})$ factors through an equiv. $\text{KL}(h)_{\text{out}} \xrightarrow{\sim} \text{FLE}_{\text{out}} \text{IndCoh}_*(\mathcal{O}_{P_{\check{h}}^{\vee}}^{\text{mer-free}})$

$$\begin{array}{ccc} & & \downarrow z_*^+ \\ \searrow \mathcal{D}S^{\text{enh}} & & \text{IndCoh}_*(\mathcal{O}_{P_{\check{h}}^{\vee}}^{\text{mer}}) \end{array}$$

Step 1 Construction of (FF)

Reduction to the universal case

$$\hat{D}_0 := \text{Spt } k[[t]]$$

group $\text{Aut} :=$ automorphism group of \hat{D}_0
ind-scheme

group $\text{Aut}^+ :=$... preserving the base point
scheme

$$\text{Aut} / \text{Aut}^+ \simeq \hat{D}_0$$

$$\begin{array}{c} \hat{X}_{dR} := \{x \in X_{dR} : \hat{D}_x \simeq \hat{D}_0\} \\ \downarrow \text{Aut} \\ X_{dR} \end{array}$$

pullback to X is induced from an Aut^+ -torsor.

$$\mathfrak{Z}_{g, \text{crit}, 0} := \text{ind}_{g[[t]]}^{\hat{g}^{\text{crit}}} (k) \llbracket t \rrbracket \hookrightarrow \text{Aut}$$

$$\text{Op}_{\check{A}, 0}^{\vee} := \text{affine scheme parametrizing } \check{A}\text{-opers on } \hat{D}_0 \hookrightarrow \text{Aut} \quad (\text{actually a double cover of Aut})$$

Suffices to construct an Aut -equiv. isom. $\text{Spec}(\mathfrak{Z}_{g, \text{crit}, 0}) \simeq \text{Op}_{\check{A}, 0}^{\vee}$

Birth of opers

$$\begin{array}{ccc} \text{Spec}(\mathfrak{Z}_{g, \text{crit}, 0}) \longrightarrow \text{Op}_{\check{A}, 0}^{\vee} & \Leftrightarrow & \check{A}\text{-oper on } \text{Spec}(\mathfrak{Z}_{g, \text{crit}, 0}) \times \hat{D}_0 \\ \text{Aut-equiv} & & \text{w Aut-equivariance} \end{array}$$

General paradigm $H^+ \subset H \rightsquigarrow Y$

$$Y \times^H H / H^+ \simeq Y / H^+$$

$$\downarrow \qquad \qquad \downarrow$$

$$Y \times^H H / H \simeq Y / H$$

Suffices to construct

1) Aut^+ -equiv. P_B^\vee and $(P_B^\vee)_T^\vee \xrightarrow{\sim} p(\Omega)$ over $\text{Spec}(\mathbb{Z}_{g, \text{out}}, 0)$

2) upgrade Aut^+ -equiv. on $(P_B^\vee)_\hbar^\vee$ to an Aut -equiv.

The $\check{\zeta}$ -bundle over $\text{Spec}(\mathbb{Z}_{g, \text{out}}, 0)$:

$$\text{Rep } \check{\zeta} \longrightarrow \mathbb{Z}_{g, \text{out}}, 0\text{-mod}$$

$$V \longmapsto \Gamma^{\text{Indcoh}}(\omega_{\check{\zeta}}, \text{Sat}^{\text{nv}}(V))^{\text{GL}(\mathbb{Z})} \quad [\text{Raskin}]$$

Consequence of the construction,

$$\text{Sat}^{\text{nv}}(V) * \text{Vac}(\mathfrak{h})_{\text{out}} \xrightarrow{\sim} \text{ev}^*(V) \otimes_{\mathcal{O}_{P_\hbar^{\text{reg}} \check{\zeta}}} \text{Vac}(\mathfrak{h})_{\text{out}}$$

$$\longrightarrow$$

Step 2

Affine Skryabin theorem (Raskin)

The functor $DS: \hat{\mathfrak{g}}^k\text{-mod} \longrightarrow \text{Vect}$ induces an equiv

$$DS^{\text{enh}}: \text{Whit}_*(\hat{\mathfrak{g}}^k\text{-mod}) \xrightarrow{\sim} (DS(\text{Vac}(\mathfrak{h})_k)\text{-mod}^{\text{fact}})_{\text{ren}}$$

At $x \in X$, $\text{Whit}_* \simeq \text{colim}_{\text{affine Kostant thm}} \text{Whit}^{\leq n}$, $\text{Whit}^{\leq n}$ is gen. by a single obj. W_k^n .
omitted.

At $k = \text{out}$, $DS(\text{Vac}(\mathfrak{h})_{\text{out}}) \simeq \mathbb{Z}_{g, \text{out}}, 0 \xrightarrow{\text{FF}} \mathcal{O}_{P_\hbar^{\text{reg}} \check{\zeta}}$

$$\text{Have } DS^{\text{enh}}: \hat{g}^{\text{unit-mod}} \longrightarrow \text{IndCoh}_*(\mathcal{O}_{P_{\check{h}}}^{\text{mer}})$$

$$\searrow \quad \downarrow$$

$$\mathcal{O}_{P_{\check{h}}}^{\text{reg-mod fact}}$$

The desired functor is

$$KL(h)_{\text{unit}} := (\hat{g}^{\text{unit-mod}})^{L^+h} \longrightarrow \hat{g}^{\text{unit-mod}} \xrightarrow{DS^{\text{enh}}} \text{IndCoh}_*(\mathcal{O}_{P_{\check{h}}}^{\text{mer}}).$$

Bonus Material B (Dennis Gaitsgory)

$$1) \text{Sph}_{\check{h}} \otimes KL \xrightarrow{*} KL$$

$$V \in \text{Rep}(\check{G}),$$

$$\boxed{\text{Sat}^{\text{nv}}(V) \otimes \text{Vac}_{\text{unit}} \simeq \tau^*(V) \otimes \text{Vac}_{\text{unit}}}$$

$$\mathcal{O}_{P_{\check{G}}} \xrightarrow{2} L\mathcal{S}_{\check{h}}^{\text{reg}} \simeq \text{pt}/\check{G}$$

$$2) \text{Vac}_{\text{unit}} \in KL(h)_{\text{unit}} \longrightarrow KM(h)_{\text{unit}}$$

$$DS: KM(h)_{\text{unit}} \longrightarrow \text{Vect}$$

$$\mathcal{O}_{P_{\check{h}}}^{\text{reg}} \xrightarrow[\text{Thm}]{\text{FF} \simeq \mathfrak{z}} DS(\text{Vac}_{\text{unit}})$$

3) DS automatically enhances to a functor

$$DS^{\text{enh}}: KM(h)_{\text{unit}} \longrightarrow DS(\text{Vac}_{\text{unit}})_{\text{-mod fact}}$$

$$\uparrow$$

$$KL(h)_{\text{unit}}^{\text{FLE}}$$

$$\xrightarrow{\sim} \text{IndCoh}^*(\mathcal{O}_{P_{\check{h}}}^{\text{non-free}})$$

$$\uparrow \mathfrak{z}$$

$$\mathfrak{z}_{\text{-mod fact}}$$

$$\xrightarrow[\text{Page 50}]{\text{FF}} \mathcal{O}_{P_{\check{h}}}^{\text{reg-mod fact}} \simeq \text{IndCoh}^*(\mathcal{O}_{P_{\check{h}}}^{\text{mer}})$$

$$L_G \quad C$$

$$\text{Sph}(C) = C^{L^+G}$$

Ex. $C = KM, \text{Sph}(KM) = KL$

$$C = D_{\text{mod}}(L_G), \text{Sph}(C) = D_{\text{mod}}(L_G/L^+G)$$

$$\text{Sph}_G = D_{\text{mod}}(L_G \backslash L_G/L^+G) \leadsto \text{Sph}(C)$$

$$C^{\text{sph-gen}} := D_{\text{mod}}(L_G) \underset{\text{Sph}_G}{\otimes} \text{Sph}(C) \xrightarrow{\quad} C$$

$$\text{Whit}^!(C) \text{ and } \text{Whit}_*(C)$$

Ex $C = D_{\text{mod}}(L_G), \text{Whit}_*(D_{\text{mod}}(L_G)) = \text{Whit}_*(G)$

$$\text{Whit}_*(KM)$$

$$\begin{array}{ccc} \text{DS}^{\text{enh}} : KM & \longrightarrow & \text{IndCoh}^*(\mathcal{O}_{P^1/\check{G}}^{\text{mer}}) \\ & \searrow & \nearrow \overline{\text{DS}^{\text{enh}}} \\ & \text{Whit}_*(KM) & \end{array}$$

Thm (Raskin) $\overline{\text{DS}^{\text{enh}}}$ is an equiv.

$$\text{Whit}(G) \underset{\text{Sph}_G}{\otimes} C^{L^+G} \simeq \text{Whit}_*(C^{\text{sph-gen}}) \hookrightarrow \text{Whit}_*(C)$$

Derived Satake

$$\begin{array}{ccc} \text{Rep}(\check{G}) & \xrightarrow{\text{Sat}^{\text{nv}}} & \text{Sph}_G \\ \downarrow & & \downarrow \text{Sat} \\ \text{Sph}_{\check{G}}^{\text{spec}} := \text{IndCoh}(LS_{\check{G}}^{\text{reg}} \times LS_{\check{G}}^{\text{reg}}) & & \end{array}$$

$$\begin{array}{c} P^1/\check{G} \\ \downarrow \\ P^1/\check{G} \times_{\check{S}_0/\check{G}} P^1/\check{G} \simeq P^1/\check{G} \times_{\check{G}/\check{G}} P^1/\check{G} \end{array}$$

$$\text{Sph}_{\check{A}}^{\text{Spec}} \longrightarrow \text{Sph}_{\check{A}, \text{temp}}^{\text{Spec}}$$

$$\text{IndCoh}(\) \xrightarrow{\Psi} \text{QCoh}(\)$$

$$\text{Sph}_{\check{A}}^{\text{Spec}} \rightsquigarrow \tilde{C} \rightsquigarrow \tilde{C}_{\text{temp}} := \text{Sph}_{\check{A}, \text{temp}}^{\text{Spec}} \otimes_{\text{Sph}_{\check{A}}^{\text{Spec}}} \tilde{C}$$

$$\begin{array}{ccc} \text{QCoh}((\text{LS}_{\check{A}}^{\text{unreg}})^{\vee}) & \xrightarrow{\quad} & \text{QCoh}(\text{LS}_{\check{A}}^{\text{reg}}) \hookrightarrow \text{Sph}_{\check{A}, \text{temp}}^{\text{Spec}} \\ & \downarrow \text{is} & \\ & \text{Rep}(\check{A}) & \end{array}$$

This bimodule defines a Morita equivalence.

$$\text{Whit}^!(\check{A}) \stackrel{\text{CS}}{\simeq} \text{Rep}(\check{A})$$

$$\boxed{\text{Whit}_*(\check{A}) \stackrel{\text{FLE}_\infty}{\simeq} \text{Rep}(\check{A})}$$

$$\text{Sph}_{\check{A}} \xrightarrow[\sim]{\text{Sat}} \text{Sph}_{\check{A}}^{\text{Spec}}$$

$$\text{QCoh}(\text{LS}_{\check{A}}^{\text{reg}}) \otimes_{\text{Sph}_{\check{A}}^{\text{Spec}}} \text{Sph}(C) \stackrel{\textcircled{1}}{\simeq} \text{Whit}_*(C^{\text{sph-gen}})$$

$$\text{QCoh}((\text{LS}_{\check{A}}^{\text{mon}})^{\wedge}_{\text{reg}})$$

$$\text{Cor } \text{Sph}(C)_{\text{temp}} \stackrel{\textcircled{2}}{\simeq} \text{QCoh}(\text{LS}_{\check{A}}^{\text{reg}}) \otimes_{\text{QCoh}((\text{LS}_{\check{A}}^{\text{mon}})^{\wedge}_{\text{reg}})} \text{Whit}_*(C^{\text{sph-gen}})$$

Apply ① to $C = KM$

$$\begin{array}{ccc} \mathcal{Q}coh(LS_{\check{\alpha}}^{reg}) \otimes_{Sph_{\check{\alpha}}^{spec}} KL \simeq \text{Whit}_*(KM^{sph-gen}) & \hookrightarrow & \text{Whit}_*(KM) \\ & \searrow \quad \quad \quad \downarrow & \\ & \text{Ind}coh^*(\mathcal{O}_{P_{\check{\alpha}}}^{mer})_{mf} & \hookrightarrow \text{Ind}coh^*(\mathcal{O}_{P_{\check{\alpha}}}^{mer}) \end{array}$$

$$\mathcal{O}_{P_{\check{\alpha}}}^{mf} = \coprod_{\check{\lambda} \in \Lambda^+} \mathcal{O}_{P_{\check{\alpha}}}^{\check{\lambda}-reg, \sim}$$

$$\forall \check{\lambda} \in KL, \quad DS(W^{\check{\lambda}}) \simeq \mathcal{O}_{P_{\check{\alpha}}}^{\check{\lambda}-reg}$$

$$\underline{\text{Cor.}} \quad KL_{temp} \simeq \mathcal{Q}coh(LS_{\check{\alpha}}^{reg}) \otimes_{\mathcal{Q}coh((LS_{\check{\alpha}}^{mer})_{reg}^{\wedge})} \text{Ind}coh^*(\mathcal{O}_{P_{\check{\alpha}}}^{mer})_{mf}$$

Recall. want $KL \simeq \text{Ind}coh^*(\mathcal{O}_{P_{\check{\alpha}}}^{mf})$

$$\mathcal{O}_{P_{\check{\alpha}}}^{mf} \longrightarrow (\mathcal{O}_{P_{\check{\alpha}}}^{mer})_{mf}^{\wedge}$$

$$\begin{array}{ccc} \downarrow & & \downarrow \\ LS_{\check{\alpha}}^{reg} & \longrightarrow & (LS_{\check{\alpha}}^{mer})_{reg}^{\wedge} \end{array}$$

$$\text{Ind}coh^*(\mathcal{O}_{P_{\check{\alpha}}}^{mf}) \simeq \mathcal{Q}coh(LS_{\check{\alpha}}^{reg}) \otimes_{\mathcal{Q}coh((LS_{\check{\alpha}}^{mer})_{reg}^{\wedge})} \text{Ind}coh^*(\mathcal{O}_{P_{\check{\alpha}}}^{mer})_{mf}$$

Prop. $KL \xrightarrow{\sim} KL_{temp}$

Tutorial 3.1 (Sam Raskin) Kac-Moody localization

Finite dim'l analogue:

H affine alg gp, $K \subset H$ subgp

$$H \curvearrowright \tilde{Y}$$

$$\downarrow$$

$$Y \leftarrow \text{stack / sm. variety}$$

$$Y = \tilde{Y} / K$$

Construction: $\text{Loc}_Y: \underbrace{H\text{-mod } K}_{\substack{K\text{-integrated} \\ H\text{-modules}}} \longrightarrow D\text{-mod}(Y)$

$$H_K^\wedge \text{ reps} \Leftrightarrow H\text{-mod } K.$$

First: $H\text{-mod} \longrightarrow D\text{-mod}(\tilde{Y})$

There is a map $h \xrightarrow{\uparrow \text{Lie alg map}} \text{Vect fields on } \tilde{Y} \quad \text{b/c } H \curvearrowright \tilde{Y}.$

$$U(h) \longrightarrow \Gamma(\tilde{Y}, \text{Diff } \tilde{Y}) = \text{End}(\text{Diff } \tilde{Y})$$

$$\text{Given } M \in H\text{-mod}, \quad \text{Loc}(M) = M \overset{\circlearrowleft}{\otimes}_{U(h)} \text{Diff } \tilde{Y} \in D\text{-mod}(\tilde{Y})$$

$$\text{Loc}(U(h)) = \text{Diff } \tilde{Y}.$$

Correspondence-by:

$$\begin{array}{ccc} & \tilde{Y}/H^\wedge & \\ \swarrow & & \searrow \\ \tilde{Y}_{dR} & & B H^\wedge \end{array}$$

} KdR acts here; quotient by KdR to get HC version

Second: Pass to k -equiv. objects

$$\text{Obtain } \mathfrak{h}\text{-mod } k \xrightarrow{\text{Loc}} D_{\text{mod}}(\tilde{y})^k = D_{\text{mod}}(\tilde{y}/k) = D\text{-mod}(\mathfrak{y})$$

Concretely, $M \in \mathfrak{h}\text{-mod } k$, can see $\text{Loc}(M) \in D_{\text{mod}}(\tilde{y})$

HC str. \rightsquigarrow descent datum

Example $k = H$, $\mathfrak{h}\text{-mod } H = \text{Rep } H \longrightarrow D_{\text{mod}}(\mathfrak{y})$

$$V \longmapsto D(H) \otimes V_{\mathfrak{y}} \quad \begin{array}{c} \tilde{y} \\ \downarrow \\ \mathfrak{y} \longrightarrow BH \end{array}$$

Generalization:

$$V \in \text{Rep } k, \quad \text{ind}_{\text{Lie } k}^{\mathfrak{h}}(V) \in \mathfrak{h}\text{-mod } k$$

$V_{\mathfrak{y}} :=$ pullback of V
a.k.a. assoc. bundle

Loc sends this to D -modules induced from $V_{\mathfrak{y}}$. $\mathfrak{y} \xrightarrow{\tilde{y}} Bk$



$$\tilde{y} = H/H_0 \quad \text{for some } H_0 \subset H \text{ subgroup, } \mathfrak{y} = k \setminus H/H_0$$

fiber of $\text{Loc}(M)$ at $1 \in k \setminus H/H_0$ is $C(\text{Lie } H_0, M) \leftarrow$ derived coinvariants.

Loc has a right adjoint (if \mathfrak{y} is quasi-cpt)

$$\Gamma \simeq \Gamma(\tilde{y}, \pi^!(-))$$

Ex $\Gamma(\delta_{1 \in H/H_0}) = \text{ind}_{H_0}^{\mathfrak{h}}(\text{triv})$



Formalism of Loc

Setup $\mathfrak{y} = \text{Bun}_G$, $x \in X$ (or $x \in \text{Ran}$)

$$\mathfrak{g} = \text{Bun}_G^{\text{loc}, x} = \{ P_G \text{ on } X \setminus x \text{ w/ triv. on } \mathring{D}_x \subset X \setminus x \}$$

origin of
Hecke action

$H = L_x G$ $\text{Bun}_G = \text{Bun}_G^{\text{loc}, x} / L_x^+ G$

∞ -dim'l version of earlier discussion:

$$\text{Loc}: \text{KL}(G) \longrightarrow \text{D-mod}(\text{Bun}_G)$$

$$\text{or } \text{Loc}: \text{KL}(G)_k \longrightarrow \text{D-mod}_k(\text{Bun}_G)$$

Ditto for any $U \subset \text{Bun}_G$ (in practice, U is quasi-cpt)

$$\begin{array}{ccc} \text{Bun}_G^{\text{loc}}(\mathring{D}_x) & & \\ \parallel & \text{Rep}(L_x^+ G) \xrightarrow{\text{ind}} & \text{KL}(G)_k \\ \text{BL}_x^+ G & \downarrow \text{evaluation bundles} & \downarrow \\ \uparrow & \text{Bun}_G \text{ Coh}(\text{Bun}_G) \xrightarrow{\text{ind}} & \text{D-mod}_k(\text{Bun}_G) \\ & \uparrow \text{tensor w/ Diff}_k & \end{array} \quad \left. \begin{array}{l} \text{commutes} \end{array} \right\} \begin{array}{l} \text{Equivariant for action of} \\ \text{Sph}_k(G)_x = \text{D-mod}_k(L^+ G \backslash L^+ G / L^+ G) \end{array}$$

Example $\text{Loc}(\mathbb{V}_k) = \text{Diff}_k$

$\text{ind}(\text{triv}) \quad \text{ind}(0)$

Example $\mathbb{V}_{k,x}^\lambda = \text{ind}_{g \in \mathbb{D}}^{\mathfrak{g}_k} (V^\lambda)$

$\text{Loc}(\mathbb{V}^\lambda) = (\text{twisted}) \text{ D-module induced from } \mathbb{E}_x^\lambda \text{ on } \text{Bun}_G$

$$\begin{array}{ccc} p_x^{\text{univ}} & \longleftarrow & p_x^{\text{univ}} \\ \downarrow & & \downarrow \\ X \times \text{Bun}_G & \xleftarrow{\text{res id}} & \text{Bun}_G \end{array} \quad \text{twist } V^\lambda \text{ by } p_x^{\text{univ}} \text{ to get } \mathbb{E}_x^\lambda$$

Application (BD), $k = \text{out}$ $\hookrightarrow \mathcal{D}_X$ - by functoriality
 $\text{Diff Bun}_k, \text{out} = \text{Loc}(\mathcal{V}_{\text{out}}^X)$
 (choose $x \in X$)

$$\text{Fun}(\mathcal{O}_{\tilde{G}, x}^{\text{res}}) = \mathcal{D}_x \curvearrowright \mathcal{V}_{\text{out}, x}$$

Rmk endos of $\text{Diff} \Leftrightarrow$ global diff'l ops

says: elements of $\mathcal{D}_x \rightsquigarrow$ global diff. ops on Bun_k .

Rmk (Preview) : Action of \mathcal{D}_x factors through an action of $\text{Fun}(\mathcal{O}_{\tilde{G}}^X(X))$.

Further: Birth ofopers Yoga:

$$\text{Sat}^{nv}(V) * \mathcal{V}_{\text{out}, x} \simeq \mathcal{V}_{\mathcal{O}_{\tilde{G}, x}^{\text{res}}} \otimes \mathcal{V}_{\text{out}, x}$$

Deduce, $\text{Sat}_x^{nv}(V) * \text{Diff}_{\text{out}} \simeq \mathcal{V}_{\mathcal{O}_{\tilde{G}, x}^{\text{res}}} \otimes \text{Diff}_{\text{out}}$

Hecke eigenproperty (in simplified form)

Aside: If you wanted an "honest" eigensheaf, believe $\text{Fun}(\mathcal{O}_{\tilde{G}}^X(X))$ act on

Diff_{out} . fix $\sigma \in \mathcal{O}_{\tilde{G}}^X(X)$, quotient Diff_{out} by cov. max'l ideal, result

call F_σ

Fact. F_σ is holonomic (in fact, regular, irred. if G is s.c.)

Tutorial 3.2 (Nick Rozenblyum) Factorization algebras

Recall X smooth ^{conn'd} proper curve

$$\text{Ran}(X) \text{ prestack, } \text{Ran}(X)(S) = \left\{ \begin{array}{l} \text{non empty finite} \\ \text{subsets of } X(S_{\text{red}}) \end{array} \right\}$$

$$\text{Ran}(X) = \text{colim}_{I \in (f_{\text{Surj}})^{\text{op}}} X_{dR}^I$$

$\mathcal{Q}\text{coh}(\text{Ran}(X))$ has some important features

$$1) \quad \pi: \text{Ran}(X) \longrightarrow \text{pt}$$

$\pi^!$ has a left adjoint $\pi_!$

$$2) \quad \text{Ran}(X) \text{ is } \underline{\text{contractible}} \Rightarrow \underbrace{\pi_! \pi^!}_{H_* (\text{Ran})} \xrightarrow{\sim} \text{id} \approx k$$

$$\exists \text{ sub prestack } (\text{Ran} \times \text{Ran})_{\text{disj}} \subset \text{Ran} \times \text{Ran}$$

$$\text{Ran is a semi-group.} \quad U: \text{Ran} \times \text{Ran} \longrightarrow \text{Ran}$$

A factorization alg is (roughly) the following data:

$$1) \quad A \in \text{Dmod}(\text{Ran})$$

$$2) \quad \text{factorization isos}$$

$$(U^! A) \big|_{(\text{Ran} \times \text{Ran})_{\text{disj}}} \simeq (A \boxtimes A) \big|_{(\text{Ran} \times \text{Ran})_{\text{disj}}}$$

+ higher coherence

$$\boxed{\begin{array}{ccc} A_x & & A_y \\ x & \rightarrow & y \end{array}}$$

Def. Let A be a factorization alg. The factorization (a.k.a. chiral) homology of A is $C^{\text{Fact}}(X, A) = \pi_!(A^{(\text{Ren})})$

Examples 1) A comm. fact alg (cat. of fact algs)

$$A \in \text{Comm Alg}(\text{Dmod}(X)) \xrightarrow{\text{FA}} \text{FactAlg}(X)$$

Fact (to be explained) A comm. fact. alg

$C^{\text{Fact}}(X, A)$ is a comm. alg and

$\text{Spec } C^{\text{Fact}}(X, A) =$ (derived) space of horizontal sections of $\text{Spec } A \rightarrow X$.

$$\text{Equiv} \quad \underset{\text{Comm Alg}}{\text{Maps}}(C^{\text{Fact}}(X, A), B) \simeq \underset{\text{Comm Alg}(\text{Dmod}(X))}{\text{Maps}}(A, B \otimes \omega_X)$$

Ex. $A = \mathcal{O} \otimes K[t]$ constant scheme $X \times \mathbb{A}^1 \rightarrow X$

$$\text{then } C^{\text{Fact}}(X, A) = \text{Sym}(H^{\text{dR}}(X))^{\vee}$$

2) Kac-Moody (at level 0)

$\text{Var}(\mathfrak{g}) \hookrightarrow$ fact alg corresponding to Kac-Moody

$$C^{\text{Fact}}(X, \text{Var}(\mathfrak{g})) \simeq C^{\text{Lie}}(H^{\bullet}(X) \otimes \mathfrak{g})$$

Exercise deduce 2) in the abelian case from 1)

Obs $p \in \text{Bun}_G$, $\text{Pit}_p \simeq \underline{C}^{\text{Lie}}(H^*(X, \mathfrak{g}_p))$

Units

What is a unital factorization algebra?

Version 0. \exists a "unit" alg $\omega_{\text{Ran } X}$.

Can ask $\omega_{\text{Ran } X} \rightarrow A$.

A unit is more!

Rough idea for every $x \in X$, have a map $k \rightarrow A_x$ compatible w factorization

e.g. $x, y \in X$, $A_x \rightarrow A_{\{x, y\}} \simeq A_x \otimes A_y$

Idea remember the inclusion relation on subsets.

Want to consider categorical prestacks (a.k.a. prestacks valued in cats)

i.e. functors $(\text{Sch}^{\text{aff}})^{\text{op}} \rightarrow \text{Cat}$

Main example.

$$\text{Ran}^{\text{unit}}(X)(S) = \left\{ \begin{array}{l} \text{poset of finite} \\ \text{subsets of } X(S^{\text{red}}) \end{array} \right\}$$

If X is a categorical prestack, can consider $\mathcal{QCoh}(X)$, $\text{IndCoh}(X)$, $\text{Dmod}(X)$

Concretely, $F \in \mathcal{QCoh}(X)$ is the data of

- $\forall x: S \rightarrow X$, $F_{S, x} \in \mathcal{QCoh}(S)$
- $\alpha: x \rightarrow x' \in X(S)$, $F_{S, x} \rightarrow F_{S, x'}$
- Compatibility w composition

Def A unital factorization algebra A is the data of

$$1) A^{(\text{Ran})} \in \text{Dmod}(\text{Ran}^{\text{unit}})$$

$$2) U^! (A^{(\text{Ran})}) \Big|_{(\text{Ran}^{\text{unit}})^2_{\text{disj}}} \cong (A^{(\text{Ran})} \boxtimes A^{(\text{Ran})}) \Big|_{(\text{Ran}^{\text{unit}})^2_{\text{disj}}}$$

$$3) A_{\phi} \cong k$$

...

Consider some k -points of $\text{Ran}^{\text{unit}}(X)$

$J \subset X$ finite subset

$$A_J \cong \bigotimes_{j \in J} A_{\{j\}} \quad \text{factorization}$$

$$J \subset J'$$

$$A_J \rightarrow A_{J'}$$

eg. $J = \emptyset, \quad J' = \{x\} \rightarrow k \rightarrow A_x$

$$\& A_{\{x\}} \rightarrow A_{\{x,y\}}.$$



Factorization homology

Ran^{unit} can also be expressed in terms of X^I .

$$\Rightarrow \pi^{\text{val}}_* : \text{Vect} \rightarrow \text{Dmod}(\text{Ran}^{\text{unit}}) \quad \text{has a left adjoint } \pi^{\text{fnc}}_*$$

Def A is a unital fact. alg $C^{\text{fact}}(X, A) = \pi_! (A^{(\text{Ran})})$

\exists nat'l map

$$t: \text{Ran} \rightarrow \text{Ran}^{\text{until}}$$

$$t^!: \text{FA}^{\text{until}}(x) \rightarrow \text{FA}(x)$$

Cr of contractibility of Ran : $\forall F \in \text{Dmod}(\text{Ran}^{\text{until}})$

$$\pi_! t^! F \simeq \pi_!^{\text{until}} F$$

Categorical version

Have unital fact. cat. defined in the evident way.

More concretely, if \mathcal{Y} is a categorical prestack, we can think of sheaves on \mathcal{Y}

as follows:

$$\begin{array}{c} \vdots \\ \text{Maps}([1], \mathcal{Y})^{\text{grpd}} \text{ "prestack of morphisms"} \\ s \downarrow \downarrow t \\ \mathcal{Y}^{\text{grpd}} = \text{"prestack of objects"} \end{array}$$

$$\mathcal{Y}^{\text{grpd}}(s) = \mathcal{Y}(s)^{\text{grpd}}$$


$$\text{Maps}([1], \mathcal{Y})^{\text{grpd}} = \left\{ \begin{array}{l} x, y \in \mathcal{Y}(s) \\ x \rightarrow y \end{array} \right\}$$

• sheaf on \mathcal{Y} is the data of a sheaf F on $\mathcal{Y}^{\text{grpd}}$ $\leadsto s^! F \rightarrow t^! F$ + compatibilities

Apply this to $\text{Ran}^{\text{until}}$

$$(\text{Ran}^{\text{until}})^{\text{grpd}} = \text{Ran}(\perp \phi)$$

$$\text{Maps}([1], \text{Ran}^{\text{until}})^{\text{grpd}}(s) = \left\{ \begin{array}{l} \underline{x}, \underline{x}' \in X(s_{\text{red}}) \\ \underline{x} \subset \underline{x}' \end{array} \right\} = \text{Ran}_{\subseteq}$$



Upshot: \mathcal{C} is a fact cat. The data of unitarity on \mathcal{C} is given by

$$P_{\text{small}}^*(\underline{e}) \rightarrow P_{\text{big}}^*(\underline{e}) + \dots$$

Fact P_{small}^* has a right adjoint $P_{\text{small},*}$

Can rewrite the str. as a map $\underline{e} \rightarrow P_{\text{small},*} P_{\text{big}}^*(\underline{e})$

More explicitly, suppose $\underline{x} \subset \underline{x}'$,

$$\Leftrightarrow \text{ins. unit } \underline{x} \subset \underline{x}' : \mathcal{C}_{\underline{x}} \rightarrow \mathcal{C}_{\underline{x}'}$$

Example 1) KL

2) A fact alg. \exists (weak) fact. cat. \mathcal{A} -mod fact

$$\text{ins. unit } \phi_{\mathcal{C}}\{x\}(k) = Ax$$

Let \underline{e} be a sheaf of cats on Ran^{unfl}

$\mathcal{C}^{\text{glob}}$ a category

Def A unital local-to-global functor is a map

$$\underline{e}^{\text{loc}} \rightarrow \underline{\text{Dmod}}(\text{Ran}^{\text{unfl}}) \otimes \mathcal{C}^{\text{glob}}$$

Note: what happens w/ units,

$$\text{ins. unit } \underline{x} \subset \underline{x'} : e_{\underline{x}}^{\text{loc}} \rightarrow e_{\underline{x'}}^{\text{loc}}$$

unital str. $\quad F_{\underline{x}} \searrow e^{\text{glob}} \swarrow F_{\underline{x'}}$

$$(*) \quad F_{\underline{x}} \xrightarrow{\sim} F_{\underline{x'}} \circ \text{ins. unit } \underline{x} \subset \underline{x'}$$

can weaken this to a "lax unital str."

where $(*)$ need not be an isom.

$$\text{Fun}^{\text{loc-glob, unit}}(e^{\text{loc}}, e^{\text{glob}}) \hookrightarrow \text{Fun}^{\text{loc-glob, lax-unital}}(e^{\text{loc}}, e^{\text{glob}}) \quad (**)$$

Claim $(**)$ has a left adjoint, given by $F \mapsto F \int^{\text{ins. unit}}$

$$F \int^{\text{ins. unit}} : e_{\underline{Z}}^{\text{loc}} \xrightarrow{\text{ins. unital}} e_{\underline{Z}^{\leq}}^{\text{loc}} \xrightarrow{F} e^{\text{glob}} \otimes D_{\text{mod}}(\underline{Z}^{\leq}) \xrightarrow{(P_{\text{small}})!} e_{\otimes D_{\text{mod}}(\underline{Z})}^{\text{glob}}$$

for $\underline{Z} \rightarrow \text{Ran}$

$$\text{where } \underline{Z}^{\leq} = \underline{Z} \times_{\text{Ran} \leftarrow P_{\text{big}}} \text{Ran}_{\subseteq}$$

Key example

$$e^{\text{loc}} = A\text{-mod fact} \quad \text{for a unital fact. alg } A$$

$$F: A\text{-mod fact} \longrightarrow \text{Vect} = e^{\text{glob}}$$

$$F(M) = \text{obli}(M)$$

Claim $F \int^{\text{ins. unital}} \simeq C^{\text{Fact}}(x, A, -)$

$$\text{for } Z \rightarrow \text{Ran},$$

$$F_{\text{ins. until}}^Z(M) = P_{\text{small}}!(M_{Z^{\leq}}) \in \text{Dmod}(Z)$$

$$P_{\text{small}}: Z^{\leq} \rightarrow Z$$

$$\text{Ran}_{f(3)} \rightarrow \begin{matrix} 4 \\ 3 \end{matrix}$$

$$b: Z \rightarrow \text{Ran}$$

M_x

$$F \rightarrow F^{\text{ins. until}}$$

$$\Rightarrow M_x \rightarrow C^{\text{Fact}}(X, A, M)$$

$M_x \quad N_y$

$$M_x \otimes N_y \rightarrow C^{\text{Fact}}(X, A, M_x \otimes N_y)$$

$$\text{if } N_y = A, \text{ then } C^{\text{Fact}}(X, A, M_x \otimes N_y) \simeq C^{\text{Fact}}(X, A, M_x).$$



Set up

$$\underline{e}^{\text{loc}} \in \text{Shv Cat}(\text{Ran}^{\text{until}})$$

$$\underline{e}^{\text{glob}}$$

$$(\text{lax-}) \text{ unitar local-to-global functor } F: \underline{e}^{\text{loc}} \rightarrow \underline{e}^{\text{glob}}$$

$$\underline{x} \in \text{Ran}(x), F_{\underline{x}}: \underline{e}_{\underline{x}}^{\text{loc}} \rightarrow \underline{e}^{\text{glob}}$$

$$\underline{x} \subset \underline{x}': \quad \underline{e}_{\underline{x}}^{\text{loc}} \xrightarrow{\text{ins. unit}} \underline{e}_{\underline{x}'}^{\text{loc}} \xrightarrow{\quad} \underline{e}^{\text{glob}}$$

$$\begin{array}{l} \text{diag} \\ \nearrow \\ Z \end{array} \rightarrow Z_{\leq} = Z \times_{\text{Ran}}^{\text{Ran}_{\leq}} P_{\text{small}}$$

$$e_z^{\text{loc}} \xrightarrow{\text{ins. unt}} e_{Z_{\leq}}^{\text{loc}} \xrightarrow{F} e^{\text{glob}} \otimes D_{\text{mod}}(Z_{\leq}) \xrightarrow{(P_{\text{small}})!} e^{\text{glob}} \otimes D_{\text{mod}}(Z)$$

$$\begin{array}{ccc} & \downarrow \text{diag!} & \downarrow \text{diag!} \\ & e_z^{\text{loc}} \xrightarrow{F} e^{\text{glob}} \otimes D_{\text{mod}}(Z) & \end{array}$$

$$\Rightarrow F \rightarrow F \int \text{ins. unt}$$

Lecture 5 (Lin Chen)

$$\begin{array}{ccc} \text{KL}(G)_{\text{crit}, \text{Ran}} & \xrightarrow[\sim]{\text{FLE}_{\text{crit}}} & \text{Indcoh}_*(\mathcal{O}_{P_{\check{G}}}^{\text{non-free}}) \\ \text{Loc} \downarrow & & \downarrow P_{\text{point}}^{\text{Spec}} \\ D(\text{Bun}_G)_{\text{crit}} & \xrightarrow{\Delta} & \text{Indcoh}_{\text{rep}}(LS_{\check{G}}) \\ \text{coet}_* \downarrow & & \downarrow \text{Spec} \\ \text{Whit}^!(G)_{\text{crit}, \text{Ran}} & \xrightarrow[\sim]{\text{CS}} & \text{Rep}(\check{G})_{\text{Ran}} \end{array}$$

Primary goal: describe $\text{coet}_* \circ \text{Loc}$ via local data (+ Ran magic)

Step 1 $\text{coet}^{\text{Var}} \circ \text{Loc}$ $\text{coet}^{\text{Var}} : D_{\text{crit}}(\text{Bun}_G) \rightarrow \text{Vect}$

Step 2 Hecke action + Ran magic

Other goals $CT_x \circ Loc, \quad \text{obv} \circ Loc.$

So Reflections

① $KL \quad KL(A) = \{ (Lg, L^+A) \text{-modules} \}$

$$Bun_A = Bun_A^{level \infty x} / L^+A$$

$$L_A \hookrightarrow Bun_A^{level \infty x} \text{ via regluing}$$

$$\leadsto Lxg \rightarrow T(Bun_A^{level \infty x})$$

$$\leadsto Lxg\text{-mod} \rightarrow D(Bun_A^{level \infty x})$$

$$M \mapsto \text{Diff} \otimes_{\underline{U(Lg)}} M$$

$$\leadsto Lxg\text{-mod}^{L^+A} \rightarrow D(Bun_A)$$

This works for any $x \in \text{Ran}$.

$$\underline{Loc}: \underline{KL}(A) \longrightarrow D(Bun_A) \otimes \underline{Vect} \quad \text{in } \text{ShvCat}(\text{Ran})$$

It is even in $\text{ShvCat}(\text{Ran}^{unrl})$

$$\underline{Ex.} \quad KL(A)_x \xrightarrow{Loc_x} D(Bun_A)$$

ins+vac. \downarrow

$$KL(A)_x = KL(A)_y \xrightarrow{Loc_y} D(Bun_A)$$

commutes

$KL(A)_y/x$ for $x < y \in \text{Ran}^{unrl}$

$$\text{Loc} : \text{KL}(G)_{\text{Ran}} \xrightarrow{\text{Loc}_{\text{Ran}}} D(\text{Bun}_G \times \text{Ran}) \xrightarrow{\int_{\text{Ran}}} D(\text{Bun}_G)$$

This is the functor Loc .

③ Coet_*

$$\text{Whit}^!(G) = D(\omega_{G,x})^{(LN,x)}$$

$$\omega_{G,x} \xrightarrow{\pi} \text{Bun}_G$$

$$\text{Poinc!}_x : \text{Whit}^!(G)_x \xrightleftharpoons[\text{Ad}_*^{\text{Whit}} \circ \pi^!]{\pi^!} D(\text{Bun}_G) : \text{Coet}_{*,x}$$

$$\text{Poinc!} : \text{Whit}^!(G) \xrightleftharpoons{\quad} D(\text{Bun}_G) \otimes \text{Vect} : \text{Coet}_*$$

$$\text{Vect} \xrightarrow{\text{unit}} \text{Whit}(G)$$

$$\begin{array}{ccc} \omega_G & \xleftarrow{p} & \omega_N \supset \exp \\ & & \downarrow \quad \uparrow \\ & & G_N \supset \exp \end{array}$$


is given by $p_!(\exp \omega_N)$

$$p_!(\exp)$$


$$\text{Exer. } ① \quad \text{unit}^R : \text{Whit}^!(G) \xrightarrow{e^!} \text{Vect}$$

$$\begin{array}{ccc} \text{Whit}^!(G) & \xrightarrow[\sim]{CS} & \text{Rep}(\check{G}) \\ e^! \searrow & & \swarrow \text{inv} \\ & \text{Vect} & \end{array}$$

Coeth_{*}:



Exer Coeth^{Vac} is given by $q_*(p^!(-) \otimes x^! \exp)$



Advanced Exer


① Poinc! is unital, Coeth_{*} is right lax unital.

② $\phi \in \text{Ran}^{\text{unfl}}$ Coeth_{*}, $\phi : D(\text{Bun}_A) \rightarrow \text{Vect}$ is Coeth^{Vac}.

— Σ

§1. Statements

Main-Thm



$$\text{Ran}^c = \{x \subset y : x, y \in \text{Ran}\}$$

$\begin{array}{cc} p_{\text{small}}^* \swarrow & \searrow p_{\text{big}}^* \\ \text{Ran} & \text{Ran} \end{array}$

$$\begin{aligned}
 e &\in \text{ShvCat}(\text{Ran}) \\
 p_{\text{big}}^*(e) &\in \text{ShvCat}(\text{Ran}^c) \\
 e_{\text{Ran}^c} &:= \Gamma(\text{Ran}^c, p_{\text{big}}^*(e))
 \end{aligned}$$

Recollection on DS.

$$DS(M) = C^{\frac{\infty}{2}}(L\mathfrak{n}, L^+\mathfrak{N}, M \otimes k\chi)$$

$$\text{For } L\mathfrak{N} = \bigcup \mathfrak{N}_i \quad (\mathfrak{N}_i = \text{Ad}_{t^{-i}P} L^+\mathfrak{N})$$

$$C^{\frac{\infty}{2}}(L\mathfrak{n}, L^+\mathfrak{N}, M) = \varinjlim_i C^*(\mathfrak{n}_i, M \otimes \det(\mathfrak{n}_i/\mathfrak{n}_0) [\dim \mathfrak{N}_i / \mathfrak{N}_0])$$

Exer Define the connecting maps.

$$\underline{DS} : \underline{KL}(\mathfrak{h}) \longrightarrow \underline{Vect}$$

is a lax unital functor.

$$DS(\mathbb{W}ac) = \mathfrak{Z} = \mathbb{Z}(\mathbb{W}ac)$$

Exercise (Tautologous)

$$\underline{DS}^{enh} : \underline{KL}(\mathfrak{h}) \longrightarrow \underline{\mathfrak{Z}\text{-mod fact}}$$

$$\underline{C_*^{fact}}(x, \mathfrak{z}, -) : \underline{\mathfrak{Z}\text{-mod fact}} \longrightarrow \underline{Vect}$$

$$\begin{array}{ccc} F : \underline{e} & \xrightarrow{F} & \underline{D} \\ \text{Fench} \swarrow & & \nearrow \text{oblv} \\ & \underline{F(1_e)\text{-mod fact}} & \end{array}$$

Rewrite \hookrightarrow as

$$KL(\mathfrak{h})_{\text{Ran}}$$

$$\downarrow \underline{DS}^{enh}$$

$$\mathfrak{Z}\text{-mod fact}_{\text{Ran}} \xrightarrow{\underline{C_*^{fact}}(x, \mathfrak{z}, -)_{\text{Ran}}} D(\text{Ran}) \xrightarrow{\int_{\text{Ran}}} \underline{Vect}$$

$$1) \quad \underline{KL_x} \xrightarrow{Loc} D_{mod}(\text{Bun}_{\mathfrak{h}})$$

$x \in \text{Ran}$

$$2) \underline{KL} \xrightarrow{\underline{Loc}}^{unital \quad \text{local-to-global}} D_{mod}(Bun_A) \otimes \underline{Vect}$$

$$3) \underline{KL}_{Ran} \xrightarrow{\underline{Loc}_{Ran}} D_{mod}(Bun_A) \otimes D_{mod}(Ran) \xrightarrow{Id \otimes \int_{Ran}} D_{mod}(Bun_A)$$

$$4) \underline{KL} \xrightarrow{DS} \underline{Vect} \quad \text{lax unital}$$

$$\underline{KL} \xrightarrow{\underline{Loc}} D(Bun_A) \otimes \underline{Vect}$$

$$\downarrow \underline{DS} \qquad \downarrow \text{coet} \otimes id$$

$$\underline{Vect} \xrightarrow{id} \underline{Vect}$$



Construction

$$\underline{KL}(A) \xrightarrow{\underline{Loc}} D(Bun_A) \otimes \underline{Vect}$$

$$\begin{array}{ccc} \textcircled{DS} \downarrow & \Downarrow & \downarrow \text{coet}^{vac} \otimes id \\ \text{lax unital } \underline{Vect} & \xlongequal{\quad} & \underline{Vect} \end{array}$$

$$\begin{array}{ccc} \underline{Vac} & \xrightarrow{\quad} & D(1) \otimes 1 \\ \downarrow & & \downarrow \\ DS(\underline{Vac}) & \neq & \text{coet}^{vac}(D(1)) \otimes 1 \\ \text{||} & & \\ \emptyset & & \end{array}$$

$$F: \underline{e} \rightarrow D \otimes \underline{Vect} \quad \text{lax unital} \quad \text{local-to-global}$$

$$F \xrightarrow{\quad} F^{ins.vac} = F^{fix}$$

$$\begin{array}{ccc} \underline{e}_x & \xrightarrow{F_x^{fix}} & D \\ \text{ins.vac} \downarrow & & \uparrow \int_{Ran_x} \\ \underline{e}_{Ran_x} & \xrightarrow{F_{Ran_x}} & D \otimes D(Ran_x) \end{array}$$

$$F_x^{fix}: \underline{e}_x \rightarrow D$$

$$\left. \begin{array}{c} \\ \end{array} \right\} F^{fix}: \underline{e} \rightarrow D \otimes \underline{Vect}$$

$$\begin{array}{ccc} \underline{DS} & \xrightarrow{\quad} & \underline{coet}^{vac} \circ \underline{Loc} \\ \searrow & & \nearrow \\ & \underline{DS}^{fix} & \end{array}$$

$$\text{Construction} \Rightarrow \underline{DS}^{fix} \xrightarrow{\quad} \underline{coet}^{vac} \circ \underline{Loc}$$

Main - Thm'

$$DS^{b'x} \xrightarrow{\quad} \underline{Loc}^{vac} \circ \underline{Loc}$$

$\xrightarrow{\quad}$

§2. Loc & obl

$$triv \in \underline{Rep}(L^+A) \xrightarrow{\underline{Loc}^{obl}} \underline{Obh}(Bun_A) \otimes \underline{Vect} \Rightarrow \underline{0} \otimes \underline{1}$$

$$\text{ind} \downarrow \uparrow \underline{obl}$$



$$\downarrow \text{ind} \otimes \text{id} \uparrow \underline{obl} \otimes \text{id}$$

$$vac \in \underline{KL}(A) \xrightarrow{\underline{Loc}} D(Bun_A) \otimes \underline{Vect} \simeq \underline{Diff} \otimes \underline{1}$$

• $\begin{array}{c} \uparrow \\ \text{---} \end{array}$ is unital

$$\left(\underline{Loc}^{obl} \circ \underline{obl} \right)^{b'x}$$



$$\underline{obl} \circ \underline{Loc}$$

• $\begin{array}{c} \text{---} \\ \downarrow \end{array}$ is lax unital

Thm 2 This is \simeq .

$\xrightarrow{\quad}$

§3. Loc & pullback

$$\begin{array}{ccc} H_1 & \simeq & Y_1 \\ \downarrow & & \downarrow p \\ H_2 & \simeq & Y_2 \end{array}$$

$$\begin{array}{ccc} h_1\text{-mod} & \xrightarrow{\underline{Loc}} & D(Y_1) \\ \uparrow \text{res} & \Rightarrow & \uparrow p^! \\ h_2\text{-mod} & \xrightarrow{\underline{Loc}} & D(Y_2) \end{array}$$

$$\begin{array}{ccc} \underline{KL}(H_1) & \xrightarrow{\underline{Loc}_{H_1}} & D(Bun_{H_1}) \otimes \underline{Vect} \\ \uparrow \text{res} & \Rightarrow & \uparrow p^! \\ \underline{KL}(H_2) & \xrightarrow{\underline{Loc}_{H_2}} & D(Bun_{H_2}) \otimes \underline{Vect} \end{array}$$

Thm 3 $(\underline{Loc}_{H_1} \circ \underline{res})^{fix} \rightsquigarrow \underline{P}^! \circ \underline{Loc}_{H_2}$

Ex $\text{Thm 2} \Rightarrow \text{Thm 3}.$

(Hint $(F \circ \phi)^{fix} \simeq F^{fix} \circ \phi$ if ϕ is unital

$(\psi \circ F)^{fix} \simeq \psi \circ F^{fix}$ if $\psi: D \otimes \underline{Vect} \rightarrow D' \otimes \underline{Vect}$)

$$\begin{array}{ccc} \underline{KL}(H_1) & \xleftarrow[\underline{Loc}_{H_1}]{\Gamma_{H_1}^{level \infty}} & D(\text{Bun}_{H_1}) \otimes \underline{Vect} \\ \downarrow \underline{res} & \searrow \underline{Loc} & \downarrow \underline{P}_* \\ \underline{KL}(H_2) & \xleftarrow[\underline{Loc}_{H_2}]{\Gamma_{H_2}^{level \infty}} & D(\text{Bun}_{H_2}) \otimes \underline{Vect} \end{array}$$

\Leftrightarrow Duality + Beck - Chevalley

$C^{\infty}_{\frac{1}{2}}(L_n, L^+N, -) = \text{BRST}_n$

Thm 4 If $H_1 \rightarrow H_2$, $\text{kernel} = N$,

$(\underline{Loc}_{H_2} \circ \underline{BRST}_n)^{fix} \rightsquigarrow \underline{P}_* \circ \underline{Loc}_{H_1}$

$\text{Thm 3} + \text{Thm 4} + \text{fact} \Rightarrow \underline{\text{Main Thm}}$

$$\begin{array}{ccc} \underline{KL}(N) & \xrightarrow{\underline{Loc}_N} & D(\text{Bun}_N) \otimes \underline{Vect} \\ \downarrow C^{\infty}_{\frac{1}{2}}(L_n, L^+N) \downarrow \text{BRST}_n & \nearrow & \downarrow \Gamma_{dR}(\text{Bun}_N) \\ \underline{KL}(N)_x & \xrightarrow{\underline{Loc}} D(\text{Bun}_x) \xrightarrow{\Gamma_{dR}} \underline{Vect} & \nearrow \uparrow \text{fact} \\ \downarrow \text{ins. val} & & \uparrow \int_{\text{Ran}_x} \\ \underline{KL}(N)_{\text{Ran}_x} & \xrightarrow{\text{BRST}_n} D(\text{Ran}_x) \supset \mathcal{O}(\text{BL}^+N)_{\text{Ran}_x} & = \Omega_n \end{array}$$

$$\text{BRST}_n(\text{Vac}) = \text{End}_{\text{Rep}(\mathbb{L}^+ N)}(\text{triv}, \text{triv}) \simeq \mathcal{O}(\mathbb{B}\mathbb{L}^+ N)$$

Thm 4': $\mathcal{O}(\text{Bun}_N) = \mathcal{C}^{\text{fact}}(X, \mathcal{O}(\mathbb{B}\mathbb{L}^+ N))$

$$\text{Bun}_N = \text{Maps}_X(X, \mathbb{B}N \times X)$$

$$\mathbb{B}\mathbb{L}^+ N = \text{Maps}_X(D, \mathbb{B}N \times X)$$

$Y \longrightarrow X$ affine D -scheme

$$\mathcal{O}(\text{Sect}_D(X, Y)) = \mathcal{C}^{\text{fact}}(X, \mathcal{O}(\mathbb{L}_D^+ Y))$$

Now \bullet Y is Weil restriction of $\mathbb{B}N \times X$ along $X \rightarrow X \times \mathbb{A}^1$

\bullet $\mathbb{B}N$ is as good as affine $\mathcal{O}(\mathbb{B}N) \simeq \mathcal{O}(\mathbb{B}N)_{\text{mod}}$

Tutorial 3-3 (Kevin Lin)

(fix a point $x \in X$)

Recall $\text{Op}_{\check{h}}^{\text{mer}} = \check{h}^{\vee}$ -opers on \mathbb{P}_x^0

eg. if $\check{h} = hL_3$

$$d + \begin{pmatrix} a_1 & a_2 & a_3 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} dt$$

$$a_1 \in k((t))$$

$$a_2 \in k((t)) dt$$

$$a_3 \in k((t)) dt^2$$

In general, $\text{Op}_{\check{h}}^{\text{mer}} = \text{colim } V_{\leq n}$

$$V_{\leq n} = \lim V_{\leq n} / V_{\leq -m} \quad (\text{fin dim. terms})$$

$$\begin{array}{ccc}
 \mathcal{O}_{\check{X}}^{m.b.} & \longrightarrow & \mathcal{O}_{\check{X}}^{mer} \\
 \downarrow & \searrow & \downarrow \\
 \mathcal{L}_{\check{X}}^{reg} & \longrightarrow & \mathcal{L}_{\check{X}}^{mer}
 \end{array}$$

Want to define a functor $\text{Poinc}_x^{\text{Spec}} : \text{IndCoh}_x(\mathcal{O}_{\check{X}}^{m.b.}) \rightarrow \text{IndCoh}(\mathcal{L}_{\check{X}}^v)$

We have:

$$\begin{array}{ccc}
 & \mathcal{O}_{\check{X}}^{mer, glob} & \\
 \swarrow \text{ev}^{mer} & & \searrow \\
 \mathcal{O}_{\check{X}}^{mer} & & \mathcal{L}_{\check{X}}^{mer, glob}
 \end{array}$$

$\mathcal{O}_{\check{X}}^{mer, glob} = \text{opens on } X \setminus x$
 $\mathcal{L}_{\check{X}}^{mer, glob} = \text{local systems on } X \setminus x$

the diagram is over $\mathcal{L}_{\check{X}}^{mer}$

Base change along $\mathcal{L}_{\check{X}}^{reg} \rightarrow \mathcal{L}_{\check{X}}^{mer}$, to get

$$\begin{array}{ccc}
 & \mathcal{O}_{\check{X}}^{m.b., glob} & \\
 \swarrow \text{ev}^{m.b.} & & \searrow \tau \\
 \mathcal{O}_{\check{X}}^{m.b.} & & \mathcal{L}_{\check{X}}^v
 \end{array}$$

"Def" $\text{Poinc}_x^{\text{Spec}} = \tau_* \circ (\text{ev}^{m.b.})^*$

Because we did base change along $\mathcal{L}_{\check{X}}^{reg} \rightarrow \mathcal{L}_{\check{X}}^{mer}$, $\text{Poinc}_x^{\text{Spec}}$ is linear over

$$\text{IndCoh}(\mathcal{L}_{\check{X}}^{reg} \times_{\mathcal{L}_{\check{X}}^{mer}} \mathcal{L}_{\check{X}}^{reg})$$

One can show directly that $\text{IndCoh}_x(\mathcal{O}_{\check{X}}^{m.b.})$ is temporized.

$$\Rightarrow \text{Poinc}_x^{\text{Spec}} \text{ lands in } \text{IndCoh}(\mathcal{L}_{\check{X}}^v)^{\text{temp}} \simeq \mathcal{QCoh}(\mathcal{L}_{\check{X}}^v)$$

Warning even if $f: X \rightarrow Y$ is a map of finite type schemes, the functor

$f_*: \text{IndCoh}(X) \rightarrow \text{IndCoh}(Y)$ need not admit a left adjoint f^*

- eg. $\text{pt} \rightarrow \text{pt} \times_{\mathbb{A}^1} \text{pt}$ has no IndCoh $*$ -pullback.

- eg. if f is quasi-smooth, f^* exists.

(In general, need f to have finite tor amplitude.)

Why $(e_{\text{mer}})^*$ exists?

$$\text{IndCoh}_*(\mathcal{O}_{P_{\check{X}}^{\text{mer}}}) = \text{colim}_{*-\text{push}} \text{IndCoh}_*(V_{\leq n})$$

$$\text{IndCoh}_*(V_{\leq n}) = \text{colim}_{*- \text{pull}} \text{IndCoh}(V_{\leq n}/V_{\leq -m})$$

$$\mathcal{O}_{P_{\check{X}}^{\text{mer, glob}}} = \text{colim } L_{\leq n} \quad (\text{each fin. dim})$$

$$\text{for } a \in L_3 \quad d + \begin{pmatrix} a_1 & a_2 & a_3 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} dt \quad a_i \in \Gamma(X \setminus x, \omega^{\otimes ?})$$

$$L_{\leq n} \hookrightarrow \mathcal{O}_{P_{\check{X}}^{\text{mer, glob}}}$$

$$\begin{array}{ccc} \downarrow & & \downarrow \\ V_{\leq n} & \longrightarrow & \mathcal{O}_{P_{\check{X}}^{\text{mer}}} \end{array}$$

$$\downarrow \\ V_{\leq n}/V_{\leq -m}$$

$*$ -pullback along $L_{\leq n} \rightarrow V_{\leq n}/V_{\leq -m}$ exists.

$$\underline{\text{Thm}} \quad \text{Ind Coh}_* (\mathcal{O}_{P_{\check{X}}^{n.b.}})_x \xrightarrow{\text{Poinc Spec}_x} \text{Ind Coh} (LS_{\check{X}})$$

$$(2^+)^* \downarrow$$

$$\text{Ind Coh}_* (\mathcal{O}_{P_{\check{X}}^{n.b.}}) \longrightarrow \mathcal{O}_{P_{\check{X}}^{n.b.}} \xrightarrow{\text{Fact C. } (-)} \text{Vect}$$

commutes.

Proof by example. Plug in $(2^+)^* \mathcal{O}_{P_{\check{X}}^{\text{reg}}} \in \text{Ind Coh}_* (\mathcal{O}_{P_{\check{X}}^{n.b.}})$

$$\begin{array}{ccc} \mathcal{O}_{P_{\check{X}}} & \longleftarrow & \mathcal{O}_{P_{\check{X}}}^{\text{glob}} \\ \downarrow 2^+ & & \downarrow \\ \mathcal{O}_{P_{\check{X}}}^{n.b.} & \longleftarrow & \mathcal{O}_{P_{\check{X}}}^{n.b., \text{glob}} \end{array}$$

upper circuit is functions on $\mathcal{O}_{P_{\check{X}}}^{\text{glob}}$

lower circuit is $C_*^{\text{Fact}}(X, \mathcal{O}_{P_{\check{X}}^{\text{reg}}})$

Nick $\Rightarrow \mathcal{O}_{P_{\check{X}}}^{\text{glob}}$ is an affine scheme w/ functions

$$C_*^{\text{Fact}}(X; \mathcal{O}_{P_{\check{X}}^{\text{reg}}})$$

$y \rightarrow X$ affine D-scheme

It is possible to find $y \rightarrow y_i$ modifying y at x , $L_{\nabla} y = \text{colim } L_{\nabla}^+ y_i$

$$\text{Ind Coh}_* (L_{\nabla} y)_x \longrightarrow \mathcal{O}_{L_{\nabla}^+ y - \text{mod } x}^{\text{fact}}$$

$$\begin{array}{ccc} \text{f-pull} \downarrow & & \downarrow C_*^{\text{Fact}} \\ \text{Ind Coh}_* (\text{Sect}_{\nabla}(X \setminus x, y)) & \xrightarrow{\quad} & \text{Vect} \end{array}$$

$$(B_{\check{G}})_{\text{Ran}} \xleftarrow{\text{ev}} LS_{\check{X}} \times \text{Ran}$$

$$\downarrow p$$

$$Loc^{\text{Spec}} = p_! \circ \text{ev}^* \quad , \quad \uparrow_{\text{Spec}} = \text{ev}_x \circ p^! = (Loc^{\text{Spec}})^R$$

Day 3 Q & A

non-free ops for G_m :

$$\mathcal{O}_{P_{G_m}} = \text{connections on trivial bundle} = \{1\text{-forms}\}$$

$$\{x\}^{cl} \text{Spec } \Gamma(\text{Bun}_{G_m}, \text{Diff}) = \text{pt}(\Omega^1) = \mathcal{O}_{P_{G_m}}(x)$$

$$\mathcal{O}_P^{mb}(D) \subset \mathcal{O}_P^{mer}(D)$$

$$\left\{ d + f dt : \begin{array}{l} a_{-1} \in \mathbb{Z}, \\ a_{-n} \text{ is nilp for } n > 1 \end{array} \right\} \left\{ d + f(t) dt : f(t) = \sum a_i t^i \in k((t)) \right\}$$

$$\begin{array}{c} a_{-1} \\ \downarrow \\ \mathbb{Z} \end{array}$$

$$\mathcal{O}_P^{mb} = \bigoplus \mathcal{O}_P^{mb,n}$$

$$\text{each cpt: } k[[t]] \times (k((t)) / t^{-1} k[[t]])^\wedge$$

$$\mathcal{Q}coh(LS_{G_m}^\vee) \otimes D(\text{Bun}_{G_m}) \xrightarrow{\text{spec. act.}} D(\text{Bun}_{G_m}) \xrightarrow{\text{coeff}} \text{Vect}$$

$$\text{dualize } D(\text{Bun}_{G_m}) \xrightarrow{\mathbb{L}_{G_m, \text{coarse}}} \mathcal{Q}coh(LS_{G_m}^\vee)^\vee = \mathcal{Q}coh(LS_{G_m}^\vee)$$

$$\text{Rep } \check{G} \otimes KL_G \xrightarrow{\text{Sat}^{nv} \otimes \text{id}} \text{Sph}(G) \otimes KL_G \xrightarrow{\text{conv}} KL_G \xrightarrow{\text{DS}} \text{Vect}$$

$$\text{DS}(M) = C^\infty(Ln, L^+n, M \otimes \psi)$$

$$\psi: n((t)) \rightarrow k, \frac{e_i}{t} \mapsto 1$$

$$V_k \xrightarrow{\text{id}} W_k$$

every other
basis vector $\mapsto 0$

$$\left\{ \begin{array}{c} L^+g \\ L^+(f+b/N) \end{array} \right\}$$

$$f+b/N \supset g/N \approx t/w$$

$$W_{\text{crit}} = \mathcal{O}_{P_{G_m}}$$

$$\text{dualize: } (KL_G \xrightarrow{DS^{enh}} (\text{Rep } \check{G})^\vee = \text{Rep } \check{G}$$

$$DS^{enh}(\text{Vac}_{out}) \approx \tau_* \mathcal{O}_{P_{\check{G}}^{res}} = R\mathcal{Z}, \quad \tau: \mathcal{O}_{P_{\check{G}}^{res}} \rightarrow LS_{\check{G}}^\vee(D) = B\check{G}$$

\uparrow
 birth of opers

$$KL_G \rightarrow \text{Rep } \check{G}$$

$$\text{Vac}_{out} \mapsto R\mathcal{Z}$$

$$KL_G \rightarrow R\mathcal{Z}\text{-mod}^{\text{fact}}(\text{Rep } \check{G}) \stackrel{\text{up to left completion}}{\approx} \text{Ind Coh}(\mathcal{O}_{P_{\check{G}}^{mer}})$$

$$\mathcal{Z}\text{-mod}^{\text{fact}} \approx \text{Ind Coh}(\mathcal{O}_{P_{\check{G}}^{mer}})$$

Tutorial 4.1 (Sam Raskin)

$G = \text{PGL}_2$

Analogy w/ no. theory:

Given $\sigma: "Gal \mathcal{O}" \rightarrow SL_2 = \check{G}$ irred (odd) rep'n, unram.

$$a_p = \text{tr}(\sigma(F_p)), \quad F_p \in "Gal \mathcal{O}"$$

$$\text{Write } q = e^{2\pi i \tau}, \quad f(\tau) = \sum a_n q^n$$

where $a_0 = 0 \leftarrow \text{const. term} = 0$ "cuspidal" \leftrightarrow irred

$$a_1 = 1$$

$$a_p = \text{tr}(\sigma(F_p))$$

$$a_n \cdot a_m = a_{nm} \quad \text{if } (n, m) = 1$$

$$a_{pn+1} + p a_{pn-1} = a_p \cdot a_{pn}$$

Langlands conj.

$f(\tau)$ (defined on $\text{Im } \tau > 0$) is a modular form of level 1

+ "Essentially every modular form has this type".

In geometry:

$\sigma \in \text{LS}_A^\vee$ (ideally irred)

Geom. Langlands: \exists eigen sheaf $F_\sigma \in \text{Dmod}(\text{Bun}_A)$

Motto: Characterize F_σ by its "q-expansion".

Analogy: $f \mapsto a_0(f) \longleftrightarrow \text{CT}_*: \text{D}(\text{Bun}_A) \rightarrow \text{D}(\text{Bun}_M)$ M. Leri
 mod forms $\longleftrightarrow \begin{cases} \text{functs} \\ \text{on } \mathbb{R}^{\geq 0} \end{cases}$

$a_0(f) = 0 \longleftrightarrow F_\sigma$ is "cuspidal" for σ irred.

$a_1(f) \longleftrightarrow \text{coeff}_0 : \text{D}(\text{Bun}_A) \rightarrow \text{Vect}$
 "vacuum wht. coeff."


$a_1(f) = 1 \longleftrightarrow \text{coeff}(F_\sigma) = k$
 "Whittaker normalization"

Analogy harmonic analysis / \mathbb{R} & adelic stuff & geometry

$$\begin{array}{ccc} \mathbb{R} & \mathbb{A}_A & \bar{\mathbb{A}}_X \\ \cup & \cup & \cup \\ \mathbb{Z} & \mathbb{O} & k(x) \\ \downarrow & & \downarrow \\ k \backslash G(\mathbb{R}) / G(\mathbb{Z}) & \longleftrightarrow & \text{Bun}_A \end{array}$$

$$a_1(f) = \int_{\mathbb{R}/\mathbb{Z}} f(\tau) e^{-2\pi i \tau} d\tau$$

$$\text{coeff}(F) = \int_{\text{Bun}_N^{\mathbb{R}}} F \cdot \exp$$



Analogy $\mathbb{R}/\mathbb{Z} \xrightarrow[\text{adelic}]{\text{int}} \mathbb{A}/\mathbb{C} \rightsquigarrow \text{Bun}_{G_A}^{\mathbb{R}} = \text{Bun}_N^{\mathbb{R}}$ $G = \text{PGL}_2$

other coeffs.

$$n > 1 \iff P_1^{\nu_1} \dots P_s^{\nu_s} \rightsquigarrow D = \sum \nu_i [P_i] \leftarrow \text{eff. divisor on } \text{Spec } \mathbb{Z}$$

Whittaker coeffs for PGL_2 indexed by $D \geq 0$ on X .

$$\text{coeff}_D: D(\text{Bun}_{\text{PGL}_2}) \longrightarrow \text{Vect}$$

$$\{0 \rightarrow \mathcal{R}(-D) \rightarrow \mathcal{E} \rightarrow 0 \rightarrow 0\} = \text{Bun}_N^{\mathcal{R}(-D)}$$

$\text{Bun}_{\text{PGL}_2}$

$$H^1(\mathcal{R}(-D)) \longrightarrow H^1(\mathcal{R}) = \mathbb{A}^2$$

$$f(q) = \sum a_n q^n \quad [a_n] \text{ given by same procedure}$$

Analogy: $\sigma \in \text{LS}_{\text{SL}_2}$

$$D = \sum n_i x_i, \quad \text{coeff}_D(F_\sigma) = \bigotimes_i \text{sgn}^{n_i}(\sigma x_i)$$

For general h

Whittaker coeffs are (naively) indexed by divisors D on X valued in Λ^+

↑
dominant
(co)weight

$$\sum \lambda_i x_i, \quad \lambda_i \in \Lambda^+$$

$$\begin{array}{c} \text{Bun}_N^{\mathcal{O}(-D)} \\ \swarrow \quad \searrow \\ \text{Bun}_h \qquad \prod_i H'(\mathcal{O}(\alpha_i(-D))) \\ \qquad \qquad \downarrow \\ \qquad \qquad \prod H'(\mathcal{O}) \\ \qquad \qquad \downarrow \text{sum} \\ A^1 = H'(\mathcal{O}) \end{array}$$

$$\text{coeff}_D(F_\sigma) = \bigotimes_i (V_\sigma^{\lambda_i})|_{x_i}, \quad \sigma \in \text{LS}_h^\vee$$

$$D = \sum \lambda_i x_i$$

Need: Smarter versions of Whittaker coefficients w/ more geometry.

Easy upgrade

$$h = \text{PGL}_2$$

$$\begin{array}{c} \text{coeff}_D \nearrow \\ \text{Vect} \\ \uparrow \text{!-fiber at } D \end{array}$$

$$\forall d \geq 0, \text{ can easily define } \text{coeff}_d: D(\text{Bun}_h) \rightarrow D(\text{Sym}^d X)$$

Can easily adapt the above to say what $\text{coeff}_D(F_\sigma)$ is. space of $D \geq 0$, $\deg D = d$


$$D(\text{Bun}_{\text{PAL}_2}) \longrightarrow \prod_{d \geq 0} D(\text{Sym}^d)$$

\uparrow
 (can't be anywhere near fully faithful)

$\{\text{mod. forms}\}$
 $\{q\text{-expansions}\}$

Fix

$$\text{Coet} : D(\text{Bun}_h) \longrightarrow \text{Rep } \check{G}_{\text{Ran}}$$



Defn of coet is more subtle.

$$D(\text{Bun}_h) \longrightarrow \text{Whit}_x(\omega_{2h}, \text{Ran})$$

\downarrow ← Casselman - Shalika
 $\text{Rep } \check{G}_{\text{Ran}}$

What happened secretly on Monday:

$$D(\text{Bun}_h) \xrightarrow{\text{IL}_h, \text{Coet}} \text{Coh}(LS_h^v)$$

$\downarrow \text{r spec}$
 $\text{Rep } \check{G}_{\text{Ran}}$

$\swarrow \text{Coet}$

True thms :

$$D(\text{Bun}_h)_{\text{cusp}} \xrightarrow{\wedge} D(\text{Bun}_h)_{\text{temp}} \subset \text{Rep } \check{G}_{\text{Ran}}$$

\nwarrow THM !

What happened yesterday

BD thm: $\text{Fun}(\mathcal{O}_{P_h^\vee}(x)) \leadsto \text{Ditt}_{\text{out}} \in D(\text{Bun}_h)$

$\sigma \in \mathcal{O}_{P_h^\vee}(x)$ defined F_σ as $\text{Ditt}_{\text{out}} / m_\sigma$

Big goal:

$$\begin{array}{ccc}
 \text{KL}(h) & \xrightarrow{\sim \text{FLE}_{\text{out}}} & \text{IndCoh}^*(\mathcal{O}_{P_h^{\text{mt}}}) \\
 \text{Loc} \downarrow & & \downarrow \text{Poinc}_*^{\text{Spec}} \\
 D(\text{Bun}_h) & \xrightarrow{\mathbb{L}_h} & \text{IndCoh}_{\text{nilp}}(LS_h^\vee) \\
 & \searrow \text{weft} & \downarrow \\
 & & \text{Rep}_{\check{h}}^{\text{Ran}}
 \end{array}$$

$$\begin{aligned}
 \text{Ditt}_{\text{out}} = \text{Loc}(W) \quad \mathbb{L}_h(\text{Ditt}_{\text{out}}) &= \text{Poinc}_*^{\text{Spec}} \text{FLE}(W_{\text{out}}) \\
 &= \text{Poinc}_*^{\text{Spec}} (\mathcal{O}_{\mathcal{O}_{P_h^\vee}^{\text{reg}}}) \\
 &= \mathcal{O}_{\mathcal{O}_P(x)} \in \text{QCoh}(LS_h^\vee)
 \end{aligned}$$

Work from yesterday $\Rightarrow \mathbb{L}_h(F_\sigma) = \text{skyscraper at } \sigma \in \mathcal{O}_P(x) \subset LS_h^\vee$

in other words, F_σ has the right left weft.

Lecture 6 (Nick Rozenblum)

Langlands functor.

$$\begin{array}{ccc}
 \text{Rep}(h)_{\text{Ran}} & \leadsto & D_{\text{mod}}(\text{Bun}_h) \xrightarrow{\text{weft}^{\text{vac}}} \text{Vect} \\
 \downarrow & & \\
 \text{QCoh}(LS_h^\vee) & &
 \end{array}$$

$$\sim \Delta_{\text{coarse}}: D_{\text{mod}}(B_{\text{un}}) \rightarrow \text{Reloh}(LS_{\check{h}}^v)$$

Δ is a renormalized version of Δ_{coarse}

$$\text{coeff}^{\text{val}}: D_{\text{mod}}(B_{\text{un}}) \longrightarrow \text{Vect}$$

$$\begin{array}{c} \nearrow \text{Whit}(\check{h})_{\text{Ran}} \searrow \\ \text{equiv. for} \end{array}$$

Satake action

\Rightarrow we have a comm. diagram

$$\begin{array}{ccc} D_{\text{mod}}(B_{\text{un}}) & \xrightarrow{\Delta} & \text{Indoh}_{\text{nilp}}(LS_{\check{h}}^v) \\ \text{coeff} \downarrow & & \downarrow \\ \text{Whit}(\check{h})_{\text{Ran}} & \xrightarrow[\sim]{\text{CS}} & \text{Rep}(\check{h})_{\text{Ran}} \\ \downarrow & & \downarrow \text{inv} \\ \text{Vect} & = & \text{Vect} \end{array}$$

$$\text{Thm} \quad \text{KL}_{\text{Ran}} \xrightarrow[\sim]{\text{FLE}} \text{Indoh}^*(\mathcal{O}_{P_{\check{h}}^{\text{mf}}})_{\text{Ran}}$$

$$\begin{array}{ccc} \text{Loc} \downarrow & & \downarrow \text{Poinc Spec} \\ D_{\text{mod}}(B_{\text{un}}) & \xrightarrow{\Delta} & \text{Indoh}_{\text{nilp}}(LS_{\check{h}}^v) \end{array}$$

canonically commutes

Recall temperedness

Derived Satake

$$\begin{array}{l} \text{Dmod}(L_{\check{h}}/L_{\check{h}}/L_{\check{h}}) \xrightarrow{\sim} \text{Sph}_{\check{h}} \simeq \text{Indoh} \left(\text{pt}/\check{h} \times_{\check{g}/\check{h}}^* \text{pt}/\check{h} \right) \\ \text{Reloh} \left(\text{pt}/\check{h} \right) = \text{Sph}_{\check{h}, \text{temp}} \end{array}$$

$$Sph_A \leadsto e, \quad e^{temp} := e \otimes_{Sph_A} Sph_A^{temp} \hookrightarrow e$$

In particular, we have $Dmod(Bun_A)^{temp} \subset Dmod(Bun_A)$

Fact: \mathbb{L} commutes w/ Sph_A -actions

$$Ind_{Nilp}^{Loc}(LS_A^v)^{temp} \simeq Qcoh(LS_A^v)$$

Upshot

$$\begin{array}{ccc} Dmod(Bun_A)^{temp} & \xrightarrow{\mathbb{L}} & Qcoh(LS_A^v) \\ \downarrow & & \downarrow \\ Dmod(Bun_A) & \xrightarrow{\mathbb{L}} & Ind_{Nilp}^{Loc}(LS_A^v) \end{array}$$

Facts. 1) $(KL)^{temp} \simeq KL$

2) $(Ind_{Loc} * (Op_A^{mf}))^{temp} = Ind_{Loc} * (Op_A^{mf})$

Upshot: we have

$$KL(A) \xrightarrow{FLE} Ind_{Loc} * (Op_A^{mf})$$

\downarrow Poinc

$$\begin{array}{ccc} Loc \downarrow & & \\ Dmod(Bun_A)^{temp} & \xrightarrow{\mathbb{L} = \mathbb{L}_{coarse}|^{temp}} & Qcoh(LS_A^v) \end{array}$$

$$\begin{array}{ccc} \downarrow & & \downarrow \\ Dmod(Bun_A) & \xrightarrow{\mathbb{L}} & Ind_{Nilp}^{Loc}(LS_A^v) \end{array}$$

$$\begin{array}{ccc} coet \downarrow & & \downarrow \Gamma Spec \\ Whit(A)_{Ran} & \xrightarrow{CS} & Rep(A^v)_{Ran} \end{array}$$

fully faithful

Need to show commutativity of

$$\begin{array}{ccc}
 KL(A) & \xrightarrow{\sim}^{FLE} & IndLoc_* (Op_{\check{A}}^{mf}) \\
 \downarrow Loc & & \downarrow p_{inc}^{Spec} \\
 Dmod(Bun_A)_{temp} & & QLoc(LS_{\check{A}}) \\
 \downarrow & & \downarrow \Gamma^{Spec} \\
 Whif(A)_{Ran} & \xrightarrow{\sim}^{CS} & Rep(\check{A})_{Ran} \\
 \downarrow \text{coet}^{vac} & & \downarrow \tau(\check{A}) \\
 Vect & \xlongequal{\quad} & Vect
 \end{array}$$

Lin $KL(A) \xrightarrow{Loc} Dmod(Bun_A) \xrightarrow{\text{coet}^{vac}} Vect$ is given by

$$KL(A) \xrightarrow{p_{Senh}} \mathfrak{z}\text{-mod fact} \xrightarrow{C^{fact}(x, \mathfrak{z}, -)} Vect$$

Kevin $IndLoc_* (Op_{\check{A}}^{mf}) \rightarrow QLoc(LS_{\check{A}}) \xrightarrow{\Gamma} Vect$ is given by

$$IndLoc_* (Op_{\check{A}}^{mf}) \xrightarrow{z_*} IndLoc_* (Op_{\check{A}}^{mer}) \rightarrow \mathfrak{z}\text{-mod fact} \xrightarrow{C^{fact}(x, \mathfrak{z}, -)} Vect$$

Point FLE is constructed so that

$$KL(A) \xrightarrow{FLE} IndLoc_* (Op_{\check{A}}^{mf}) \xrightarrow{i_*} IndLoc_* (Op_{\check{A}}^{mer})$$

$$\text{agrees w } KL(A) \rightarrow KM(A) \rightarrow Whif_*(KM) \xrightarrow{p_{Senh}} IndLoc_*(Op_{\check{A}}^{mer})$$


$$\text{so. } KL(A) \xrightarrow{FLE} IndLoc_* (Op_{\check{A}}^{mf})$$

$$\begin{array}{ccc}
 \downarrow & & \downarrow i_* \\
 Whif_*(KM) & \xrightarrow{p_{Senh}} & IndLoc_*(Op_{\check{A}}^{mer}) \longrightarrow \mathfrak{z}\text{-mod fact}
 \end{array}$$

commutes.

Lecture 7. (Justin Campbell) Eisenstein series and constant terms

$P \subset G$ parabolic, $P = MN_P$



Constant term: $CT_* : D_{\text{mod}}(\text{Bun}_M) \longrightarrow D_{\text{mod}}(\text{Bun}_N)$

$$CT_* = q_* P^! [\pm]$$

Prop CT_* admits a left adjoint $Eis^!$.

Ref. Braverman-Laitsgory, Drinfeld-Laitsgory

Remark can modify $Eis^!$, CT_* so that $D_{\text{mod}_{\text{crit}}}(\text{Bun}_M) \xrightleftharpoons[CT_*]{Eis^!} D_{\text{mod}_{\text{crit}}}(\text{Bun}_N)$

The cuspidal part

$D_{\text{mod}}(\text{Bun}_M)_{Eis} \subset D_{\text{mod}}(\text{Bun}_M)$ full subcat. gen. by images of $Eis_{P,*}^!$

for all proper parabolic subgrp. $P \subsetneq G$

$$D_{\text{mod}}(\text{Bun}_M)_{\text{cusp}} = \{ F \in D_{\text{mod}}(\text{Bun}_M) : CT_{P,*} F = 0, \forall P \subsetneq G \}$$

Theorem (Drinfeld-Laitsgory) \exists open substack $j: U \hookrightarrow \text{Bun}_M$ s.t.

1) U has quasi-cpt intersection w/ each conn. comp of Bun_M .

2) $\forall F \in D_{\text{mod}}(\text{Bun}_M)_{\text{cusp}}, j_* j^! F \simeq F \simeq j_* j^* F$.

The spectral side

$$\begin{array}{ccc} & LS_{\check{P}}^{\vee} & \\ p^{\text{Spec}} \swarrow & & \searrow q^{\text{Spec}} \\ LS_{\check{h}}^{\vee} & & LS_{\check{M}}^{\vee} \end{array}$$

q^{Spec} quasi-smooth $\Rightarrow q^{\text{Spec},*}$ exists (for IndCoh)

p^{Spec} proper $\Rightarrow p_{*}^{\text{Spec}}$ is left adjoint to $p^{\text{Spec},!}$

Prop (Arikin-Hartshorne)

$p_{*}^{\text{Spec}} q^{\text{Spec},*}$ and $q_{*}^{\text{Spec}} p^{\text{Spec},!}$ preserve nilpotent

singular supp.

Key point: if $\alpha \in g^{*}$ is s.t. its image in p^{*} lands in m^{*} and is nilp, then α is nilpotent.

$$E_{\text{is}}^{\text{Spec}} := p_{*}^{\text{Spec}} q^{\text{Spec},*} : \text{IndCoh}_{\text{Nilp}}(LS_{\check{M}}^{\vee}) \rightleftarrows \text{IndCoh}_{\text{Nilp}}(LS_{\check{h}}^{\vee}) : q_{*}^{\text{Spec}} p^{\text{Spec},!} =: C_T^{\text{Spec}}$$

Thm (Arikin-Hartshorne)

- 1) $\mathcal{O}\text{Coh}(LS_{\check{h}}^{\text{ired}}) \xrightarrow{\sim} \text{IndCoh}_{\text{Nilp}}(LS_{\check{h}}^{\text{ired}})$
- 2) $\text{IndCoh}_{\text{Nilp}}(LS_{\check{h}}^{\vee})$ is gen. by $\mathcal{O}\text{Coh}(LS_{\check{h}}^{\vee})$ and essential image $E_{\text{is}}^{\text{Spec}}(\mathcal{O}\text{Coh}(LS_{\check{M}}^{\vee}))$ for all $\check{P} \neq \check{h}$.



Spectral

Constant term via factorization homology

Local analogue of constant term:

$$\begin{array}{ccc} & LS_{\check{P}}^{\text{reg}} & \\ p_{\text{loc}}^{\text{Spec}} \swarrow & & \searrow q_{\text{loc}}^{\text{Spec}} \\ LS_{\check{h}}^{\text{reg}} & & LS_{\check{M}}^{\text{reg}} \end{array}$$

$$J^{Spec, !} := (q^{Spec})_* (p^{Spec})^* : \mathcal{Q}coh(LS_{\check{h}}^{reg}) \rightarrow \mathcal{Q}coh(LS_{\check{m}}^{reg})$$

More concretely, $J^{Spec, !} : \text{Rep}(\check{h}) \xrightarrow{\text{res}_{\check{p}}^{\check{h}}} \text{Rep}(\check{p}) \xrightarrow{\text{inv}_{\check{M}\check{p}}} \text{Rep}(\check{m})$

lax unital factorization functor.

$$\begin{array}{ccc} \text{Rep}(\check{h}) & \xrightarrow{J^{Spec, !}} & \text{Rep}(\check{m}) \\ \text{Loc}_{\check{h}}^{Spec} \downarrow & \eta \swarrow & \downarrow \text{Loc}_{\check{m}}^{Spec} \\ \text{Ind Coh}_{\text{Miep}}(LS_{\check{h}}^{\vee}) \otimes \underline{Vect} & \xrightarrow{CT^{Spec} \otimes \text{id}} & \text{Ind Coh}_{\text{Miep}}(LS_{\check{m}}^{\vee}) \otimes \underline{Vect} \end{array}$$

$\text{Prop} \exists$ nat'l trans. η of lax unital local-to-global functors which induces

$$(Loc_{\check{m}}^{Spec} \circ J^{Spec, !})^{\text{ins. unit}} \xrightarrow{\sim} (CT^{Spec} \otimes \text{id}) \circ Loc_{\check{h}}^{Spec}$$

Proof (sketch) By def'n.

$$\begin{array}{ccc} \text{Rep}(\check{h}) & & \\ \text{Loc}_{\check{h}}^{Spec} \downarrow & \searrow \text{Loc}_{\check{h}}^{Spec} & \\ & \cong & \mathcal{Q}coh(LS_{\check{h}}^{\vee}) \otimes \underline{Vect} \\ & \swarrow & \\ & & \text{Ind Coh}_{\text{Miep}}(LS_{\check{h}}^{\vee}) \otimes \underline{Vect} \end{array}$$

Atkin - waits going:

$$\begin{array}{ccc}
 \mathcal{O}_{\text{coh}}(LS_{\check{h}}) & \xrightarrow{CT_{\mathcal{O}_{\text{coh}}}^{\text{Spec}}} & \mathcal{O}_{\text{coh}}(LS_{\check{m}}) \\
 \downarrow & \swarrow & \downarrow \\
 \text{Ind Coh}_{\text{Nilp}}(LS_{\check{h}}) & \xrightarrow{CT^{\text{Spec}}} & \text{Ind Coh}_{\text{Nilp}}(LS_{\check{m}})
 \end{array}$$

$$CT_{\mathcal{O}_{\text{coh}}}^{\text{Spec}} := q_*^{\text{Spec}} p^{\text{Spec},*}$$

$$\begin{array}{ccc}
 \text{Rep}(\check{h}) & \xrightarrow{\text{res}_{\check{p} \rightarrow \check{h}}} & \text{Rep}(\check{p}) \\
 \downarrow & \swarrow p^{\text{Spec},*} \otimes \text{id} & \downarrow \\
 \mathcal{O}_{\text{coh}}(LS_{\check{h}}) \otimes \underline{\text{Vect}} & \longrightarrow & \mathcal{O}_{\text{coh}}(LS_{\check{p}}) \otimes \underline{\text{Vect}}
 \end{array}$$

Draw the same square, replacing $\check{p} \rightarrow \check{h}$ by $\check{p} \rightarrow \check{m}$. Then pass to right adjoints of the horizontal functors.

$$\begin{array}{ccc}
 \text{Rep}(\check{p}) & \xrightarrow{\text{inv}_{N_{\check{p}}}} & \text{Rep}(\check{m}) \\
 \downarrow & \swarrow & \downarrow \\
 \mathcal{O}_{\text{coh}}(LS_{\check{p}}) \otimes \underline{\text{Vect}} & \xrightarrow{q_*^{\text{Spec}} \otimes \text{id}} & \mathcal{O}_{\text{coh}}(LS_{\check{m}}) \otimes \underline{\text{Vect}}
 \end{array}$$

$$\text{WTS } (L_{\check{m}}^{\text{Spec}} \circ \text{inv}_{N_{\check{p}}})^{\text{ins. unit}} \simeq (q_*^{\text{Spec}} \otimes \text{id}) \circ L_{\check{p}}^{\text{Spec}}$$

Enough to show after base changing along $\sigma: \text{Spec } k \rightarrow LS_{\check{m}}$.

For simplicity, take $\sigma = \sigma_{\text{triv}}$.

Lecture 8 (Dennis Gaitsgory)

Goal of this talk

$$D_{\text{mod}}(Bun_M) \xrightarrow{\mathbb{L}_M} \text{Ind Coh}_{\text{nilp}}(LS_M^\vee)$$

$$\downarrow \text{Eis!} \uparrow \text{CT}^*$$

$$\downarrow \text{Eis}^{\text{Spec}} \uparrow \text{CT}^{\text{Spec}} + \text{induction step}$$

$$\begin{array}{ccccc} D_{\text{mod}}(Bun_A) & \xrightarrow{\mathbb{L}_A} & \text{Ind Coh}_{\text{nilp}}(LS_A^\vee) & & \\ \text{Whit}^!(\omega_M)_{\text{Ran}} & \xrightarrow{\quad} & \text{Rep}(X)_{\text{Ran}} & & \\ \text{coJ}^{\text{fix}^\vee} \downarrow & & \uparrow \text{ind}(\text{fix}^\vee) & & \\ D_{\text{mod}}(Bun_M) & \xrightarrow{\text{Whit}^!(\omega_M)_{\text{Ran}} \mathbb{L}_{A, \text{coarse}}} & \text{Coh}(LS_M^\vee) & \xrightarrow{\quad} & \text{Rep}(X)_{\text{Ran}} \\ \text{Eis!} \searrow & \text{coJ} \uparrow & \text{Eis}^{\text{Spec, coarse}} \uparrow & \Gamma_A^{\text{Spec}} & \\ D_{\text{mod}}(Bun_A) & \xrightarrow{\mathbb{L}_{M, \text{coarse}}} & \text{Coh}(LS_A^\vee) & & \end{array}$$

Need $\text{coJ} : \text{Whit}^!(\omega_M) \rightarrow \text{Whit}^!(\omega_A)$ so that

- the upper lid commutes
- the left lid commutes

Define $\text{Whit}^!(\omega_M) \xrightarrow{\text{CS}_M} \text{Rep}(M^\vee)$

$$\begin{array}{ccc} \text{coJ} \downarrow & & \downarrow \text{ind} \\ \text{Whit}^!(\omega_A) & \xrightarrow{\text{CS}_A} & \text{Rep}(A^\vee) \end{array}$$

$$\text{Whit}^!(\omega_A) \xrightarrow{\text{CS}_A} \text{Rep}(A^\vee)$$

$$\begin{array}{ccc} \downarrow \text{J} & & \downarrow \text{inv}_{N_P} \\ \text{Whit}^!(\omega_M) & \xrightarrow{\text{CS}_M} & \text{Rep}(M^\vee) \end{array}$$

Thm (Sith) The diagram commutes.

$$\text{Whit}^!(\omega_a) \otimes \text{Dmod}(\omega_a)^{LN_{p-1} \cdot L^+ M} \longrightarrow \text{Whit}^!(\omega_m)$$

↓

$$\text{Whit}^!(\omega_a) \otimes \text{Dmod}(\omega_a \times \omega_m)^{LP^-}$$

↓

$$\text{Dmod}(\omega_a) \otimes \text{Dmod}(\omega_a) \otimes \text{Dmod}(\omega_m) \xrightarrow{\langle, \rangle_{\omega_a} \otimes \text{Id}} \text{Dmod}(\omega_m)$$

↓ \langle, \rangle

Verf



$$\Delta^{-, \frac{\infty}{2}} = \text{is} \cdot \omega_s^-$$

$$\bar{S} \xrightarrow{i} \omega_a$$

||


$$LN_{p-1}$$

$$\overline{\text{Bun}}_p^- \times \text{Bun}_a^{\Omega(u)} \xleftarrow{\quad} \text{Zast}$$



$$- J_{KM} = \text{BRST} = \left(\frac{\omega}{2} (L\eta_p, -) \right)$$

- J_{Op} = pull-push



$$\begin{array}{ccc} KL(M) & \xrightarrow{FLE_M} & \text{Indoh}^*(O_{P\check{m}}^{mf}) \\ \uparrow J_{KM} & & \uparrow J_{Op} \\ KL(a) & \xrightarrow{FLE_h} & \text{Indoh}^*(O_{P\check{h}}^{mf}) \end{array}$$

Bottom lid VERY HARD

Cor. Δ_a admits a left adjoint.

Proof.

$$C \xrightarrow{F} D$$

WTS that $(\Delta_a)^L$ is defined on a generating set of objects.

$$\begin{array}{ccc} C_1 & \xrightarrow{F_1} & D_1 \\ i_C \uparrow & & \uparrow i_D \\ C & \xrightarrow{F} & D \end{array} \quad \begin{array}{ccc} C_1 & \xleftarrow{F_1^L} & D_1 \\ i_C^L \downarrow & & \downarrow i_D^L \\ C & \xleftarrow{F^L} & D \end{array}$$

Lemma F^L is defined on the essential image of i_D^L so that commutes

$$\begin{array}{ccc} \text{Whit}^!(\omega_a) & \xrightarrow[\sim]{CS} & \text{Rep}(\check{h})_{\text{Ran}} \\ \text{Point}_a \downarrow \text{coet}_a \uparrow & & \text{Loc}_{\check{h}}^{\text{Spec}} \uparrow \uparrow \text{Spec } \check{h} \\ \text{Dmod}(\text{Bun}_a) & \xrightarrow{\Delta_a} & \text{Indoh}_{\text{Nilp}}(LS_a^{\vee}) \end{array}$$

page 44

Obtain Π_G^L is defined on $\mathcal{O}(\text{Loc}) (LS_G^\vee) \subset \text{IndLoc}_{\text{Nrep}} (LS_G^\vee)$

$$\begin{array}{ccc} \text{Whit}! (W_G) & \xrightarrow{LS_G^\vee} & \text{Rep}(\check{G})_{\text{Ran}} \\ \text{Poinc}! \downarrow & & \downarrow \text{Loc}^{\text{Spec}}_{\check{G}} \\ \text{Dmod}(Bun_G) & \xleftarrow{\Pi_G^L} & \text{IndLoc}_{\text{Nrep}} (LS_G^\vee) \end{array}$$

$$\begin{array}{ccc} \text{Dmod}(Bun_M) & \xrightarrow{\Pi_M} & \text{IndLoc}_{\text{Nrep}} (LS_M^\vee) \\ \text{Eis}! \downarrow \uparrow \text{CT}_\alpha & & \text{Eis}^{\text{Spec}} \downarrow \uparrow \text{CT}^{\text{Spec}} \\ \text{Dmod}(Bun_G) & \xrightarrow{\Pi_G} & \text{IndLoc}_{\text{Nrep}} (LS_G^\vee) \end{array}$$

Obtain Π_G^L is defined on the essential image of Eis^{Spec} and

$$\begin{array}{ccc} \text{Dmod}(Bun_M) & \xleftarrow{\Pi_M^L} & \text{IndLoc}_{\text{Nrep}} (LS_M^\vee) \\ \text{Eis}! \downarrow & & \downarrow \text{Eis}^{\text{Spec}} \\ \text{Dmod}(Bun_G) & \xleftarrow{\Pi_G^L} & \text{IndLoc}_{\text{Nrep}} (LS_G^\vee) \end{array}$$

Lemma [AG] $\text{IndLoc}_{\text{Nrep}} (LS_G^\vee)$ is generated by $\mathcal{O}(\text{Loc}) (LS_G^\vee)$ and Eis^{Spec} .

Def $\text{Dmod}(Bun_G)_{\text{Eis}} = \langle \text{Eis}!, \text{proper parabolics} \rangle$

$$\text{Dmod}(Bun_G)_{\text{Eis}} \xrightleftharpoons{\quad} \text{Dmod}(Bun_G) \xrightleftharpoons[e]{e^L} \text{Dmod}(Bun_G)_{\text{cusp}}$$

Def $\text{Ind Coh}_{\text{nilp}}(LS_X)_{Eis} = \langle Eis^{\text{Spec}}, \text{proper parabolics} \rangle$

$$\text{Ind Coh}_{\text{nilp}}(LS_X^v)_{Eis} \begin{matrix} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{matrix} \text{Ind Coh}_{\text{nilp}}(LS_X^v) \begin{matrix} \xrightarrow{(e^{\text{Spec}})^L} \\ \xleftarrow{e^{\text{Spec}}} \end{matrix} \text{Ind Coh}_{\text{nilp}}(LS_X^v)_{\text{cusp}}$$

$$\parallel$$

$$\text{Ind Coh}_{\text{nilp}}(LS_X^v)_{\text{red}}$$

$$\parallel$$

$$\text{Ind Coh}_{\text{nilp}}(LS_X^{\text{ired}})$$

$$e^{\text{Spec}} = (j^{\text{Spec}})_*$$

$$(e^{\text{Spec}})^L = (j^{\text{Spec}})^*$$

$$\parallel$$

$$\text{Coh}(LS_X^{\text{ired}})$$

$$\text{Dmod}(\text{Bun}_G)_{Eis} \begin{matrix} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{matrix} \text{Dmod}(\text{Bun}_G) \begin{matrix} \xrightarrow{e^L} \\ \xleftarrow{e} \end{matrix} \text{Dmod}(\text{Bun}_G)_{\text{cusp}}$$

$$\mathbb{L}_{G,Eis}^L \uparrow \downarrow \mathbb{L}_{G,Eis}$$

$$\mathbb{L}_G^L \uparrow \downarrow \mathbb{L}_G$$

$$\uparrow \downarrow$$

$$\text{Ind Coh}_{\text{nilp}}(LS_X^v)_{\text{red}} \begin{matrix} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{matrix} \text{Ind Coh}_{\text{nilp}}(LS_X^v)$$

$$\text{Ind Coh}_{\text{nilp}}(LS_X^v)$$

$$\begin{matrix} \xrightarrow{(j^{\text{Spec}})^*} \\ \xleftarrow{(j^{\text{Spec}})_*} \end{matrix}$$

$$\text{Coh}(LS_X^{\text{ired}})$$

Thm $(\mathbb{L}_{G,Eis}^L, \mathbb{L}_{G,Eis})$ are mutually inverse equivalences.

$$Eis^{\text{Spec}} \longrightarrow \mathbb{L}_{G,Eis} \cdot \mathbb{L}_{G,Eis}^L \cdot Eis^{\text{Spec}}$$

SI

$$\mathbb{L}_G \cdot Eis ! \cdot \mathbb{L}_M^L$$

IS

$$Eis^{\text{Spec}} \longrightarrow Eis^{\text{Spec}} \cdot \mathbb{L}_M \cdot \mathbb{L}_M^L$$

$$\begin{array}{ccc}
 \text{Whit}'(W_n)_{\text{Ran}} & \cong & \text{Rep}(\check{G})_{\text{Ran}} \\
 \uparrow & & \uparrow \vdash \text{Spec } \check{G} \\
 D_{\text{mod}}(Bun_n) & & \text{Indlo}_{\text{nilp}}(LS_{\check{G}}) \\
 \uparrow e & & \uparrow (\check{J}^{\text{Spec}})_* \\
 D_{\text{mod}}(Bun_n)_{\text{cusp}} & \longrightarrow & \mathcal{O}lo_{\check{G}}(LS_{\check{G}}^{\text{inv}})
 \end{array}$$

For $G = GL_n$

Bonus Material D (Johan Fargnason)

Recall $D(Bun_n)_{\text{Eis}} \cong \text{Indlo}_{\check{G}}(LS_{\check{G}})_{\text{Eis}}$

leaves $D(Bun_n)_{\text{cusp}} \xrightarrow{\mathbb{L}_{G, \text{cusp}}} \mathcal{O}lo_{\check{G}}(LS_{\check{G}}^{\text{inv}})$

Goal. $\mathbb{L}_{G, \text{cusp}}$ is conservative: $\mathbb{L}_{G, \text{cusp}}(F) = 0 \Rightarrow F = 0$.

In fact, we'll prove that $\mathbb{L}_{G, \text{temp}}: D(Bun_n)_{\text{temp}} \rightarrow \mathcal{O}lo_{\check{G}}(LS_{\check{G}}^{\text{inv}})$ is conservative.

Rmk. 1) To prove hLC, it remains to show that $\mathbb{L}_{G, \text{temp}}^L$ is fully faithful.

2) The above proof is microlocal in nature.

Reduction Claim. It suffices to show that $\mathbb{L}_{G, \text{temp}}: D_{\text{nilp}}(Bun_n)_{\text{temp}} \rightarrow \mathcal{O}lo_{\check{G}}(LS_{\check{G}}^{\text{rest}})$ is conservative.

Pf Given $F \in D(Bun_n)_{\text{temp}}$, $\exists \sigma: \text{Spec } k \rightarrow LS_{\check{G}}^{\text{inv}}$, $\sigma \otimes F \neq 0$.

$$\sigma: \text{Spec } k \rightarrow \text{LS}_{\check{h}}^{\text{rest}} \rightarrow \text{LS}_{\check{h}}^{\vee}$$

$$\text{so } \sigma \otimes F \in D(\text{Bun}_{\check{h}}) \otimes_{\mathcal{O}(\text{LS}_{\check{h}}^{\vee})} \mathcal{O}(\text{LS}_{\check{h}}^{\text{rest}}) \xrightarrow{[\text{AKRRV}]} D_{\text{nilp}}(\text{Bun}_{\check{h}})$$

$$\text{If } 0 \neq \mathbb{L}_{\check{h}, \text{temp}}(\sigma \otimes F) = \sigma \otimes \underbrace{\mathbb{L}_{\check{h}, \text{temp}}(F)}_{\neq 0}$$

going to explicitly describe $\text{Ker}(\mathbb{L}_{\check{h}, \text{coarse}} = D_{\text{nilp}}(\text{Bun}_{\check{h}}) \rightarrow \mathcal{O}(\text{LS}_{\check{h}}^{\text{rest}}))$

Def'n $x \in g^* \approx g$ regular if $(g(x))$ minimal dim ($= \text{rk}(g)$)

Better for us: fix some non-degenerate character $\psi \in \Pi^*$,

consider $g^*/N = g^*/_{\check{h}} \times_{\text{pt}/\check{h}} \text{pt}/N$

Define Kos_{ψ} as sitting in a Cartesian diagram:

$$\begin{array}{ccc} \text{Kos}_{\psi} & \longrightarrow & \psi/N \\ \downarrow & \lrcorner & \downarrow \\ g^*/N & \longrightarrow & \Pi^*/N \end{array}$$

Identifying $g^* \approx g$, $\Pi^* \approx \Pi$, $\psi = \mathbf{f}$, then $\text{Kos}_{\psi} \approx \mathbf{f} + \mathbf{b}/N$ ($\approx \mathbf{f} + g^e$)

The map $\mathbf{f} + \mathbf{b}/N \rightarrow g_{\text{reg}}/\check{h}$ bijection on k -pts

$$g_{\text{irreg}} = g \setminus g_{\text{reg}}, \quad N_{\text{irreg}} = N \cap g_{\text{irreg}}$$

Globally Define $N_{\text{irreg}} \subset T^* \text{Bun}_{\check{h}}$ to be pairs

$(p_{\check{h}}, \psi \in \Gamma(X, g_{p_{\check{h}}}^* \otimes \Omega_X^1))$ s.t. ψ is nilp. irreg. (ie. locally factors through N_{irreg})

$$\Gamma_{T_{P_h} \text{Bun}_h} = H^*(X, g_{P_h})[1] \quad \text{Using Serre Duality,}$$

$$H^0 T_{P_h}^* \text{Bun}_h = \Gamma(X, g_{P_h}^* \otimes \omega_X')$$

┘

Thm A $\text{Ker}(\mathbb{L}_{h, \text{coarse}} : D_{\text{Nilp}}(\text{Bun}_h) \rightarrow D_{\text{Nilp}_{\text{ineg}}}(\text{Bun}_h)) = D_{\text{Nilp}_{\text{ineg}}}(\text{Bun}_h)$

Thm B $D_{\text{Nilp}}(\text{Bun}_h)_{\text{temp}} \cap D_{\text{Nilp}_{\text{ineg}}}(\text{Bun}_h) = 0.$

$G = SL_2$: $\text{Nilp}_{\text{ineg}} = \{0\}$

$$D_{\text{Nilp}_{\text{ineg}}}(\text{Bun}_h) = \{F \in D(\text{Bun}_h) : H^i(F) = 0 \text{ (ext's of } w)\}$$

$$\begin{array}{ccc} & \mathbb{L}_{h, \text{coarse}} & \text{Coh}(LS_X) \\ & \nearrow & \downarrow \Gamma \end{array}$$

Thm A $\text{coet}_0 : D(\text{Bun}_h) \rightarrow \text{Vect}$

$$\begin{array}{ccc} \text{Bun}_N^{\Omega} & \xrightarrow{P} & \text{Bun}_h \\ \psi \downarrow & & \\ \mathbb{A}^1 & & \end{array} \quad \text{coet}_0(F) = \text{Car}(\text{Bun}_N^{\Omega}, p^! F \otimes \psi^! (\exp))$$

Thm 1A $\text{coet}_0|_{D_{\text{Nilp}}(\text{Bun}_h)}$ satisfies

- 1) t-exact
- 2) commutes w/ Verdier duality
- 3) $F \in D_{\text{Nilp}}(\text{Bun}_h)^b$, then $\dim \text{coet}_0(F) = \text{mult}(\text{Nilp}^{\text{Kos}}, \text{cc}(F))$

Defining Nilp^{Kos} :

$$\psi: \text{Bun}_N^{\Omega} \rightarrow \mathbb{A}^1$$

$$\sim d\psi: \text{Bun}_N^{\Omega} \rightarrow T^* \text{Bun}_N^{\Omega}$$

Define $\text{Kos}_{\psi}^{\text{glob}}$ by

$$\begin{array}{ccc} \text{Kos}_{\psi}^{\text{glob}} & \xrightarrow{\quad} & \text{Bun}_N^{\Omega} \\ \downarrow & \lrcorner & \downarrow d\psi \\ T^* \text{Bun}_N \times_{\text{Bun}_N} \text{Bun}_N^{\Omega} & \xrightarrow{dp} & T^* \text{Bun}_N^{\Omega} \end{array}$$

Key properties

$$\text{Kos}_{\psi} \times_{g^*/h} \mathcal{N}/h = \{f\}$$

$$\text{Kos}_{\psi}^{\text{glob}} \times_{T^* \text{Bun}_N} \text{Nilp} = \{f^{\text{glob}}\}$$

Define $\text{Nilp}^{\text{Kos}} \subset \text{Nilp}$ the irred. comp. containing f^{glob} .

$G = \text{SL}_2$

$$\text{Nilp} = \bigcup_{d \geq 1-g} \overline{\text{Nilp}^d}$$

$$\text{deg } L = d$$

$$\text{Nilp}^d = \left\{ 0 \rightarrow L \rightarrow \mathcal{E} \rightarrow L^{\vee} \rightarrow 0, \varphi: L^{\vee} \rightarrow L \otimes \mathcal{N} \right\}_{0 \neq}$$

$$\mathcal{E} \rightarrow L^{\vee} \xrightarrow{\varphi} L \otimes \mathcal{N}'_x \rightarrow \mathcal{E} \otimes \mathcal{N}'_x$$

$$\text{Nilp}^{1-g} = \text{Nilp}^{\text{kos}}$$

$$f^{gbb} = (\Omega^{-1/2} \oplus \Omega^{1/2}, \quad \varphi: \Omega^{-1/2} \oplus \Omega^{1/2} \rightarrow (\Omega^{-1/2} \oplus \Omega^{1/2}) \otimes \Omega \\ = \Omega^{1/2} \oplus \Omega^{3/2})$$

$$\varphi = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

Thm 2A If $F \in D_{\text{Nilp}}(\text{Bun}_h)$ s.t. $SS(F) \notin \text{Nilp}_{\text{reg}}$, then $\exists \check{\lambda} \in \check{\Lambda}^+$

$$SS(H_{V_x^{\check{\lambda}}} + F) \supset \text{Nilp}^{\text{kos}}$$

$$\forall x \in X$$

Exer formulate and prove analogous result for $D_X(g/h)$ using Fourier transform

$$\text{Thm 1A} + 2A \Rightarrow \text{Thm A.}$$

\uparrow

$$D_{\text{Nilp}_{\text{reg}}}(\text{Bun}_h) = \text{Ker}(\mathbb{L}_{h, \text{coarse}} | D_{\text{Nilp}}(\text{Bun}_h))$$

" \subset " If $F \in D_{\text{Nilp}_{\text{reg}}}(\text{Bun}_h)$, we get that $\text{coeth}_0(H_{V_x^{\check{\lambda}}} + F) = 0$

$$\text{This means } \Gamma(LS_{\check{h}}^{\text{rest}}, \mathbb{L}_{h, \text{coarse}}(F) \otimes E_x^{\check{\lambda}}) = 0$$

$$\Rightarrow \mathbb{L}_{h, \text{coarse}}(F) = 0 \text{ because } LS_{\check{h}}^{\text{rest}} \text{ is a formal affine scheme.}$$

" \supset " Given $F \in \text{Ker}(\quad)$, $\Gamma(LS_{\check{h}}^{\text{rest}}, E_x^{\check{\lambda}} \otimes \mathbb{L}_{h, \text{coarse}}(F)) = 0, \forall \check{\lambda}$

$$\parallel \\ \text{coeth}_0(H_{V_x^{\check{\lambda}}} + F)$$

$$\Rightarrow SS(F) \subset \text{Nilp}_{\text{reg}} \quad \square$$

$$\text{Thm A} + \text{Thm B} \Rightarrow \ker(\mathbb{L}_A, \text{coarse} \mid D_{\text{nilp}}(B_{\text{un}})) \cap D_{\text{nilp}}(B_{\text{un}})_{\text{temp}} = \emptyset.$$

Lecture 9 (Dennis Gaitsgory) Ambidexterity - I

$$\begin{array}{ccc}
 D_{\text{mod}}(B_{\text{un}}) \xrightleftharpoons[\text{Eis}]{\text{Eis}} D_{\text{mod}}(B_{\text{un}}) & \xrightleftharpoons[e]{e^L} & D_{\text{mod}}(B_{\text{un}})_{\text{cusp}} \\
 \mathbb{L}_A^L \uparrow \downarrow \mathbb{L}_A & & \mathbb{L}_{A, \text{cusp}}^L \uparrow \downarrow \mathbb{L}_{A, \text{cusp}} \\
 \text{Ind}_{\text{nilp}}^{\text{Loh}}(LS_A^{\vee}) \xrightleftharpoons[\text{red}]{\text{red}} \text{Ind}_{\text{nilp}}^{\text{Loh}}(LS_A^{\vee}) & \xrightleftharpoons[(j^{\text{spec}})^*]{(j^{\text{spec}})^*} & \text{Ind}_{\text{nilp}}^{\text{Loh}}(LS_A^{\text{ired}}) \\
 & & \text{"} \\
 & & \text{Loh}(LS_A^{\text{ired}})
 \end{array}$$

$$\mathbb{L}_{A, \text{temp}} : D_{\text{mod}}(B_{\text{un}})_{\text{temp}} \rightarrow \text{Loh}(LS_A^{\vee}) \text{ is conservative}$$

\Downarrow

$$\mathbb{L}_{A, \text{cusp}} : D_{\text{mod}}(B_{\text{un}})_{\text{cusp}} \rightarrow \text{Loh}(LS_A^{\text{ired}}) \text{ is conservative}$$

Goal to show $\mathbb{L}_{A, \text{cusp}}^L$ is fully faithful.

$$\mathbb{L}_{A, \text{temp}} \circ \mathbb{L}_{A, \text{temp}}^L : \text{Loh}(LS_A^{\vee}) \rightarrow \text{Loh}(LS_A^{\vee})$$

SI

$$A_A \otimes - , \quad A_A \in \text{Assoc Alg}(\text{Loh}(LS_A^{\vee}))$$

$$\mathbb{L}_{A, \text{cusp}} \circ \mathbb{L}_{A, \text{cusp}}^L$$

"

$$A_{A, \text{cusp}} \otimes - ,$$

$$A_{A, \text{cusp}} = (j^{\text{spec}})^*(A_A)$$

Fully-faithfulness $\Leftrightarrow \mathcal{O}_{LS_h^*} \Rightarrow A_h$ is an isom.

on cusp part $\Leftrightarrow \mathcal{O}_{LS_h^{\text{ined}}} \Rightarrow A_{h, \text{cusp}}$ is an isom.

Know already: $(\iota^{\text{spec}})^* (\mathcal{O}_{LS_h^*}) \Rightarrow (\iota^{\text{spec}})^* (A_h)$

Thm (Ambidexterity) $\mathbb{L}_{h, \text{cusp}}$ is ambidexterous $\Leftrightarrow \mathbb{L}_{h, \text{cusp}}^L \cong \mathbb{L}_{h, \text{cusp}}^R$

eg. $\text{Vect} \xrightarrow{\otimes V} \text{Vect}$

Cor $A_{h, \text{cusp}}$ is perfect $\in \text{Qcoh}(LS_h^{\text{ined}})$ and self-dual.

$$A_{h, \text{cusp}} \otimes - = \mathbb{L}_{h, \text{cusp}} \mathbb{L}_{h, \text{cusp}}^L$$

$$(\mathbb{L}_{h, \text{cusp}} \cdot \mathbb{L}_{h, \text{cusp}}^L)^R = (\mathbb{L}_{h, \text{cusp}}^L)^R \cdot \mathbb{L}_{h, \text{cusp}}^R = \mathbb{L}_{h, \text{cusp}} \cdot \mathbb{L}_{h, \text{cusp}}^L$$

$$C^V = \text{Func}_{\text{cts}}(C, \text{Vect})$$

$$C = \text{Ind}(C^c)$$

$$C^V = \text{Ind}((C^c)^{\text{op}})$$

$$C \otimes C^V \xrightarrow{\text{ev}} \text{Vect}$$

$$\text{unit}_C \in C \otimes C^V$$

$$C \xrightarrow{F} D$$

$$C^V \xleftarrow{F^V} D^V$$

$$C \begin{array}{c} \xrightarrow{F} \\ \xleftarrow{F^R} \end{array} D$$

F preserves compactness

$$C^c \xrightarrow{F} D^c$$

$$(C^c)^{op} \xrightarrow{F^{op}} (D^c)^{op}$$

Lemma (F^{op}, F^v) is an adjoint pair.

$$\left. \begin{array}{c} \} \\ C^v \xrightarrow{F^{op}} D^v \end{array} \right\}$$

$$ID_C : C^c \rightarrow (C^v)^c$$

$$F^{op} = ID_D \circ F \circ ID_C$$

Example $C = D_{mod}(\mathcal{Y})$, \mathcal{Y} - qc alg stack

$$D_{mod}(\mathcal{Y}) \otimes D_{mod}(\mathcal{Y}) \xrightarrow{\langle, \rangle} Vect$$

$$F_1 \quad F_2 \quad \longmapsto \text{CIR}(\mathcal{Y}, F_1 \overset{!}{\otimes} F_2)$$

$$\Delta_*(W_{\mathcal{Y}}) \in D_{mod}(\mathcal{Y} \times \mathcal{Y}) \simeq D_{mod}(\mathcal{Y}) \otimes D_{mod}(\mathcal{Y})$$

$$(D_{mod}(\mathcal{Y})^c)^{op} = D_{mod}(\mathcal{Y})^c, \quad ID_{\mathcal{Y}}$$

$$\mathcal{Y}_1 \xrightarrow{f} \mathcal{Y}_2$$

$$f_* : D_{mod}(\mathcal{Y}_1) \rightarrow D_{mod}(\mathcal{Y}_2)$$

$$(f_*)^v : D_{mod}(\mathcal{Y}_2) \rightarrow D_{mod}(\mathcal{Y}_1)$$

$$\parallel$$

$$f^!$$

$$(f_*)^{op} = f^! \simeq f_* \quad , \quad f \text{ proper}$$

$$f \text{ smooth}, \quad (f^!)^{op} = f^*$$

U c Bung

We obtain $D_{\text{mod}}(u)$ is self-dual.

Take U to be large enough so that $F \in \text{Dmod}(\text{Bun}_n)_{\text{cusp}} \Rightarrow F \xrightarrow{\sim} j_* j^* F$.

$$C_1 \xrightleftharpoons[e]{e^L} C$$

$$c_1^{\nu} \xrightleftharpoons[(e^L)^{\nu}]{e^{\nu}} c^{\nu}$$

$$D_{\text{mod}}(\text{Bun}_G)_{\text{usp}} \begin{array}{c} \xleftarrow{(e_u)^2} \\ \xrightarrow{e_u} \end{array} D_{\text{mod}}(u) \xrightarrow{\hat{J}_*} D_{\text{mod}}(\text{Bun}_G)$$

$$D_{\text{mod}}(Bun_n)_{\text{usp}}^v \begin{matrix} \xleftarrow{(e_n)^v} \\ \xrightarrow{(e_n^L)^v} \end{matrix} D_{\text{mod}}(n)$$

Lemma $D^{\text{mod}}(\text{Bun}_G)_{\text{cusp}}^{\vee} = D^{\text{mod}}(\text{Bun}_G)_{\text{cusp}}$

$\uparrow \quad \uparrow$
 $D^{\text{mod}}(U)$

$$D^{\text{mod}}(\text{Bun}_n)_{\text{usp}} \subset D^{\text{mod}}(U) \text{ can be realized as } (j^*, E_i!)^{\perp}$$
$$(D^{\text{mod}}(\text{Bun}_n)_{\text{usp}})^{\vee} \subset D^{\text{mod}}(U) \quad \text{can be realized as } (ID_U \cdot (j^* \cdot \text{Fis}) \cdot ID_{\text{Bun}_n})^{\perp}$$

$$\begin{array}{ccc} & \text{Bun}_P & \\ \swarrow p & & \searrow q \\ \text{Bun}_G & & \text{Bun}_M \end{array}$$

$$E(x) = p \cdot q^x, \quad E(x) = p_x \cdot q^x$$


Claim $E(i)_!$ and $E(i)_*$ after composing w/ j^* are finite extensions of copies of each other

$$D_{\text{mod}}(\text{Bun}_G)_{\text{cusp}} \xrightarrow{\mathbb{L}_{G, \text{cusp}}} \text{Coh}(LS_{\check{G}}^{\text{red}})$$

$$D_{\text{mod}}(\text{Bun}_G)_{\text{cusp}} \xleftarrow{\mathbb{L}_{G, \text{cusp}}^{\vee}} \text{Coh}(LS_{\check{G}}^{\text{red}})$$

Thm 1 $\mathbb{L}_{G, \text{cusp}}^{\vee} = (\mathbb{L}_{G, \text{cusp}})^L$

Thm 2 $\mathbb{L}_{G, \text{cusp}}^{\vee} = (\mathbb{L}_{G, \text{cusp}})^R$



$$HN^n = \{ 0 \rightarrow L \rightarrow E \rightarrow L^{\vee} \rightarrow 0 \mid \deg L = n > 0 \}$$

$$\mathbb{L}_{G, \text{cusp}}^L \simeq (\mathbb{L}_{G, \text{cusp}}^L)^{\circ P} \Leftrightarrow \text{Thm 1}$$

$$n \gg 0, H^1(L^2) = 0.$$

Σ HN^n unip gerbe over $\text{Bun}_{G_m}^n, n \gg 0$

$\Sigma_{\text{ex}} CT_{\Sigma}^{\vee} = ! - \text{rest. to } HN^n \text{ for } n \gg 0$

$$\begin{array}{ccc} \text{Whit}^!(G_m)_{\text{Ran}} & \xrightarrow{CS_G} & \text{Rep}(\check{G})_{\text{Ran}} \\ \uparrow \text{cleft} & & \uparrow \Gamma_{\check{G}}^{\text{spec}} \\ D_{\text{mod}}(\text{Bun}_G) & & \text{IndCoh}_{\text{Miep}}(LS_{\check{G}}^{\vee}) \\ \uparrow e & & \uparrow (\text{ispec})_{\Sigma} \\ D_{\text{mod}}(\text{Bun}_G)_{\text{cusp}} & \xrightarrow{\mathbb{L}_{G, \text{cusp}}} & \text{Coh}(LS_{\check{G}}^{\text{red}}) \end{array}$$

$$\begin{array}{ccc} & \text{Rep}(\check{G})_{\text{Ran}} & \\ & \uparrow \Gamma_{\check{G}}^{\text{spec}} \cdot (\text{ispec})_{\Sigma} & \\ CS \cdot \text{cleft} \cdot e & \nearrow & \\ D_{\text{mod}}(\text{Bun}_G)_{\text{cusp}} & \xrightarrow{\mathbb{L}_{G, \text{cusp}}} & \text{Coh}(LS_{\check{G}}^{\text{red}}) \end{array}$$

$$\begin{array}{ccc}
 & \text{Vect} & \\
 \begin{array}{c} \downarrow \\ ((\text{coett}^{\text{vac}}, e)^L)^{\text{op}} \end{array} & \begin{array}{c} \downarrow \\ ((\Gamma(LS_{\check{A}}, -) \cdot j_x^{\text{Spec}})^L)^{\text{op}} \end{array} & \begin{array}{c} \downarrow \\ (\text{coett}^{\text{vac}}, e)^L \end{array} \\
 \text{Dmod}(\text{Bun}_{\check{A}})_{\text{cusp}} & \xleftarrow{(\mathbb{L}_{\check{A}}, \text{cusp})^{\text{op}}} \mathcal{O}(\text{coh}(LS_{\check{A}}^{\text{inred}})) & \xleftarrow{(\mathbb{L}_{\check{A}}, \text{cusp})^L} \mathcal{O}(\text{coh}(LS_{\check{A}}^{\text{inred}}))
 \end{array}$$

$$\begin{array}{c}
 \Gamma(LS_{\check{A}}, -) \cdot j_x^{\text{Spec}} \\
 \parallel \\
 \mathcal{O}_{LS_{\check{A}}^{\text{inred}}}
 \end{array}
 \begin{array}{c}
 \downarrow \\
 \Gamma(LS_{\check{A}}, -) \cdot j_x^{\text{Spec}}
 \end{array}
 = (j_x^{\text{Spec}})^L \cdot \Gamma(LS_{\check{A}}, -)^L = (j^{\text{Spec}})^* \cdot \mathcal{O}_{LS_{\check{A}}^{\text{inred}}}$$

$$\begin{array}{c}
 \Gamma(LS_{\check{A}}, -) \cdot j_x^{\text{Spec}} \\
 \parallel \\
 \mathcal{O}_{LS_{\check{A}}^{\text{inred}}}
 \end{array}
 \begin{array}{c}
 \downarrow \\
 \Gamma(LS_{\check{A}}, -) \cdot j_x^{\text{Spec}}
 \end{array}
 = (j_x^{\text{Spec}})^V \cdot \Gamma(LS_{\check{A}}, -)^V = (j^{\text{Spec}})^* \cdot \mathcal{O}_{LS_{\check{A}}^{\text{inred}}}$$

$$(\text{coett}^{\text{vac}}, e)^L = e^L \cdot (\text{coett}^{\text{vac}})^L = e^L (\text{Poinc}^{\text{vac}}!) \in \text{Dmod}(\text{Bun}_{\check{A}})_{\text{cusp}}$$

$$((\text{coett}^{\text{vac}}, e)^L)^{\text{op}} \simeq (\text{coett}^{\text{vac}}, e)^L$$

$$\parallel \\
 e^L(\text{Poinc}^{\text{vac}}!)$$

$$\text{D}(\text{Bun}_{\check{A}}) \xrightarrow{j^*} \text{Dmod}(\mathcal{U})$$

$$e^L(\text{Poinc}^{\text{vac}}!) = e^L(\text{Poinc}^{\text{vac}}!)$$

$$e^L(\text{Poinc}^{\text{vac}}!) = e^L(\text{Poinc}^{\text{vac}}_*)$$

$$\begin{array}{ccc}
 & \text{Bun}_{\check{A}} & \\
 \swarrow p & & \searrow \text{ev} \\
 \text{Bun}_{\check{A}} & & \mathbb{A}^1
 \end{array}$$

Lecture 10 (Lin Chen)

Goal 1: $KL_{Ran} \xrightarrow{Loc} D(Bun_A) \xrightarrow{j^*} D(U)$ is a quotient For $U \xrightarrow{j} Bun_A$

Goal 2: $\mathbb{L}_{A, cusp}^V \xrightarrow{\sim} \mathbb{L}_{A, cusp}^R$

Goal 3: $\mathbb{R}_{A, cusp} \otimes - \stackrel{\text{def}}{=} \mathbb{L}_{A, cusp} \circ \mathbb{L}_{A, cusp}^L : \mathcal{Qcoh}(LS_{\check{X}}^{ired}) \rightarrow \mathcal{Qcoh}(LS_{\check{X}}^{ired})$
 $\sigma \in LS_{\check{X}}^{ired}, \quad \mathbb{R}_{A, cusp}|_{\sigma} = C. (\quad)$

Goal 1, General paradigm $F: \mathcal{C} \rightarrow \mathcal{D}$
 $F^R: \mathcal{D} \rightarrow \mathcal{C}$

$$(F^V)^L = (F^R)^V = F^{conj}: \mathcal{C}^V \rightarrow \mathcal{D}^V$$

Exer $unit_1 \in \mathcal{C} \otimes \mathcal{C}^V$
 $unit_2 \in \mathcal{D} \otimes \mathcal{D}^V$

Construct $F \otimes F^{conj} (unit_1)$
 \downarrow
 $unit_2$

Exer TFAE

- F is a Verdier quot
- F^{conj} is a Verdier quot

- $F \otimes F^{conj} (unit_1) \xrightarrow{\sim} unit_2$

- $F \circ F^R \xrightarrow{\sim} id$

Prop 1 $j^* \circ \text{Loc}$ is a quotient.

Proof.

$$\begin{array}{ccccc}
 \text{KL}_{\text{Ran}} & \xrightarrow{\text{Loc}} & D(\text{Bun}_h) & \xrightarrow{j^*} & D(U) \\
 \uparrow \text{ind} & & \uparrow \text{ind} & & \uparrow \text{ind} \\
 \text{Rep}(L^t_h)_{\text{Ran}} & \xrightarrow[\text{Loc}]{\text{Qcoh!}} & \text{Qcoh}(\text{Bun}_h) & \xrightarrow{j^*} & \text{Qcoh}(U)
 \end{array}$$

$j^*, \text{Loc}^{\text{Qcoh}}$
preserves cpt's.

$$\begin{array}{ccc}
 & \swarrow \text{r-pull} & \searrow \int_{\text{Ran}} \\
 \text{IBL}^t_h \text{Ran} & U \times \text{Ran} & U
 \end{array}$$

$\Rightarrow j^*, \text{Loc}$ preserves cpt's

$$\text{KL} \simeq \text{KL}^\vee$$

$$D(U) \simeq D(U)^\vee$$

Claim (Exer.) $j^* \circ \text{Loc}$ is self-conjugate via the above self-dualities

$$h\text{-mod } K \xrightarrow[\text{Loc}]{\ell/2} D^{\ell/2}(\tilde{y}/K)$$

$$\text{Loc}^\ell \xleftrightarrow{\text{conj.}} \text{Loc}^\ell$$

Apply paradigm, only need $(j^* \otimes j^*)(\text{Loc} \otimes \text{Loc})(\text{unit}_{\text{KL}}) \xrightarrow{\sim} \Delta_{*, \text{dR}} \omega_{U/U}$

$$\Delta: U \rightarrow U \times U$$

Stronger claim:

Key Cal 1

$$(\text{Loc} \otimes \text{Loc})(\text{unit}_{\text{KL}}) \xrightarrow{\sim} \Delta_{*, \text{dR}} \omega_{\text{Bun}_h}, \quad \Delta: \text{Bun}_h \rightarrow \text{Bun}_h \times \text{Bun}_h.$$

Rmk. ① unit_{KL} is CADO chiral alg. diff. operator

$$\begin{array}{ccc}
 D(L_h) & \xrightarrow{\Gamma_{\text{Ind}^h}} & L_g\text{-mod} \otimes L_g\text{-mod} \\
 \downarrow \text{ind} & & \downarrow \text{ind} \\
 L_h \sim L_h \hookrightarrow L_h & \xrightarrow[\text{(fact. unit)}]{\delta_{L^t_h}} & \text{CADO} \quad (L^t_h \times L^t_h - \text{integrable}) \\
 (L^t_h \times L^t_h\text{-inv}) & & \text{Paralog}
 \end{array}$$

$$② \quad \text{Rep}(L^+G) \longrightarrow \mathcal{Q}\text{coh}(U) \checkmark$$

$$\boxed{\text{Key Cal 2}} \quad \text{Loc}^{\mathcal{Q}\text{coh}} \otimes \text{Loc}^{\mathcal{Q}\text{coh}}(\text{unit}_{\text{Rep}(L^+G)})$$

is

$$\Delta^* \mathcal{O}_{\text{Bun}_G}$$

$\Leftrightarrow j^*, \text{Loc}^{\mathcal{Q}\text{coh}}$ is a quotient

$$\forall j: U \rightarrow \text{Bun}_G$$

$$③ \quad \text{Ex. Key 2} \Rightarrow \text{Key 1.}$$

$$(\text{id} \otimes \text{id})(\text{unit}_{\text{Rep}(L^+G)}) \simeq (\text{id} \otimes \text{obv})(\text{unit}_{K_L})$$

$$\left[\begin{array}{l} F \otimes \text{id}(\text{unit}_1) \\ = \text{id} \otimes F^V(\text{unit}_2) \end{array} \right]$$

$$④ \quad \text{Rep}^V(\check{G})_{\text{Ran}} \xrightarrow{\text{Loc}^{\text{Spec}}} \mathcal{Q}\text{coh}(LS_G^V)$$

$$\boxed{\text{Key Cal 3}} \quad \text{Loc}^{\text{Spec}} \otimes \text{Loc}^{\text{Spec}}(\text{unit}_{\text{Rep}^V(\check{G})}) \longrightarrow \Delta^* \mathcal{O}_{LS_G^V}$$

$\Leftrightarrow \text{Loc}^{\text{Spec}}$ is a quotient

⑤ relative Nish

$$\text{Fun}(\text{Sect}_{\nabla}(X, Z)) = C^{\text{fact}}(X, \text{Fun}(L^+_{\nabla} Z))$$

$$\begin{array}{c} Z \\ \downarrow \\ \text{pt} \end{array}$$

$$Z \rightarrow Y$$

$$(IB_{\check{G}}^V \rightarrow IB_{\check{G}}^V \times IB_{\check{G}}^V)$$

for Key 3

$$w(B_G) \rightarrow w(B_G \times B_G) \text{ for Key 2}$$

w is restr. along $X \rightarrow X_{dR}$

$$⑥ \quad \text{Poinc}! \otimes \text{Poinc}_*(\text{unit}_{\text{unit}}) \xrightarrow{?} \Delta^* w_{B_{\check{G}}}^V \text{ can't prove.}$$

Goal 2

$$\begin{array}{ccc}
 \text{KL}_{\text{Ran}} & \xrightleftharpoons{\text{FLE}} & \text{IndCoh}^*(\mathcal{O}_{\check{X}}^{\text{mf}})_{\text{Ran}} \\
 \text{Loc}^{\text{cusp}} \downarrow \uparrow \text{F}^{\text{cusp}} & & \downarrow \text{Poinc}^{\text{ired}} \uparrow \text{coct}^{\text{ired}} \\
 \text{D}(\text{Bun}_h)_{\text{cusp}} & \xrightleftharpoons[\text{L}^{\text{R}}_{\text{cusp}}]{\text{L}^{\text{cusp}}} & \text{QCoh}(\text{LS}_{\check{X}}^{\text{ired}})
 \end{array}$$

Claim Every cat above is self-dual.

Every functor is self-conjugate.

① $\text{KL} \simeq \text{KL}^\vee$

② $\text{D}_{\text{cusp}} \simeq \text{D}_{\text{cusp}}^\vee$

③ $\text{QCoh}(\text{LS}_{\check{X}}^{\text{ired}}) \simeq \text{QCoh}(\text{LS}_{\check{X}}^{\text{ired}})^\vee$

④* $\text{IndCoh}^*(\mathcal{O}_{\check{X}}^{\text{mf}}) \simeq \text{IndCoh}^*(\mathcal{O}_{\check{X}}^{\text{mf}})^\vee$

Cheat: FLE is an equivalence ① \Rightarrow ④

$$\text{IndCoh}_* \simeq (\text{IndCoh}^!)^\vee$$

$$\text{IndCoh}^!(Y) \rightsquigarrow \text{IndCoh}_*(Y)$$

Need ω^{take} in $\text{IndCoh}_*(Y)$

⑤ FLE \checkmark

⑥ $\text{Loc}^{\text{cusp}} \checkmark$

⑦ $\text{Poinc}^{\text{ired}}$ is self-conjugate


⑧ L_{cusp} is ...

⑦ \Rightarrow ⑧ \checkmark

(Goal 2)

$$\text{L}_{\text{cusp}}^{\omega_j} = \text{L}_{\text{cusp}}$$

$$(\text{L}_{\text{cusp}}^{\text{R}})^\vee \rightsquigarrow \text{L}_{\text{cusp}}^{\text{R}} \simeq \text{L}_{\text{cusp}}^\vee$$



By def'n, $\text{Poinc}^{\text{ined}} = \pi'_* \cdot \boxed{s'^*}$

$(\text{Poinc}^{\text{ined}})^{\text{conj}} = \pi'_! \cdot \boxed{(s')^*}$

Lemma. π' is proper.

$$\begin{aligned}
 A_{h, \text{cusp}} \varphi - &= \mathbb{L}_{\text{cusp}} \cdot \mathbb{L}_{\text{cusp}}^L \\
 &= \mathbb{L}_{\text{cusp}} \cdot \mathbb{L}_{\text{cusp}}^R \quad (\text{as endomorphisms}) \\
 &\quad \text{is (b/c } \Gamma_{\text{cusp}} \text{ is f.f.)}
 \end{aligned}$$

$$\mathbb{L}_{\text{cusp}} \cdot \Gamma_{\text{cusp}}, \text{Loc}^{(\text{cusp})} \cdot \mathbb{L}_{\text{cusp}}^R$$

is

$$\text{Poinc}^{\text{ined}} \cdot \text{FLE}^{-1} \cdot \text{FLE} \cdot \text{coeff}^{\text{ined}}$$

is

$$\text{Poinc}^{\text{ined}} \cdot \text{coeff}^{\text{ined}}$$

Want. $\text{Poinc}^{\text{ined}} \cdot \text{coeff}^{\text{ined}} (\cup \mathcal{L}_{\check{A}}^{\text{ined}}) = A_{h, \text{cusp}}$

$$\pi'_! \cdot s^* \cdot s_* \cdot \pi'^!$$

Prop 1 . $\pi_! \cdot s^* \cdot s_* \cdot \pi^! \Rightarrow \pi_! \cdot \pi^!$

Prop 2

$$\begin{array}{ccc}
 \mathcal{O}_{\tilde{U}, \text{Ran}}^{mb, \text{glob}, \text{inval}} & & \mathcal{O}_{\sigma, \text{Ran}} \\
 \downarrow \subset & & \downarrow \\
 \pi \left(\begin{array}{c} (LS_{\tilde{U}}^{\text{inval}})^{\wedge} \\ \downarrow \omega \\ LS_{\tilde{U}}^{\text{inval}} \end{array} \right) & & (\mathcal{O}_{\sigma, \text{Ran}})_{dR} \\
 & \Rightarrow & \downarrow \\
 & & \sigma
 \end{array}$$

Prop 2 . $\pi_! \pi^! \Rightarrow \omega_! \omega^!$

Prop 2' $\text{End}_{\text{IndCoh}}(\omega_{\mathcal{O}_{\sigma, \text{Ran}}}) = \text{End}_{\text{Dmod}}(\omega_{\mathcal{O}_{\sigma, \text{Ran}}}) = C.(\mathcal{O}_{\sigma, \text{Ran}})$

$$\mathcal{O}_{\tilde{U}}^{\vee} \longrightarrow \text{IB}_{\tilde{U}}^{\vee}$$

$$\mathbb{Z} \longrightarrow X \quad \text{affine D-scheme}$$

$$\begin{array}{ccc}
 \text{Sect}_{\triangleright}(X-\underline{x}, \mathbb{Z})_{\text{Ran}} & & \\
 \downarrow \pi & & \downarrow s \\
 \text{Sect}_{\triangleright}(X-\underline{x}, \mathbb{Z})_{\text{Ran}, dR} & & L_{\triangleright}(\mathbb{Z})_{\text{Ran}} \\
 \nwarrow \pi_{dR} & & \searrow
 \end{array}$$

Prop 1 . $\pi_! \cdot s^* \cdot s_* \cdot \pi^! \Rightarrow \pi_! \pi^!$

Prop 2 $\pi_! \circ \pi^! = \pi_{!, dR} \cdot \pi_{dR}^!$

Lecture 11 (Kevin Lin)

$$D(\text{Bun}_g) \xrightarrow{\mathbb{L}_g} \text{IndCoh}_X(\text{LS}_g^\vee)$$

$\text{Coh}(\text{LS}_g^\vee)$ - linear

$$\begin{array}{ccc} \text{Vect}^{\text{vac}} & & \\ \text{coet}^* \downarrow & & \downarrow r \\ \text{Vect} & = & \text{Vect} \end{array}$$

$$\begin{array}{ccc} \text{Vect} & = & \text{Vect} \\ \text{Poinc}^{\text{vac}} \downarrow & & \downarrow \mathcal{O}_{\text{LS}_g^\vee} \\ D(\text{Bun}_g) & \xleftarrow{\mathbb{L}_g^L} & \text{IndCoh}_X(\text{LS}_g^\vee) \end{array}$$

A_g algebra in $\text{Coh}(\text{LS}_g^\vee)$ producing the monad $\mathbb{L}_g \circ \mathbb{L}_g^L$.

We need to show $\mathcal{O}_{\text{LS}_g^\vee} \rightarrow A_g$ is an isom.

Its restriction along $(\text{ind})^\wedge : (\text{LS}_g^\vee)_{\text{LS}_g^\vee}^\wedge \rightarrow \text{LS}_g^\vee$ is an isom.

It remains to study $\mathcal{O}_{\text{LS}_g^\vee} \rightarrow A_g^{\text{cusp}}$

Facts about A_g^{cusp}

(i) it's a classical vector bundle

(ii) it admits a connection w/ finite monodromy.

Proof of (i) Dennis explained A_g^{cusp} is a self-dual perfect complex.

Lin explained $(\sigma)^* A_g^{\text{cusp}} \simeq C_g(\text{generic oper str. on } \sigma)$

□

Dima showed this space is non empty.

[BKS] proved contractibility for classical groups



\check{G} simply conn'd, $g \geq 2$.

More facts: \check{G} simply conn'd, $g \geq 2$,

I. $LS_{\check{G}}$ is a classical li stack [BD1]

II. $LS_{\check{G}}^{\text{inv}} \subset LS_{\check{G}}$ has codim ≥ 2

III. $LS_{\check{G}}^{\text{inv}}$ is simply conn'd.

IV. $\Gamma(LS_{\check{G}}, A_G) = k$.

Why does this imply $\mathcal{O}_{LS_{\check{G}}^{\text{inv}}} \Rightarrow A_G^{\text{cusp}}$?

III $\Rightarrow A_G^{\text{cusp}} \simeq (\mathcal{O}_{LS_{\check{G}}^{\text{inv}}})^{\oplus 2k}$

$H^0(LS_{\check{G}}^{\text{inv}}, A_G^{\text{cusp}}) \simeq H^0(LS_{\check{G}}^{\text{inv}}, \mathcal{O}_{LS_{\check{G}}^{\text{inv}}})^{\oplus 2k}$

(I+II) \Rightarrow

$H^0(LS_{\check{G}}, A_G) \simeq k$



Proof of II

smooth

$LS_{\check{G}}^{\text{stable}} \subset LS_{\check{G}}^{\text{inv}} \subset LS_{\check{G}}$

affine
fibration

\downarrow

stable

$Bun_{\check{G}} \xrightarrow{\text{codim} \geq 2 \text{ complement}} Bun_{\check{G}}$

if $\check{G} = SL_n$, $(\mathcal{E}, \wedge^n \mathcal{E} \simeq \mathcal{O})$ is stable iff every subbundle $\mathcal{E}' \subset \mathcal{E}$ has $\deg(\mathcal{E}') < 0$.

The obstruction to P_A^\vee admitting a connection is the Atiyah class

$$\alpha_{P_A} \in H^1(X, \Omega^1 \otimes P_A \otimes \check{g}^\vee) \simeq H^0(X, P_A \otimes \check{g}^\vee)$$

Bun_G is simply conn'd

Proof sketch of III.

$$\Gamma(LS_A^\vee, A_A) = \text{Maps}_{\text{IndMod}_G(LS_A^\vee)}(\mathcal{O}_{LS_A^\vee}, \mathbb{L}_A = \mathbb{L}_A^L \otimes \mathcal{O}_{LS_A^\vee})$$

$$\simeq \text{End}_{D(\text{Bun}_G)}(\underbrace{\mathbb{L}_A^L \otimes \mathcal{O}_{LS_A^\vee}}_{\text{Poinc}^\vee!})$$

$$\begin{array}{ccc} \text{Bun}_G^\Omega & \rightarrow & \text{Bun}_G^\Omega / T \xrightarrow{\text{locally closed}} \text{Bun}_G \\ \downarrow \text{continuous fibers} & & \downarrow \\ \mathbb{A}^2 & \xrightarrow{\pi} & \mathbb{A}^2 / T \\ \downarrow \text{sum} & & \\ \mathbb{A}^1 & & \end{array}$$

this End group is $\text{End}(\pi_! \cdot \text{sum}^*(\exp))_{D(\mathbb{A}^2/T)}$

$$\text{for } G = \text{SL}_2, \quad \text{Bun}_G^\Omega = \{\Omega^{1/2} \rightarrow \mathcal{E} \rightarrow \Omega^{-1/2}\} \simeq H^1(X, \Omega^1) \simeq \mathbb{A}^1$$

We are looking at $\pi_! \cdot \exp \simeq j_* k$ [shift]

$$\mathbb{A}^1 / \mathcal{G}_m \xleftarrow{j} \mathcal{G}_m / \mathcal{G}_m = \text{pt}$$