

Enriques surfaces

Cole Franklin

$$S, \quad b_2(S) = 10, \quad \kappa(S) = 0, \quad 2K_S = 0, \quad \chi(\mathcal{O}_S) = 1$$

Def (1) If $h^{0,1} = 0$, we say S is classical

(2) If $h^{0,1} = 1$, & $\mathbb{F}: H^1(S, \mathcal{O}_S) \rightarrow H^1(S, \mathcal{O}_S)$ bijective, S is ordinary

(3) — " — $\mathbb{F} = 0$, we say S is supersingular

Prove ① all Enriques have $h^{0,1} = p_g = 0$ unless $\text{char } k = 2$.

$\text{char } k = 0$,

$$0 = \Delta_S = 2h^{0,1} - \begin{array}{c} 1 \\ 1 \\ 0 \end{array} \Rightarrow h^{0,1} = 0$$

$\text{char } k = p > 0$: assume $h^{0,1} = 1$, take basis $e \in H^1(\mathcal{O}_S)$

$$\mathbb{F}e = \lambda e, \quad \mathbb{F} + \lambda: \mathcal{O}_{S,1} \rightarrow \mathcal{O}_{S,1}, \quad \alpha_\lambda = \ker(\mathbb{F} - \lambda)$$

$$0 \rightarrow \alpha_\lambda \rightarrow \mathcal{O}_S \xrightarrow{\mathbb{F} - \lambda} \mathcal{O}_S \rightarrow 0$$

$$H^1(S, \alpha_\lambda) \rightarrow \ker(\mathbb{F} - \lambda) \rightarrow 0$$

\exists nontrivial α_λ for $X \xrightarrow{\pi} S$

Claim (i) X is Cohenstein & $\omega_X \simeq \mathcal{O}_X$.

(ii) $\pi^* \mathcal{O}_X$ has a compo series & p factors \mathcal{O}_S .

$$\chi(\mathcal{O}_X) \leq h^0(\mathcal{O}_X) + h^2(\mathcal{O}_X) = 2$$

$$\chi(\mathcal{O}_X) = \chi(\pi_* \mathcal{O}_X) = p \chi(\mathcal{O}_S) = p \Rightarrow p \leq 2.$$

@ [Thm 3 in Mumford III] $\exists S \rightarrow \mathbb{P}^1$ w/ $P_n = 1$ fibers

Def A curve $D = \sum n_i E_i$ is of canonical type if $K \cdot E_i = 0 = D \cdot E_i$.

D is indecomposable if $\gcd(n_i) = 1$.

Pr @: ETS $\exists D$ of canonical type.

$$\text{Let } p_0 = \min_{c \text{ inv'd}} p_g(c) \quad \text{s.t. } c^2 > 0$$

$$= \min \left(\frac{1}{2} c^2 + 1 \right)$$

non singular rat'l

Take $D \in |C|$ s.t. D is reducible but not $E + E'$

Step 1: Let E be a comp of D , $D = E + E'$

$$0 < c^2, \quad D \cdot E' = C \cdot E' \geq 0$$

$$D \cdot E = \underbrace{D^2}_{\geq 0} - \underbrace{D \cdot E'}_{\geq 0} \leq c^2$$

But $D \cdot E = E^2 + E \cdot E'$ so either $E \cdot E' > 0$ or $E^2 < 0$ in either case,

$E^2 < c^2$ so $E^2 \leq 0$ so $p_g(E) \in \{0, 1\}$. $\Leftrightarrow E = 0$ or -2 .

Step 2. $D = \sum m_i E_i$ $E_i^2 = 0 \Rightarrow E_i$ is of canonical type

so wlog $E_i^2 = -2$.

- $\sum m_i \geq 3$

- $E_i \cdot E_j \geq 2$ ^{Assume} $(E_i + E_j)^2 = E_i^2 + E_j^2 + 2E_i \cdot E_j \geq 2$

$1 \leq \dim |E_i + E_j|$ By R-R,

$$\chi(E_i + E_j) = \chi(D) + \frac{1}{2} (E_i + E_j)^2 \geq 2$$

But $|E_i + E_j| + |D - E_i - E_j| \leq |C|$

Step 3. If $E_i \cdot E_j = 2$, then $E_i + E_j$ canonical

so wlog $E_i \cdot E_j \in \{0, 1\}$.

Graph: vertices E_i , Edge if $E_i \cdot E_j = 1$

$$\tilde{A}_n, \tilde{D}_n, \tilde{E}_6, \tilde{E}_7, \tilde{E}_8$$

or is Dynkin diagram $D^2 < 0 < C^2 = D^2 \Rightarrow C = \emptyset$.

③ $p = b_2 = 10$. $\forall S$ Enriques.

Pl [Mum I, Thm 4] Thm. $p = b_2$ \forall quasi-elliptic surfaces.

Pl $\exists Y \xrightarrow{\quad} S$ Y smooth ruled surface
 \uparrow
 proper of degree p .

$$\text{Pic}(Y) \otimes \mathbb{Z}_\ell \longrightarrow H^2(Y, \mathbb{Z}_\ell)$$

$$\pi^* \uparrow \downarrow \pi_* \quad \pi^* \uparrow \downarrow \pi_*$$

$$\text{Pic}(S) \otimes \mathbb{Z}_\ell \longrightarrow H^2(S, \mathbb{Z}_\ell)$$

Prop If S Enriques surface is elliptic $S \xrightarrow{\theta} \mathbb{P}^1$, it suffices to show

$$P = b_2 \text{ on } J(S) \xrightarrow{f} \mathbb{P}^1.$$

Fact : $b_i(J(S)) = b_i(S)$ [Prop 4.3.874 in Dalg I]

$$P(S) = P(J(S)).$$

[Thm 2 of Mum I] gives $\omega_{J(S)} = f^*((R^1 f_* \mathcal{O}_S)^{-1} \otimes \omega_{\mathbb{P}^1})$

so $\omega_{J(S)} = f^* \mathcal{O}(-1)$, so all plurigenera of $J(S)$ vanish

by Castelnuovo's rationality crit, $J(S)$ is rat'l

$$\text{so } P(J(S)) = b_2(J(S)). \quad \square$$

$$\textcircled{4} \quad \text{Pic}_{S/k}^\tau = \begin{cases} \mathbb{Z}/2\mathbb{Z} & S \text{ classical} \\ \mu_2 & \text{ordinary} \\ \alpha_2 & \text{supersingular} \end{cases}$$

Pr S classical $H^1(S, \mathcal{O}_S) = 0 \Rightarrow \text{Pic}_S$ smooth

$$\text{Pic}_S^\tau = \text{Tors}(\text{Pic}(S))$$

τ - torsion

$$1 = \chi(F)$$

so $F \simeq F^\vee \otimes \omega_S$ has a section

$$\text{s. } F = \mathcal{O}_S \otimes \omega_S$$

So Pic_S^τ supported at a point if S non-classical.

Thm [1.3.1, Delg I] Any Enriques surface has a degree 2 cover $\pi: X \rightarrow S$ which is a torsor for $(\text{Pic}_S^\tau)^\vee$.

Classical:

$$1 \rightarrow \mu_2 \rightarrow G_m \rightarrow G_m \rightarrow \mathbb{Z}$$

$$H^1(S, \mu_2) \rightarrow H^1(S, G_m)[2] \rightarrow 0$$

non-classical. \square

So we call $X \rightarrow S$ the "K3" cover of S .

Thm. X is Gorenstein $\iff H^1(\mathcal{O}_X) = 0, \omega_X \cong \mathcal{O}_X$.

(1) If $p \neq 2$ or S ordinary, X is smooth

(2) Else X is not smooth.

In case II, $\pi: X \rightarrow S$ is purely inseparable $\implies \pi$ is a homeomorphism $X_{\text{ét}} \rightarrow S_{\text{ét}}$

$$\text{so } c_2(X) = c_2(S) = 12$$

$$\chi(\mathcal{O}_X) = 2\chi(\mathcal{O}_S) = 2$$

But if X smooth, $\underset{\substack{\uparrow \\ 2\chi}}{12} \chi(\mathcal{O}_X) = \chi_X^2 + c_2(X) = 12$

The first part of the paper is devoted to the study of the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the

of the system of equations (1) and (2) and the