IMPACTLENS

Impact Evaluation Report

Flagship clinic booking system

EFFECT DETECTED

EFFECT PER PERIOD

+25.0

95% interval +13.6 to +28.6

RELATIVE LIFT

+11.9%

+6.5% to +13.6%

CUMULATIVE EFFECT

+650

over 26 post-intervention periods

Analysis setup

Method	Synthetic Control
Inference	in-space placebo permutation
p-value	0.1250
Treated unit	flagship_clinic
Intervention date	30 June 2025
Periods (pre / post)	78 / 26
Control units	8
Outcome	weekly appointments

1 caveat(s) were raised during the analysis and are listed in full on the next page. Read them before acting on the number above.

Causal estimates are conditional on their assumptions. This report states them rather than hiding them.

Executive summary

What we found

The intervention increased weekly appointments by about 25.0 per period (+11.9%, interval +6.5% to +13.6%). The 95% placebo interval runs from 13.6 to 28.6 and excludes zero.

Over the 26 periods since the intervention, the cumulative difference is approximately 649.6. That total inherits all the uncertainty of the per-period estimate, so it should be read as an order of magnitude rather than an exact figure.

Method. The counterfactual is a weighted blend of control units chosen to track the treated unit before the intervention. The reported interval comes from re-running the analysis on each control unit as a placebo, so it answers 'how unusual is this gap among units that received nothing?'

Data. 104 periods of W-MON data from 01 Jan 2024 to 22 Dec 2025: 78 before the intervention and 26 after, with 8 control unit(s) in the donor pool.

Caveats. Placebo interval built from only 7 donor placebos. The smallest p-value attainable with this donor pool is 0.125; finer significance claims are not supported by the data.


Do the methods agree?


All 4 control-based methods agree in direction, and the spread between their point estimates (17.1) is small relative to their own uncertainty. The conclusion is not an artefact of one modelling choice. The naive before/after comparison reports 42.3 per period -- 1.7x the control-based estimate of 24.9. The extra 17.4 is movement the control units experienced too, so it was never attributable to the intervention.

Caveats raised by the analysis


Placebo interval built from only 7 donor placebos. The smallest p-value attainable with this donor pool is 0.125; finer significance claims are not supported by the data.


The effect over time







Accumulation and panel context







Diagnostics: can this result be trusted?







Method comparison





Every method side by side

Where the methods agree, the finding is robust to the modelling choice. Where they disagree, the headline number is an artefact of assumptions and should not be quoted without explanation. The naive row is a benchmark showing what a before/after dashboard would have reported.

	effect	ci_lower	ci_upper	relative_effect	p_value	significant	n_warnings
	24.91575776335414	14.971851178695049	34.32240099669261	0.11871707928533078	0.0140140140140140	True	1
	24.984680345420313	13.598812232962428	38.60929208004161	0.1190845845425300	0.125	True	1
style)	25.658099400094727	21.359133608517929	29.930396873662616	0.1226904033044470	0.0	True	0
	42.04065816033283	34.362014305542949	49.719302015122	0.2218109554806177	0.0	True	2
	42.27828548121631	35.522806162896495	49.033764799535404	0.2196131247852560	0.0	True	1

Validation: does the tool recover a known effect?





Recovery study results

Each estimator was run on simulated data where the true effect was inserted by us, so the correct answer is known. 'coverage' is the share of runs whose interval contained the truth and should be close to the nominal level; materially lower means the intervals are too narrow. 'power' is the share of runs that detected the effect.

	label	n_runs	true_effect	mean_estimate	bias	rmse	coverage	power
	Difference-in-Differe	250	23.97	24.14	0.1685	2.354	1	0.98
_control	Synthetic Contro	250	23.97	24.15	0.1852	2.501	0.936	0.868
	Bayesian Counterfac	250	23.97	24.97	0.9964	2.828	0.9	1
	Interrupted Time Se	250	23.97	44.85	20.88	21.19	0	1
	Naive pre/post (bench	250	23.97	41	17.03	17.17	0	1

Assumptions and limits

What must be true for this number to be valid

Good pre-period fit: the weighted donors must track the treated unit closely before the intervention. A poor pre-fit makes the counterfactual worthless.

No interference: donor units were not themselves affected by the intervention.

Convex hull: the treated unit's behaviour must be reachable by a weighted average of the donors. An outlier treated unit cannot be matched.

Stable relationship: the pre-period relationship between treated unit and donors continues to hold through the post period.

When this tool gives the wrong answer

The control units were affected by the intervention too. Spillover shrinks the measured gap and the estimate understates the effect.

Something else changed at the same date and affected only the treated unit. That change is indistinguishable from the intervention and will be credited to it.

The pre-period is short. The baseline is then poorly characterised and the counterfactual is close to a guess.

The treated unit was selected because it was already improving. Selection on the outcome biases every method here upward.

Dataset facts

N units: 9

N controls: 8

N periods: 104

N pre: 78

N post: 26

Frequency: W-MON

First date: 2024-01-01 00:00:00

Last date: 2025-12-22 00:00:00

Missing cells: 0

Treated pre mean: 192.51256281953496

Treated pre std: 17.85546654682822

Dropped rows: 0