

Lecture 10: Distributions

Ethan Ebbighausen

July 22nd, 2026

1 Introduction

How far can we push weak solutions? We have defined a weak derivative, which may go well beyond just Sobolev spaces, and we have also observed that there are some interesting objects that behave like measures. Our case-in-point is the heat kernel $H_t(x)$, which converged pointwise as $t \rightarrow 0$ to

$$\begin{cases} \infty & x = 0 \\ 0 & \text{else} \end{cases}$$

but when we measured it via integration, $\lim_{t \rightarrow 0} \int H_t(y) f(x - y) dy = f(x)$, so this limit acts like the probability distribution assigning probability 1 to 0 and probability 0 to everything else. We noticed before that H_t looked like a derivative of a smoothed-out step function, so we are saying that this “point density” looks like the derivative of the step function (which is not differentiable, so we may try using weak definitions).

Surely, the weird pointwise behavior isn’t enough to describe this, because we can generate other functions with the same pointwise behavior where this fails. We can see this behavior arise in another way physically, in electrostatics, through Coulomb’s law.

1.1 Model: Coulomb’s Law

Charles-Augustin de Coulomb published an empirical observation of electristatics in the 18th century, describing that a particle with electric charges q_0 located at the origin generates an electric field given by

$$E(x) = \frac{kq_0 x}{|x|^3}$$

for a constant k . Recall that Gauss’ law also told us $\nabla \cdot E = 4\pi k\rho$. The weird part about this condition is that, except at $x = 0$,

$$\nabla \cdot \frac{x}{|x|^3} = \frac{\nabla \cdot x}{|x|^3} - \frac{3x}{|x|^4} \nabla |x| = 0$$

so the behavior on a null set must somehow cause Gauss’ law. We rectify this by using the weak form of Gauss’ law,

$$\int_{\mathbb{R}^3} E \cdot \nabla \psi dx = -4\pi k \int_{\mathbb{R}^3} \rho \psi dx$$

for all test functions ψ . As long as we exclude the origin, we can integrate by parts, so we write

$$\begin{aligned} \int_{\mathbb{R}^3} E \cdot \nabla \psi dx &= \lim_{\epsilon \rightarrow 0} \int_{|x| \geq \epsilon} E \cdot \nabla \psi dx \\ &= \lim_{\epsilon \rightarrow 0} \int_{|x|=\epsilon} \nu \cdot E \psi dS = - \int_{|x|=\epsilon} \frac{kq_0}{\epsilon^2} \psi dS \\ &= -4\pi kq_0 \psi(0) \end{aligned}$$

so this requires

$$\int_{\mathbb{R}^3} \rho \psi dx = q_0 \psi(0)$$

another instance of a point density.

Learning Objectives:

- Define a distribution and compute distributional derivatives and limits.
- Define fundamental solutions, use them to solve PDEs.
- Compute Green's functions on some domains.

2 Definitions, Convergence, Derivatives

The theory of distributions was developed independently in the mid-20th century by Sergei Sobolev and Laurent Schwartz, as the concept of the dual to the test functions.

A *distribution* on a domain $\Omega \subset \mathbb{R}^n$ is a linear function $C_c^\infty(\Omega) \rightarrow \mathbb{C}$ that is continuous. We write the function as u , and write evaluations as pairings (u, ψ) .

By continuity here, we mean that if $\{\psi_n\} \rightarrow \psi$ in C_c^∞ , then $(u, \psi_n) \rightarrow (u, \psi)$, where converging in $C_c^\infty(\Omega)$ means that there exists some compact K so all the ψ_n are supported in K and all of their partial derivatives converge uniformly to those of ψ on K .

We denote the space of distributions by $D'(\Omega)$ (the prime signifying dual). Notice that $L_{loc}^1(\Omega) \subset D'(\Omega)$ via

$$(f, \psi) = \int_{\Omega} f \psi dx$$

and we have also seen the Dirac delta $(\delta_x, \psi) = \psi(x)$.

We may place the *weak topology* on $D'(\Omega)$, by stating $u_k \rightarrow u$ in $D'(\Omega)$ if

$$\lim_{k \rightarrow \infty} (u_k, \psi) = (u, \psi)$$

for all $\psi \in C_c^\infty$. We may finally formalize the idea behind the heat kernel:

[2.1] Given $f \in L^1(\mathbb{R}^n)$ satisfying $\int_{\mathbb{R}^n} f dx = 1$, define $f_a(x) = a^n f(ax)$. Then,

$$\lim_{a \rightarrow \infty} f_a = \delta_0$$

as a distributional limit.

Proof. For $\psi \in C_c^\infty(\mathbb{R}^n)$,

$$(f_a, \psi) = \int_{\mathbb{R}^n} a^n f(ax) \psi dx = \int_{\mathbb{R}^n} f(x) \psi(x/a) dx$$

so

$$|(f_a, \psi) - \psi(0)| \leq \int_{\mathbb{R}^n} |f(x)(\psi(x/a) - \psi(0))| dx$$

We now use the decay of f and the continuity of ψ . Pick $\epsilon > 0$. Assume $|\psi| \leq M$. Then, we may find R so

$$\int_{|x| > R} |f| dx \leq \frac{\epsilon}{2M}$$

and $\delta > 0$ so $|\psi(y) - \psi(0)| < \epsilon/2$ for $|y| < \delta$. Setting $a \geq R/\delta$,

$$\int_{\mathbb{R}^n} |f(x)(\psi(x/a) - \psi(0))| dx \leq \int_{|x| > R} |f|(2M) dx + \int_{|x| \leq R} |f| \epsilon/2 dx < \epsilon$$

so $(f_a, \psi) \rightarrow \psi(0)$ as desired. $\square$

Remark: This is a version of mollification, a process introduced by Kurt Otto Friedrichs in 1944.

We may also extend many operations to D' . Because the idea of the integral pairing is so central to the construction of the distributions, we define many operations by mimicking that case. For example, when $f \in C^\infty(\Omega)$, we define uf to be the distribution so

$$(uf, \psi) = (u, f\psi)$$

for all ψ . Similarly, we define the *distributional derivative* $\partial^\alpha u$ to be the distribution

$$(\partial^\alpha u, \psi) = (u, (-1)^{|\alpha|} \partial^\alpha \psi)$$

Example: Notice that

$$\int_{\mathbb{R}} -1_{[0, \infty)} \psi' dx = \int_0^\infty -\psi'(x) dx = \psi(0)$$

so that

$$\frac{d}{dx}(-1_{[0, \infty)}) = \delta_0$$

as distributions. We may compute further derivatives by noticing

$$(\partial^\alpha \delta_0, \psi) = (-1)^{|\alpha|} \partial^\alpha \psi(0)$$

Example:

The function $\ln|x|$ is locally integrable and so it defines a distribution on $\mathbb{R}$. We may expect the distributional derivative to be $\frac{1}{x}$ for $x \neq 0$, but this is not locally integrable. To understand this, we go back to the integral pairing:

$$\begin{aligned} \int_{\mathbb{R}} -\psi' \ln|x| dx &:= \lim_{\epsilon \rightarrow 0} - \int_{|x| \geq \epsilon} \psi' \ln|x| dx \\ &= \lim_{\epsilon \rightarrow 0} \int_{|x| \geq \epsilon} \frac{\psi(x)}{x} dx + [\psi(\epsilon) - \psi(-\epsilon)] \ln(\epsilon) \\ &= \lim_{\epsilon \rightarrow 0} \int_{|x| \geq \epsilon} \frac{\psi(x)}{x} dx \end{aligned}$$

since

$$\lim_{\epsilon \rightarrow 0} [\psi(\epsilon) - \psi(-\epsilon)] \ln(\epsilon) = \psi'(0) \lim_{\epsilon \rightarrow 0} 2\epsilon \ln|\epsilon| = 0$$

The limit definition of the integral is given a special name, the principal value distribution $PV[x^{-1}]$, and only converges because it is taken symmetrically.

Example:

Gauss' law from the introduction, combined with Coulomb's law, may be rewritten as

$$\nabla \cdot \frac{x}{|x|^3} = 4\pi\delta$$

3 Fundamental Solutions

Imagine we have a matrix L , and vectors f_i so $Lf_i = e_i$. How do we solve $Lf = v$? We superimpose solutions $f = \sum v_i f_i$ so $Lf = \sum v_i e_i = v$.

Now, if instead we have a differential operator L and solutions Φ_x so $L\Phi = \delta_0$, we can solve $Lu = f$ by superimposing solutions

$$u(x) = \int \Phi_x(y) f(y) dy$$

Then, as long as we are justified in swapping the integral and derivatives,

$$Lu = \int L\Phi_x(y) f(y) dy = f(x)$$

as desired. We therefore call the solution Φ_x the *fundamental solution* at x . With linear operator L , we only need to compute Φ_0 and notice $\Phi_x(y) = \Phi_0(x - y)$ (so we have a convolution above).

We're treating the fundamental solution like a function here, but that may not always be so. To justify these actions, we need to consider convolutions with distributions and how distributional derivatives work with convolutions.

In the first case, when $f, g \in L^1_{loc}(\mathbb{R}^n)$ and $\psi \in C_c^\infty(\Omega)$,

$$(f * g, \psi) = \int \int f(y) g(x - y) \psi(x) dy dx = \int f(y) \int g(y - x) \psi(x) dx dy$$

so that we may move the convolution off f and onto ψ by taking $g^-(x) = g(-x)$. Then,

$$(f * g, \psi) = (f, g^- * \psi)$$

Using the right hand side, we may define the convolution of $u \in D'(\mathbb{R}^n)$ and a test function $\phi \in C_c^\infty(\mathbb{R}^n)$ by

$$(u * \phi, \psi) = (f, \phi^- * \psi)$$

for all $\psi \in C_c^\infty(\mathbb{R}^n)$. In particular, the δ_0 distribution gives

$$(\delta_0 * \phi, \psi) = \phi^- * \psi(0) = (\phi, \psi)$$

so $\delta_0 * \phi = \phi$.

For the second complication, we have the following

[3.1] For $w \in D'(\mathbb{R}^n)$ and $\psi \in C_c^\infty(\mathbb{R}^n)$,

$$D^\alpha(w * \psi) = (D^\alpha w) * \psi = w * (D^\alpha \psi)$$

Proof:

For ϕ, ψ both test functions,

$$\begin{aligned} D^\alpha(\phi * \psi) &= D^\alpha \int_{\mathbb{R}^n} \phi(x - y) \psi(y) dy \\ &= \int_{\mathbb{R}^n} D^\alpha \phi(x - y) \psi(y) dy = (D^\alpha \phi) * \psi \end{aligned}$$

as desired. Next, for $w \in D'(\mathbb{R}^n)$ and $\phi, \psi \in C_c^\infty(\mathbb{R}^n)$,

$$\begin{aligned} (D^\alpha(w * \phi), \psi) &= (-1)^{|\alpha|} (w * \phi, D^\alpha \psi) = (-1)^\alpha (u, \phi^- * D^\alpha \psi) = (-1)^\alpha (u, D^\alpha(\phi^- * \psi)) \\ &= ((D^\alpha u), \phi^- * \psi) = ((D^\alpha u) * \phi, \psi) \end{aligned}$$

or

$$(-1)^\alpha(u, D^\alpha(\phi^- * \psi)) = (u, D^\alpha\phi^- * \psi) = (u * D^\alpha\phi, \psi)$$

as desired.

Q.E.D.

[3.2] For $\phi \in C_c^\infty(\mathbb{R}^n)$, $u \in \mathcal{D}'(\mathbb{R}^n)$, $u * \phi \in C^\infty(\mathbb{R}^n)$.

Proof:

We claim that

$$u * \phi = \langle u, \phi(x - \cdot) \rangle$$

so it is a function. Then, the above result shows it is smooth. To prove this is a function, notice that for $\psi \in C_c^\infty$, we have

$$\lim_{m \rightarrow \infty} \sum_{x \in \frac{1}{m}\mathbb{Z}^n} \frac{1}{m} \psi(x) \phi(x - \cdot) = \int_{\mathbb{R}^n} \psi(x) \phi(x - \cdot) dx$$

where the limit converges in C_c^∞ . Then,

$$\begin{aligned} \langle \langle u, \phi(x - \cdot) \rangle, \psi \rangle &= \int \langle u, \phi(x - \cdot) \rangle \psi(x) dx = \langle u, \int \phi(x - \cdot) \psi(x) dx \rangle \\ &= \langle u, \phi^- * \psi \rangle = \langle u * \phi, \psi \rangle \end{aligned}$$

as desired.

Q.E.D.

Example: A distribution $u \in \mathcal{D}'(\mathbb{R}^n)$ with $\partial_j u = 0$ for all j is a constant function.

To see this, let $\phi \in C_c^\infty(\mathbb{R}^n)$ so $u * \phi_a$ is smooth. Since $\partial_j(u * \phi_a) = 0$ for all j , this is actually a constant C_a . Further, since $\lim_{a \rightarrow \infty} u * \phi_a = u$ as a distribution, and $u * \phi_a = C_a$ as a distribution, we have

$$\langle u, \psi \rangle = \lim_{a \rightarrow \infty} \langle C_a, \psi \rangle = (\lim_{a \rightarrow \infty} C_a) \int \psi dx$$

so that u is the constant $C = \lim_{a \rightarrow \infty} C_a$.

We now formally state our heuristic

[3.3] If L is a constant coefficient operator on $\mathbb{R}^n$ with fundamental solution $\Phi \in D'(\mathbb{R}^n)$ satisfying $L\Phi = \delta_0$, then for $f \in C_c^\infty(\mathbb{R}^n)$,

$$Lu = f$$

is solved by

$$u = \Phi * f$$

Proof:

Let

$$Lf = \sum_{|\alpha| \leq m} a_\alpha D^\alpha f$$

Then,

$$L(\Phi * f) = \sum_{|\alpha| \leq m} a_\alpha D^\alpha(\Phi * f) = (L\Phi) * f = \delta_0 * f = f$$

Q.E.D.

3.1 Laplace Solution

To motivate our fundamental solution, we recall that the Laplacian is rotationally invariant. It may then make sense to try to reduce to an ODE in r , in which case we have

$$r^{1-n} \partial_r (r^{n-1} \partial_r \rho) = \partial_r^2 \rho + \frac{n-1}{r} \partial_r \rho = \delta_0$$

From Coulomb's law in the introduction, we saw that $\frac{1}{Cr^2}$ gives a solution in 3D for the right constant. We then may try a general monomial r^{-l} , in which case

$$(-l)(-l-1)r^{-l-2} + (n-1)(-l)r^{-l-2}$$

is only zero for $r \neq 0$ if $l = n-1$. We then must check at $r = 0$:

[3.4] On $\mathbb{R}^n$, the operator $-\Delta$ has the fundamental solution

$$\Phi(x) = \begin{cases} -\frac{1}{2\pi} \ln(r) & n = 2 \\ \frac{1}{(n-2)A_n r^{n-2}} & n \geq 3 \end{cases}$$

for $A_n = \text{vol}(B(0,1))$ in $\mathbb{R}^n$.

Proof:

We wish to show

$$(-\Delta \Phi, \psi) = \psi(0)$$

First, we notice that

$$\nabla \Phi(x) = -\frac{x}{A_n r^n}$$

so

$$\begin{aligned} \int_{\mathbb{R}^n} \Delta \Phi \psi dx &= - \int_{\mathbb{R}^n} \nabla \Phi \cdot \nabla \psi dx \\ &= \frac{1}{A_n} \int_{\mathbb{R}^n} \frac{x}{r^n} \cdot \nabla \psi dx \\ &= \frac{1}{A_n} \int_{\mathbb{R}^n} \frac{1}{r^{n-1}} \partial_r \psi dx \\ &= \frac{1}{A_n} \int_{S^{n-1}} \int_0^\infty \partial_r \psi dr dS = -\psi(0) \end{aligned}$$

as desired.

Q.E.D.

When the domain is not $\mathbb{R}^n$, it becomes a bit harder to use this symmetry to our advantage. George Green developed a method for doing so in 1828, by connecting fundamental solutions to boundary value problems through an integral formula.

Let Φ be the fundamental solution of the Laplacian on $\mathbb{R}^n$ given above. Let $\Phi_y(x) = \Phi(x-y)$.

[3.5] Green's Representation Formula Suppose that $\Omega \subset \mathbb{R}^n$ is a bounded domain with piecewise C^1 boundary. For $u \in C^2(\overline{\Omega})$,

$$u(y) = - \int_{\Omega} \Phi_y \Delta u dx + \int_{\partial\Omega} [\Phi_y \partial_\eta u - u \partial_\eta \Phi_y] dS$$

for $y \in \Omega$.

Proof:

Because the point y is fixed, we assume $y = 0$. Pick some $\epsilon > 0$ small enough so $\overline{B(0, \epsilon)} \subset \Omega$. Then, on $\overline{\Omega} - B(0, \epsilon)$, Φ is smooth and has $\Delta\Phi = 0$. We then have

$$\begin{aligned} \int_{\Omega - \overline{B(0, \epsilon)}} \Phi \Delta u dx &= \int_{\partial\Omega} (\Phi \partial_\eta u - u \partial_\eta \Phi) dS + \\ &\quad \int_{\partial B(0, \epsilon)} (\Phi \partial_\eta u - u \partial_\eta \Phi) dS \end{aligned}$$

Because Δu is continuous and Φ is locally integrable,

$$\lim_{\epsilon \rightarrow 0} \int_{\Omega - \overline{B(0, \epsilon)}} \Phi \Delta u dx = \int_{\Omega} \Phi \Delta u dx$$

so to prove the representation formula, we need to show

$$\lim_{\epsilon \rightarrow 0} \int_{\partial B(0, \epsilon)} (\Phi \partial_\eta u - u \partial_\eta \Phi) dS = u(0)$$

We treat the two terms separately. First, ∇u is bounded on $\overline{\Omega}$, so we take

$$|\partial_\eta u| \leq |\nabla u| \leq M$$

so

$$\left| \int_{\partial B(0, \epsilon)} \Phi \partial_\eta u dS \right| \leq \frac{M}{n-1} \epsilon$$

(or $M\epsilon \ln(\epsilon)$ if $n = 1$), which disappears as $\epsilon \rightarrow 0$.

However, $\partial_\eta \Phi = \partial_r \Phi = -\frac{1}{A_n r^{n-1}}$ implies

$$-\int_{\partial B(0, \epsilon)} u \partial_r \Phi dS = \frac{1}{|\partial B(0, \epsilon)|} \int_{\partial B(0, \epsilon)} u dS$$

which converges to $u(0)$ as $\epsilon \rightarrow 0$ by continuity.

Q.E.D.

The original goal of this representation formula was to solve the Poisson problem with inhomogeneous Dirichlet boundary condition. Suppose there exists a family of functions $H_y \in C^2(\overline{\Omega})$ for $y \in \Omega$ such that

$$\begin{aligned} \Delta H_y &= 0 \\ H_y|_{\partial\Omega} &= \Phi_y|_{\partial\Omega} \end{aligned}$$

Then, $G_y = \Phi_y - H_y$ has homogeneous boundary conditions and has $\Delta G_y = \delta_y$ in Ω . This is called the Green's function for Ω .

[3.6] Suppose $\Omega \subset \mathbb{R}^n$ is a bounded domain with piecewise C^1 boundary condition that admits Green's function G_y . Then, for $f \in C^0(\Omega)$ and $g \in C^0(\partial\Omega)$,

$$\begin{aligned} -\Delta u &= f \\ u|_{\partial\Omega} &= g \end{aligned}$$

is solved by

$$u(y) = - \int_{\Omega} f G_y dx - \int_{\partial\Omega} g \partial_\eta G_y dS$$

Proof:

By Green's second identity,

$$\int_{\Omega} H_y \Delta u - u \Delta H_y dy = \int_{\partial\Omega} H_y \partial_{\eta} dS$$

Using the representation formula,

$$\begin{aligned} u(y) - \int_{\Omega} \Phi_y \Delta u dx + \int_{\partial\Omega} [\Phi_y \partial_{\eta} u - u \partial_{\eta} \Phi_y] dS - 0 \\ = - \int_{\Omega} \Phi_y \Delta u dx + \int_{\partial\Omega} [\Phi_y \partial_{\eta} u - u \partial_{\eta} \Phi_y] dS - \int_{\Omega} H_y \Delta u dy + \int_{\partial\Omega} H_y \partial_{\eta} dS \end{aligned}$$

which, combining terms, is the formula above.

Q.E.D.

The *method of images* is a trick to help compute the Green's function. It involves trying to place charges outside of the working domain to cancel the boundary terms of the normal fundamental solution. In general, we try to construct $G_y(x)$ by starting with $\Phi(y-x)$, then finding locations $\{\bar{x}_j\}$ and magnitudes (charges) q_j so

$$\sum_j q_j \Phi(y - \bar{x}_j)$$

exactly cancels $\Phi(y-x)$ for $y \in \partial\Omega$

Example: The Half-Plane

If we consider $\Omega = \mathbb{R}_+^n = \{x \in \mathbb{R}^n \mid x^n > 0\}$, then the boundary has $x^n = 0$. Hence,

$$\Phi(y-x) = \Phi(y-\bar{x})$$

for $y \in \partial\mathbb{R}_+^n$ and $\bar{x} = (x^1, \dots, x^{n-1}, -x^n)$ the reflection across the boundary. Therefore, we set

$$G_y(x) = \Phi(y-x) - \Phi(y-\bar{x})$$

to remove boundary conditions. Notice that since $\bar{x}$ is outside of the domain, $\Phi(y-\bar{x})$ is harmonic and we satisfy the differential condition as well.

To use this definition in the representation formula, we only need to compute

$$\partial_{y^d} G_y(x) = -\frac{1}{nA_n} \frac{y^n - x^n}{|y-x|^n} + \frac{1}{nA_n} \frac{x^n + y^n}{|y-\bar{x}|^n}$$

so that for $y = (y', 0)$ and $x = (x', x^n)$,

$$-\eta(y) \cdot D_y G_y(x) = \frac{2x^n}{nA_n} \frac{1}{(|y' - x'|^2 + (x^n)^2)^{n/2}}$$

For example, a harmonic function u with boundary condition g on the upper half plane has, for continuous bounded g ,

$$u(x) = \frac{2x^n}{nA_n} \int_{\mathbb{R}^{n-1}} \frac{g(y')}{(|y' - x'|^2 + (x^n)^2)^{n/2}} dy'$$

Example: The Unit Ball

A similar reflection argument gives the Green's function on $B(0, 1)$, where the reflection instead must be taken across the circle. For $x \in B(0, 1)$, this reflection is

$$\tilde{x} = \frac{x}{|x|^2}$$

Then, for $y \in \partial B(0, 1)$,

$$|\tilde{x} - y|^2 = \left| \frac{x}{|x|^2} - y \right|^2 = \left| \frac{x - |x|^2 y}{|x|^2} \right|^2 = \frac{1}{|x|^2} (|x|^2 - 2x \cdot y + 1)$$

but

$$|x - y|^2 = |x|^2 - 2x \cdot y + 1$$

then implies

$$|y - x| = |x||y - \tilde{x}|$$

Since $\Phi(x) = -\frac{1}{2\pi} \ln |x|$, taking $\Phi(y - x) - \Phi(y - \tilde{x})$ gives

$$G_y(x) = \begin{cases} -\frac{1}{2\pi} \ln \left(\frac{|y-x|}{|x||y-\tilde{x}|} \right) & x \neq 0 \\ = \frac{1}{2\pi} \ln |y| & x = 0 \end{cases}$$

In this version, the outward normal is simply y , so we compute (for $r = |y|$)

$$\partial_r \ln(|y - x|) = x \cdot \nabla \ln |y - x| = y \cdot \frac{y - x}{|y - x|^2}$$

$$\partial_r G_y(x) = -\frac{1 - |x|^2}{2\pi |y - x|}$$

The formula for a harmonic u with boundary value g is then

$$u(x) = \frac{1}{2\pi} \int_{\partial B(0,1)} \frac{1 - |x|^2}{|x - y|^2} g(y) dS(y)$$

(In polar coordinates, this is the Poisson formula we derived in lecture 8).

4 Time-Dependent Fundamental Solutions

We adapt to the time dependence by considering families of distributions, indexed by time. In other words, we have a map $\mathbb{R} \rightarrow D'(\mathbb{R}^n)$ by $t \mapsto w_t$. This map is differentiable with respect to time if there is a family $\partial_t w_t \in D'(\mathbb{R}^n)$ so

$$\frac{d}{dt}(w_t, \psi) = (\partial_t w_t, \psi)$$

for all $\psi \in C_c^\infty(\mathbb{R}^n)$.

We now call W_t a fundamental solution of $\partial_t + L$ if

$$\begin{aligned} \partial_t W_t + L W_t &= 0 \\ W_t|_{t=0} &= \delta \end{aligned}$$

as before, or a solution to

$$\partial_t^2 + L$$

if

$$\begin{aligned}\partial_t^2 W_t + L W_t &= 0 \\ W_t|_{t=0} &= 0 \\ \partial_t W_t|_{t=0} &= \delta\end{aligned}$$

This may seem awkward at first, but this change allows us to take initial conditions as opposed to boundary conditions.

4.1 Wave Equation (1D)

Recall D'Alembert's formula

$$\begin{aligned}u(t, x) &= \frac{1}{2} [g(x+t) + g(x-t)] + \frac{1}{2} \int_{x-t}^{x+t} h(\tau) d\tau \\ &= \frac{1}{2} [g(x+t) + g(x-t)] + \frac{1}{2} 1_{[-t, t]} * h(x)\end{aligned}$$

where we have converted the integral to a convolution. Further, notice that

$$\partial_t 1_{[-t, t]}(x) = \delta_t + \delta_{-t}$$

and

$$u(t, x) = \frac{1}{2} g * \partial_t 1_{[-t, t]} + \frac{1}{2} 1_{[-t, t]} * h$$

this motivates us to guess that $W_t = \frac{1}{2} 1_{[-t, t]}$. Indeed, for a test function ψ ,

$$(\partial_t^2 W_t, \psi) = \frac{1}{2} (-\delta'_{-t} + \delta'_t, \psi) = \frac{1}{2} [\psi'(t) - \psi'(-t)]$$

while

$$(\partial_x^2 W_t, \psi) = (W_t, \partial_x^2 \psi) = \frac{1}{2} [\psi'(t) - \psi'(-t)]$$

by the Fundamental Theorem of Calculus. Therefore,

$$\begin{aligned}\partial_t^2 W_t - \partial_x^2 W_t &= 0 \\ W_t|_{t=0} &= 0 \\ \partial_t W_t|_{t=0} &= \delta_0\end{aligned}$$

as desired.

Fundamental solutions to higher-dimensional wave equations are known, but take an awkward form that arises from studying homogeneous distributions, so we will not display the general case here. For example, in 3D the solution is

$$E_t(x) = -\frac{1}{2\pi} 1_{(0, \infty)}(t) \delta_0(t^2 - |x|^2)$$

4.2 Heat Equation

The heat equation fundamental solution is simple for us because we already found a convolution solution formula

$$u(t, x) = \int_{\mathbb{R}^n} H_t(x-y) h(y) dy$$

Therefore, we would guess that $H_t(x)$ is the fundamental solution. Well, we have proven with some effort that

$$\lim_{t \rightarrow 0} H_t(x) = \delta_0$$

as distributions in the past, and $H_t(x)$ was defined to be a solution to the heat equation. A more historically accurate presentation of the material would have waited to define $H_t(x)$ until this point, and only solved the heat equation with separation of variables until now.