Analyses performed for DAOP well-defined residues.

Procheck analysis, RMSD calculation and structure superimposition are based on DAOP with S(phi)+S(psi)>=1.8 : 2A-60A, 68A-70A, 77A-107A

NESG ID: NAME

PDB ID:

Deposition date:

Common Name:

Class:

Length (a.a.): 108

Organism:
SwissProt /
TrEMBL ID:
models: 5

Oligomerization: monomer

Molecular

weight:

Secondary Structure Elements:

alpha helices: 9A-22A, 29A-40A, 44A-58A, 67A-72A, 79A-104A

beta strands:

FIDs deposited in the BMRB? no

Comparison of core atoms:

DAOP > 1.8 Å: A:2..A:60, A:68..A:70, A:77..A:107

FindCore2: A:4..A:61 **CYRANGE**: 7..54

RMSD All residues Ordered residues² Selected residues³

 All backbone atoms
 16.6 Å
 14.9 Å
 14.9 Å

 All heavy atoms
 15.6 Å
 16.1 Å
 16.1 Å

Ramachandran Plot Summary for selected residues³ from Procheck

Most favoured regions Additionally allowed regions Generously allowed regions Disallowed regions

93.0% 7.0% 0.0%

Ramachandran Plot Summary for selected residues³ from Richardson Lab's Molprobity

Most favoured regions Allowed regions Disallowed regions View plot View model summary

95.3% 4.5% 0.2%

08/03/22



| Program | Verify3D | ProsaII (-ve) | Procheck (phi-psi) ³ | Procheck (all) ³ | MolProbity Clashscore |
|----------------------|----------|---------------|---------------------------------|-----------------------------|-----------------------|
| Raw score | 0.05 | N/A | 0.26 | 0.20 | 0.00 |
| Z-score ¹ | -6.58 | N/A | 1.34 | 1.18 | 1.53 |

Close Contacts and Deviations from Ideal Geometry (from PDB validation software)


Number of close contacts (within 1.6 & Aring for H atoms, 2.2 & Aring for heavy atoms):

RMS deviation for bond angles: 2.0°

RMS deviation for bond lengths:

 $0.013~\mathrm{\AA}$


³ Selected residues DAOP with S(phi)+S(psi)>=1.8: 2A-60A, 68A-70A, 77A-107A




¹ With respect to mean and standard deviation for a set of 252 X-ray structures < 500 residues, of resolution <= 1.80 Å, R-factor <= 0.25 and R-free <= 0.28; a positive value indicates a 'better' score

² Ordered residues (DAOP): 2A-60A, 68A-70A, 77A-107A






Procheck G-factor for phi-psi










Procheck G-factor for all dihedral angles











Residue Plot of Ramachandran anlysis(based on data from Richardson Lab's Molprobity)

References:

- 1. Bhattacharya A, Tejero R and Montelione GT, "Evaluating protein structures determined by structural genomics consortia". Proteins 2007, 66:778-795
- 2. Tejero R, Snyder D, Mao B, Aramini JM and Montelione GT, "PDBStat: a universal restraint converter and restraint analysis software package for protein NMR". J Biomol NMR 2013, 56:337-351
- 3. Luthy R, Bowie JU and Eisenberg D, "Assessment of protein models with three-dimensional profiles". Nature 1992, 356:83-85
- 4. Bowie JU, Luthy R and Eisenberg D, "A Method to identify protein sequences that fold into a known three-dimensional structure". Science 1991, 253:164-169
- 5. Sippl MJ, "Recognition of errors in three-dimensional structures of proteins". Proteins 1993, 17:355-362
- 6. Sippl MJ, "Calculation of conformation ensembles from potentials of mean force". J Mol Biol 1990, 213:859-883
- 7. Laskowski RA et al, "AQUA and PROCHECK_NMR: programs for checking the quality of proteins structures solved by NMR". J Biomolec NMR 1996, 8:477-486
- 8. Laskowski RA et al "PROCHECK: a program to check the stereochemical quality of protein structures". J Appl Cryst 1993, 26:283-291
- 9. Word JM et al, "Exploring steric constrains on protein mutations using MAGE / PROBE". Prot Sci 2000, 9:2251-2259
- 10. Word JM et al, "Asparagine and glutamine: using hydrogen atom contacts in the choice of side-chain amide orientation". J Mol Biol 1999, 285:1735-1747
- 11. Word JM et al, "Visualizing and quantifying molecular goodness-of-fit: small-probe contact dots with explicit hydrogens". J Mol Biol 1999, 285:1711-1733
- 12. Luthy R, McLachlan AD and Eisenberg D, "Secondary structure-based profiles: use of structure-conserving scoring tables in searching protein sequence databases for structural similarities". Proteins 1991, 18:229-239
- 13. Richardson DC, Richardson J S, "The kinemage: a tool for scientific communication". Prot Sci 1992, 1(1):3-9
- 14. Guntërt P, Mumenthaler, C & Wüthrich, K "Torsion angle dynamics for NMR structure calculation with the new program DYANA". J. Mol. Biol 1997, 273:283-298
- 15. Lovell SC et al, "Structure validation by Calpha geometry: phi,psi and Cbeta deviation". Proteins 2003, 50:437-450
- 16. Kabsch W, Sander C, "Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features". Biopolymers 1983, 22:2577-2637
- 17. Bagaria A, Jaravine, V, Huang YJ, Montelione, GT, and Guntert, P "Protein structure validation by generalized linear model root-mean-square deviation prediction". Protein Sci 2012) 21:229-238.



Protein Structure Validation Suite (PSVS)



Software Environment

Software for structure quality evaluation:

DSSP DsspCMBI-April-2000 pdbstat PdbStat-5.20.8 Version

AutoAssign Version 2.4.0 (uses only AVS scripts)

RPF analysis ASDP-2.3



Verify3D Version 1.0 corrected by Aneerban ProsaII Prosa2003
PROCHECK Version 3.5.4

MolProbity programs:

1999 cluster

clashlistcluster 1999 (corrected by Aneerban)

mage Version 6.35.040409
prekin Version 6.35.040406
reduce Version 2.14
probe Version 2.6

probe Version 2.6

Other Software:

PERL Version 5.16.3

convert ImageMagick 6.7.8
ps2pdf Ghostscript 9.25
htmldoc v1.8.28
gnuplot Version 4.6.2
jpegtopnm netpbm-progs 10.79.00
pnmcrop netpbm-progs 10.79.00 pnmtojpeg netpbm-progs 10.79.00

Information updated on October, 2021. RTT

08/03/22 6