



Structure Quality Analysis for NAME

Analyses performed for DAOP well-defined residues.

Procheck analysis, RMSD calculation and structure superimposition are based on DAOP with $S(\phi)+S(\psi)\geq 1.8$: 2A-99A, 104A-106A

NESG ID: NAME

PDB ID:

Deposition date:

Common Name:

Class:

Length (a.a.): 111

Organism:


SwissProt /

TrEMBL ID:

models: 5

Oligomerization: monomer

Molecular weight: 13095



Secondary Structure Elements:

alpha helices: 3A-16A, 19A-36A, 46A-59A, 62A-81A

beta strands:

FIDs deposited in the BMRB? no

Comparison of core atoms:

DAOP > 1.8 Å : A:2..A:99, A:104..A:106

FindCore2 : A:1..A:96

CYRANGE : 1..91

RMSD	All residues	Ordered residues ²	Selected residues ³
All backbone atoms	1.5 Å	0.9 Å	0.9 Å
All heavy atoms	1.8 Å	1.0 Å	1.0 Å

Ramachandran Plot Summary for selected residues³ from Procheck



Structure Quality Analysis for NAME

Most favoured regions

92.0%

Additionally allowed regions

7.8%

Generously allowed regions

0.2%

Disallowed regions

0.0%

Ramachandran Plot Summary for selected residues³ from Richardson Lab's MolProbity

Most favoured regions

95.4%

Allowed regions

4.2%

Disallowed regions

0.4%

[View plot](#) [View model summary](#)

Global quality scores

Program	Verify3D	ProsaII (-ve)	Procheck (phi-psi) ³	Procheck (all) ³	MolProbity Clashscore
Raw score	0.13	N/A	0.26	0.33	1.94
Z-score ¹	-5.30	N/A	1.34	1.95	1.19

Close Contacts and Deviations from Ideal Geometry (from PDB validation software)

Number of close contacts (within 1.6 Å for H atoms, 2.2 Å for heavy atoms): 0


RMS deviation for bond angles: 1.9 °

RMS deviation for bond lengths: 0.012 Å

¹ With respect to mean and standard deviation for a set of 252 X-ray structures < 500 residues, of resolution ≤ 1.80 Å, R-factor ≤ 0.25 and R-free ≤ 0.28; a positive value indicates a 'better' score

² Ordered residues (DAOP): 2A-99A, 104A-106A


³ Selected residues DAOP with S(phi)+S(psi) ≥ 1.8 : 2A-99A, 104A-106A






Structure Quality Analysis for NAME

S(psi) v/s Residue number




Procheck G-factor for phi-psi







Procheck G-factor for all dihedral angles



Verify3D score over window of 7 residues




Residual VdW violations from MolProbity





Structure Quality Analysis for NAME



Residue Plot of Ramachandran analysis(based on data from Richardson Lab's Molprobit)

References:

1. Bhattacharya A, Tejero R and Montelione GT, "Evaluating protein structures determined by structural genomics consortia". *Proteins* 2007, 66:778-795
2. Tejero R, Snyder D, Mao B, Aramini JM and Montelione GT, "PDBStat: a universal restraint converter and restraint analysis software package for protein NMR". *J Biomol NMR* 2013, 56:337-351
3. Luthy R, Bowie JU and Eisenberg D, "Assessment of protein models with three-dimensional profiles". *Nature* 1992, 356:83-85
4. Bowie JU, Luthy R and Eisenberg D, "A Method to identify protein sequences that fold into a known three-dimensional structure". *Science* 1991, 253:164-169
5. Sippl MJ, "Recognition of errors in three-dimensional structures of proteins". *Proteins* 1993, 17:355-362
6. Sippl MJ, "Calculation of conformation ensembles from potentials of mean force". *J Mol Biol* 1990, 213:859-883
7. Laskowski RA et al, "AQUA and PROCHECK_NMR: programs for checking the quality of proteins structures solved by NMR". *J Biomolec NMR* 1996, 8:477-486
8. Laskowski RA et al "PROCHECK: a program to check the stereochemical quality of protein structures". *J Appl Cryst* 1993, 26:283-291
9. Word JM et al, "Exploring steric constraints on protein mutations using MAGE / PROBE". *Prot Sci* 2000, 9:2251-2259
10. Word JM et al, "Asparagine and glutamine: using hydrogen atom contacts in the choice of side-chain amide orientation". *J Mol Biol* 1999, 285:1735-1747
11. Word JM et al, "Visualizing and quantifying molecular goodness-of-fit: small-probe contact dots with explicit hydrogens". *J Mol Biol* 1999, 285:1711-1733
12. Luthy R, McLachlan AD and Eisenberg D, "Secondary structure-based profiles: use of structure-conserving scoring tables in searching protein sequence databases for structural similarities". *Proteins* 1991, 18:229-239
13. Richardson DC, Richardson J S, "The kinemage: a tool for scientific communication". *Prot Sci* 1992, 1(1):3-9
14. Gunttört P, Mumenthaler, C & Wüthrich, K "Torsion angle dynamics for NMR structure calculation with the new program DYANA". *J. Mol. Biol* 1997, 273:283-298
15. Lovell SC et al, "Structure validation by Calpha geometry: phi,psi and Cbeta deviation". *Proteins* 2003, 50:437-450
16. Kabsch W, Sander C, "Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features". *Biopolymers* 1983, 22:2577-2637
17. Bagaria A, Jaravine, V, Huang YJ, Montelione, GT, and Guntert, P "Protein structure validation by generalized linear model root-mean-square deviation prediction". *Protein Sci* 2012) 21:229-238.



Protein Structure Validation Suite (PSVS)



Software Environment

Software for structure quality evaluation:

DSSP	DsspCMBI-April-2000
pdostat	PdbStat-5.20.8 Version
AutoAssign	Version 2.4.0 (uses only AVS scripts)
RPF analysis	ASDP-2.3
PDB validation	Version 8.061
Verify3D	Version 1.0 corrected by Aneerban
ProsaII	Prosa2003
PROCHECK	Version 3.5.4

MolProbit programs:

cluster	1999
clashlistcluster	1999 (corrected by Aneerban)
mage	Version 6.35.040409
prekin	Version 6.35.040406
reduce	Version 2.14
probe	Version 2.6

Other Software:

PERL	Version 5.16.3
convert	ImageMagick 6.7.8
ps2pdf	Ghostscript 9.25
htmldoc	v1.8.28
gnuplot	Version 4.6.2
jpegtopnm	netpbm-progs 10.79.00
pnmcrop	netpbm-progs 10.79.00
pnmtjpeg	netpbm-progs 10.79.00

Information updated on October, 2021. RTT