
Analyses performed for DAOP well-defined residues.

Procheck analysis, RMSD calculation and structure superimposition are based on DAOP with S(phi)+S(psi)>=1.8:8A-57A



Secondary Structure Elements:

alpha helices: 9A-25A, 29A-40A, 44A-55A

108

20

12640

beta strands:

NESG ID:

Deposition date: Common Name:

Length (a.a.):

Organism: SwissProt/ **TrEMBL ID:** # models:

Molecular

weight:

PDB ID:

Class:

FIDs deposited in the BMRB? no

Comparison of core atoms:

A:8..A:25, DAOP > 1.8 Å: A:27..A:57

> **FindCore2**: A:7..A:59 **CYRANGE:** 10..55

| RMSD All residues | Ordered residues ² | Selected residues ³ |
|-------------------|-------------------------------|--------------------------------|
|-------------------|-------------------------------|--------------------------------|

19.1 Å $0.6\,\mathrm{\AA}$ $0.6\,\mathrm{\AA}$ All backbone atoms 17.6 Å 1.3 Å 1.3 Å All heavy atoms

Ramachandran Plot Summary for selected residues³ from Procheck

Most favoured Additionally allowed regions

Generously allowed regions Disallowed regions regions

97.1% 0.0% 2.9% 0.0%

Ramachandran Plot Summary for selected residues³ from Richardson Lab's Molprobity

Most favoured regions Allowed regions Disallowed regions View plot View model summary

98.4% 1.5% 0.1%

Global quality scores

| Program | Verify3D | ProsaII (-ve) | Procheck (phi-psi) ³ | $Procheck\ (all)^3$ | MolProbity Clashscore |
|----------------------|----------|---------------|---------------------------------|---------------------|-----------------------|
| Raw score | 0.02 | 0.28 | 0.33 | 0.20 | 12.47 |
| Z-score ¹ | -7.06 | -1.53 | 1.61 | 1.18 | -0.61 |

Close Contacts and Deviations from Ideal Geometry (from PDB validation software)


14 Number of close contacts (within 1.6 & Aring for H atoms, 2.2 & Aring for heavy atoms): 0.6 ° RMS deviation for bond angles: $0.004 \, \text{Å}$ RMS deviation for bond lengths:


With respect to mean and standard deviation for a set of 252 X-ray structures < 500 residues, of resolution <= 1.80 Å, R-factor <= 0.25 and R-free <= 0.28; a positive value indicates a 'better' score

² Ordered residues (DAOP): 8A-57A

³ Selected residues DAOP with S(phi)+S(psi)>=1.8:8A-57A








08/03/22 3











Procheck G-factor for all dihedral angles





08/03/22 4


















08/03/22 5





Residue Plot of Ramachandran anlysis(based on data from Richardson Lab's Molprobity)

References:

- 1. Bhattacharya A, Tejero R and Montelione GT, "Evaluating protein structures determined by structural genomics consortia". Proteins 2007. 66:778-795
- 2. Tejero R, Snyder D, Mao B, Aramini JM and Montelione GT, "PDBStat: a universal restraint converter and restraint analysis software package for protein NMR". J Biomol NMR 2013, 56:337-351
- 3. Luthy R, Bowie JU and Eisenberg D, "Assessment of protein models with three-dimensional profiles". Nature 1992, 356:83-85
- 4. Bowie JU, Luthy R and Eisenberg D, "A Method to identify protein sequences that fold into a known three-dimensional structure". Science 1991, 253:164-169
- 5. Sippl MJ, "Recognition of errors in three-dimensional structures of proteins". Proteins 1993, 17:355-362
- 6. Sippl MJ, "Calculation of conformation ensembles from potentials of mean force". J Mol Biol 1990, 213:859-883
- 7. Laskowski RA et al, "AQUA and PROCHECK_NMR: programs for checking the quality of proteins structures solved by NMR". J Biomolec NMR 1996, 8:477-486
- 8. Laskowski RA et al "PROCHECK: a program to check the stereochemical quality of protein structures". J Appl Cryst 1993, 26:283-291
- 9. Word JM et al, "Exploring steric constrains on protein mutations using MAGE / PROBE". Prot Sci 2000, 9:2251-2259
- 10. Word JM et al, "Asparagine and glutamine: using hydrogen atom contacts in the choice of side-chain amide orientation". J Mol Biol 1999, 285:1735-1747
- 11. Word JM et al, "Visualizing and quantifying molecular goodness-of-fit: small-probe contact dots with explicit hydrogens". J Mol Biol 1999, 285:1711-1733
- 12. Luthy R, McLachlan AD and Eisenberg D, "Secondary structure-based profiles: use of structure-conserving scoring tables in searching protein sequence databases for structural similarities". Proteins 1991, 18:229-239
- 13. Richardson DC, Richardson J S, "The kinemage: a tool for scientific communication". Prot Sci 1992, 1(1):3-9
- 14. Guntërt P, Mumenthaler, C & Wüthrich, K "Torsion angle dynamics for NMR structure calculation with the new program DYANA". J. Mol. Biol 1997, 273:283-298
- 15. Lovell SC et al, "Structure validation by Calpha geometry: phi,psi and Cbeta deviation". Proteins 2003, 50:437-450
- 16. Kabsch W, Sander C, "Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features". Biopolymers 1983, 22:2577-2637
- 17. Bagaria A, Jaravine, V, Huang YJ, Montelione, GT, and Guntert, P "Protein structure validation by generalized linear model root-mean-square deviation prediction". Protein Sci 2012) 21:229-238.



Protein Structure Validation Suite (PSVS)



Software Environment

Software for structure quality evaluation:

DSSP DsspCMBI-April-2000 pdbstat PdbStat-5.20.8 Version

AutoAssign Version 2.4.0 (uses only AVS scripts)
RPF analysis ASDP-2.3
PDB validation Version 8.061

Version 1.0 corrected by Aneerban

PDB Various

Verify3D Version 1

Prosa2003 PROCHECK Version 3.5.4

MolProbity programs:

1999

clashlistcluster 1999 (corrected by Aneerban)

Version 6.35.040409 Version 6.35.040406 prekin

reduce Version 2.14 probe Version 2.6

Other Software:

PERL Version 5.16.3 ImageMagick 6.7.8 convert Ghostscript 9.25 ps2pdf

v1.8.28 htmldoc

gnuplot Version 4.6.2

netpbm-progs 10.79.00 jpegtopnm netpbm-progs 10.79.00 pnmcrop netpbm-progs 10.79.00 pnmtojpeg

Information updated on October, 2021. RTT