

Wellcure Pharmaceuticals Ltd

Package Insert
惠康藥業有限公司 藥品說明書

Unit 8, 15/F, Kwai Chung Industrial Building, Kwai Chung, N.T., Hong Kong | Tel: 2487-XXXX | Approval No: HK-46829

Name-CN (說明書名稱-中文)	鹽酸二甲雙胍片		
Generic Name (通用名稱)	Metformin Hydrochloride Tablets	Specification (規格)	500mg × 60 tablets
Manufacturer (廠商名稱)	Wellcure Pharmaceuticals Ltd 惠康藥業有限公司	Approval Number (批准文號)	HK-46829

Composition (成分)

Each film-coated tablet contains Metformin Hydrochloride 500mg. 每片含鹽酸二甲雙胍500毫克。

Description (性狀)

White, round, film-coated tablets. 白色圓形薄膜衣錠。

Indications and Usage (適應症/功能主治)

Indicated for the management of Type 2 Diabetes Mellitus, particularly in overweight patients, as an adjunct to diet and exercise. May be used as monotherapy or in combination with other oral hypoglycemic agents or insulin.
用於2型糖尿病的治療，尤其適用於超重患者，作為飲食及運動治療的輔助。可單獨使用或與其他口服降糖藥或胰島素合併使用。

Administration and Dosage (用法用量)

Oral administration with meals. Initial dose 500mg twice daily, may be gradually increased according to blood glucose response up to a maximum of 2000mg/day.
口服，隨餐服用。起始劑量為每日兩次，每次500毫克，可根據血糖反應逐漸增加，最高每日劑量不超過2000毫克。

Adverse Reactions (不良反應)

Common: nausea, diarrhoea, abdominal discomfort, metallic taste. Rare but serious: lactic acidosis, particularly in patients with renal impairment.
常見：噁心、腹瀉、腹部不適、口中金屬味。罕見但嚴重：乳酸性酸中毒，尤見於腎功能不全患者。

Contraindication (禁忌)

Severe renal impairment (eGFR < 30 mL/min), diabetic ketoacidosis, severe hepatic impairment, acute conditions with risk of tissue hypoxia.
嚴重腎功能不全（eGFR低於30 mL/min）、糖尿病酮症酸中毒、嚴重肝功能不全、有組織缺氧風險之急性病症。

Precautions (注意事項)

Use with caution in patients with renal impairment; monitor renal function regularly. Discontinue temporarily before and after administration of iodinated contrast media. Not recommended during pregnancy or lactation unless clinically indicated.
腎功能不全患者慎用，並須定期監測腎功能。使用含碘造影劑前後應暫停服用。除非臨床有需要，否則不建議於懷孕或哺乳期間使用。

Storage / Packaging / Validity (貯藏/包裝/有效期)	
Storage (貯藏)	Store below 25°C, protect from moisture 存放於25°C以下，防潮
Packaging (包裝)	Blister pack, 60 tablets per box 鋁塑包裝，每盒60片
Validity (有效期)	36 months 36個月
Executive Standard (執行標準)	Hong Kong Pharmaceutical Standard 香港藥典標準
Manufacturer Address (廠商地址)	Unit 8, 15/F, Kwai Chung Industrial Building, Kwai Chung, N.T., Hong Kong 香港新界葵涌葵涌工業大廈15樓8室

This document is for reference only. Please consult a physician or pharmacist before use. 本文件僅供參考，使用前請諮詢醫生或藥劑師。