

Spatiotemporal analysis of post-fire vegetation recovery in Bastrop County

Ajay Jayaraman, Asha Karmakar, Ethan McFarlin, Fiona Pollack, Pranav Rajan, Iris Xia, Sean Yang



1. BACKGROUND RESEARCH

Wildfires occur naturally in many forest ecosystems and promote species diversity by clearing flammable undergrowth, thinning older trees, accelerating the carbon cycle, and replenishing soil nutrients. The aftermath of these wildfires varies widely depending on environmental variables including species and soil composition, weather, and seed production. Considering the impracticalities of surveying large areas through field work, remote sensing through satellite imagery allows for a more holistic overview of the entire region.

2. THE BASTROP COUNTY WILDFIRE

In 2011, central Texas experienced its most extreme drought since the 1950s, as well as the hottest U.S. summer on record, with an average temperature of 86.8° F. These already-extreme conditions created an environment ideal for the propagation of a spreading conflagration. After the original burns were ignited from sparks from fallen power lines, the wildfire converged from three smaller fires into one major entity over a period of several days in early September 2011.



Fig. 1. False color composite images showing progression of vegetation recovery

Due to the winds of Tropical Storm Lee, the fire continued to spread for over a month, killing two people, destroying 96% of Bastrop State Park (home to the indigenous Loblolly Pine), and burning \$325 million worth of property before being fully extinguished on October 29, 2011.

3. RESEARCH OBJECTIVES

Although vegetation regrowth in the immediate aftermath of the wildfire is well-documented, longer periods of time are typically more effective in capturing the extended cycle of forest recovery. This investigation aims to explore the following topics:

- How can remote sensing techniques be used to evaluate the long-term effects of the Bastrop County wildfire and determine whether these trends are temporary or lasting?
- What are the relationships between burn severity and postfire vegetation recovery in the aftermath of a wildfire as tracked using NDVI and dNBR?
- What environmental factors influence the nature of post-fire vegetation dynamics?

4. DATA ACQUISITION AND PROCESSING


To survey the large Bastrop Complex fire region, we utilized and processed 18 Landsat 5 and Landsat 8 images from 2009 to 2018. Landsat measures multispectral surface reflectance in the Near Infrared (NIR), Short Wave Infrared (SWIR), and Visible Bands bands. We selected the dates for our data using the Landlook Viewer and USGS Earth Explorer based on the image quality, cloud cover, and time of year in relation to the date of the 2011 fire.

The Difference Normalized Burn Ratio (dNBR) and Normalized Difference Vegetation Index (NDVI) are common indices used to quantify burn severity and vegetation growth, respectively. dNBR is calculated with NIR and SWIR bands from Landsat while NDVI incorporates the NIR and Visible Red bands. We quantified burn severity using the dNBR index, revealing patterns regarding the progression and impact of the fire.

$$NBR = \frac{NIR - SWIR}{NIR + SWIR} \cdot 1,000 \quad dNBR = NBR_{Prefire} - NBR_{Postfire} \quad NDVI = \frac{NIR - Red}{NIR + Red} \cdot 1,000$$

Data was processed using the following computational methods in Python/QGIS: clipping the raw imagery to the burn scar region, calculating a map of dNBR and NDVI, defining thresholds for burn severity based on dNBR value frequencies, and generating statistics (mean and standard deviation) and graphs of NDVI for each burn severity.

Fig. 2. The dNBR heat map exhibits a vertical strip of high severity, indicating a southerly direction of the wind that propagated the flames. Pockets of varying burn severity are influenced by other factors including species, geographic context, and fire movement.



5. VEGETATION REGROWTH

The resulting plot clearly illustrates the gradual decline in prefire NDVI values due to the worsening Texas drought, leading up to the significant drop in NDVI due to the Bastrop Complex Fire in 2011. The general trend also depicts fluctuations due to varying drought and precipitation levels, including drops in 2015 and 2018 due to hotter and drier conditions. The 2011 drop in NDVI is followed by a significant increase, with levels in 2013 and onwards appearing comparable to prefire NDVI levels, indicating a rapid rate of recovery.




Fig. 3. Plot of mean NDVI over time per burn severity. Each of the colored lines corresponds to the defined burn severities of Unburned, Low, Moderate, and High Severity. The patterns displayed by the various burn severities are fairly similar to the general trend, but the Unburned and Low Severity lines appear to have many more fluctuations, indicating that they are more influenced by changes in heat and precipitation while more severely burned areas were more focused on regrowth.




Fig. 4. Graph of percent change in NDVI over time per burn severity class for three chosen clusters of time. A clear trend can be observed with the first two clusters that the higher severity levels experienced greater percent changes in NDVI, but the most recent cluster shows the four severity levels experiencing similar percent changes, suggesting a plateau point.

6. GROUND TRUTHING

To confirm the geographical features observed by the spectral analysis of Landsat bands, qualitative field data was collected across various sampling points in the Griffith League Scout Ranch. The variance of vegetation regrowth appears to be largely consistent with predictions of NDVI as classified by burn severity using satellite imagery.



Fig. 5. Sampling transect near edge of burn scar. Organized recovery efforts involving professional replanting were more effective than logging at expanding the area of regrowth, but clumps of natural regeneration experienced the most rapid rates of development in all.

7. CONCLUSIONS

Both the satellite analysis and field investigation validate the expected contrast of behavior between the burned versus unburned areas and the different levels of burn severity.

- Areas of higher burn severity tended to have greater changes in NDVI, although this may be temporary as the NDVI leveled off in more recent years.
- Mean NDVI over time indicated a rapid recovery, but the field study indicated that there are still drastic differences by burn severity in terms of the vegetation count and form. The longer rebound period can be attributed to the severity of the disaster.
- Natural and anthropogenic factors such as areas with targeted replanting, streams, and natural regeneration tended to recover faster than areas with post-fire logging.

8. FUTURE RESEARCH

The positive correlation between burn severity and vegetation regrowth opens up investigations regarding the severity of prescribed controlled burns best suited for maximizing ecosystem health. Similar ground-truthing methodologies may also provide scientists with avenues to explore the effects of species diversity on regrowth.

9. ACKNOWLEDGEMENTS

We would like to thank Teresa Howard of the University of Texas at Austin Center for Space Research for her mentorship, along with the Texas Space Grant Consortium for directing the Stem Enhancement in Earth Science (SEES) program.