

Math 55a: Honors Abstract Algebra

Homework Assignment #3 (16 September 2016):

Linear Algebra III

The expression δ_{ij} is called the Kronecker delta (after the mathematician Leopold Kronecker [1823–1891], who made more substantial contributions to mathematics than this).¹

— Corwin and Szczarba, *Calculus in Vector Spaces*, p.124

A bit more about the structure of infinite-dimensional vector spaces:

1. i) Prove that a vector space with a countable² spanning set over a countable field is countable.
ii) Prove that a vector space with a countable spanning set over any field does not have an uncountable linearly independent set.
2. Suppose V is a vector space and U a subspace with basis B_0 . Suppose that for some (finite) n we can extend B by n vectors to obtain a basis for V . Prove that if B is any basis for U , and B' any basis for V that contains B , then $\{v \in B' : v \notin B\}$ has cardinality n . [This is clear when B is also finite (equivalently: when U , and thus also V , is finite dimensional), once we've shown that the dimension is well-defined when finite; but the point is that the result still holds without that assumption.]

Some basics about linear transformations and their matrices:

- 3.–4. Solve Exercises B-11 (page 68) and D-9, 10, 16 (page 89) from Chapter 3 of the textbook. For B-11, if $S_1 \cdots S_n$ is injective, what if anything can be said of S_1, S_2, \dots, S_n ? For the other three exercises, note that “ $\mathcal{L}(V)$ ” is Axler’s abbreviation for “ $\mathcal{L}(V, V)$ ” (it is also known as $\text{End}(V) = \text{Hom}(V, V)$).
5. Let \mathcal{P}_n be the (**R**- or **C**-)vector space of polynomials of degree at most n , and $L : \mathcal{P}_n \rightarrow \mathcal{P}_n$ be the linear transformation taking any polynomial $P(x)$ to the polynomial
$$(L(P))(x) = (x - 3)P''(x)$$
(here P'' is the second derivative d^2P/dx^2). Exhibit a matrix for L relative to a suitable basis for \mathcal{P}_n , and determine the kernel, image, and rank of L .
6. Let V, W be arbitrary vector spaces over the same field. Show that, for any vector v in V , the evaluation map $E_v : \mathcal{L}(V, W) \rightarrow W$ defined by $E_v(L) = L(v)$ for all $L \in \mathcal{L}(V, W)$ is a linear transformation. If V, W are finite dimensional, what is the dimension of $\ker E_v$?

¹It is the (i, j) entry of an identity matrix, that is, $\delta_{ij} = 1$ if $i = j$ and 0 otherwise; also $\delta_{ij} = \varphi_j(v_i)$ where $(v_i)_{i=1}^n$ is a basis for a finite-dimensional vector space, and $(\varphi_j)_{j=1}^n$ its dual basis, see 3.96 on page 102.

²For us “countable” means “finite or countably infinite”.

7. Let V, W be vector spaces over the rational field \mathbf{Q} . Prove that a map $T : V \rightarrow W$ is linear if and only if $T(v + v') = Tv + Tv'$ for all $v, v' \in V$. (Cf. the italicized note to Exercise 9 on p.58 of the textbook.)

More about duality:

8. We saw that, for any vector spaces V, W , the dual of $V \oplus W$ is naturally identified with $V^* \oplus W^*$. What is the dual of $\oplus_{i \in I} V_i$? Use this to construct a vector space V over some field F such that V is not isomorphic with V^* .
9. Let x_0, \dots, x_m be distinct elements of F . Recall that the $m + 1$ vectors $v_i := (x_0^i, x_1^i, \dots, x_m^i)$ ($0 \leq i \leq m$) constitute a basis of F^{m+1} . Describe the dual basis.
10. Finally, suppose F is a finite field of q elements, and let e be a positive integer such that $2e < q$. Then we can regard \mathcal{P}_{q-2e} as a subspace of F^F by evaluation at the elements of F . Call this subspace U . Show that for any $v \in V$ there is at most one $u \in U$ that differs from v in fewer than e coordinates; that is, there is at most one polynomial $P \in \mathcal{P}_{q-2e}$ such that $P(x) \neq v_x$ holds for fewer than e elements $x \in F$. Suppose such P exists and is nonzero, and let $d < e$ be the number of $x \in F$ such that $P(x) \neq v_x$. Prove that d is also the smallest integer n such that the intersection of \mathcal{P}_{q-2e+d} with the vector space

$$\mathcal{P}_d v := \{Pv : P \in \mathcal{P}_d\}$$

contains some vector $w \neq 0$. Explain how this can be used to recover u in fewer than q calculations of such w .

(The point of this is that each such calculation can be done “in polynomial time” [i.e. there exist C and k such that the calculation requires at most Cq^k field operations in F , regardless of what q, e, v might be — one way to do this is “Gaussian elimination”]; while trying all e -element subsets of F certainly cannot be done in polynomial time. Of course the exceptional case $u = P = 0$ *can* be detected in polynomial time.)

Problem set is due Friday, Sep. 23 in class.