

Math 55a, Fall 2004

Fourth Assignment, due October 19

1. Let (X, \mathcal{T}_X) , (Y, \mathcal{T}_Y) be topological spaces, \mathcal{S}_Y a sub-base for the topology \mathcal{T}_Y , and $F : X \rightarrow Y$ a map between the two spaces. By definition, a *neighborhood* of a point in a topological space is a set – not necessarily an open set! – whose interior contains the point in question. By definition, F is continuous at a given point $x \in X$ if, for every neighborhood $N_{F(x)}$ of $F(x)$, there exists a neighborhood N_x of x such that $F(N_x) \subset N_{F(x)}$. Show that the following conditions on F are equivalent:

- i) F is continuous (i.e., $F^{-1}(U) \in \mathcal{T}_X$ for every $U \in \mathcal{T}_Y$).
- ii) $F^{-1}(S)$ is closed for every closed set $S \subset Y$.
- iii) $F^{-1}(V) \in \mathcal{T}_X$ for every $V \in \mathcal{S}_Y$.
- iv) F is continuous at every point $x \in X$.

2. In this problem, \mathcal{T}_{reg} denotes the usual topology on the real line \mathbb{R} , and \mathcal{T}_{cof} the cofinite topology on \mathbb{R} – the topology for which a set, other than \mathbb{R} itself, is closed if and only if it is finite or empty.

- a) Is \mathcal{T}_{cof} a Hausdorff topology?
- b) Characterize the sequences in \mathbb{R} which are convergent with respect to the \mathcal{T}_{cof} topology, in as simple terms as possible.
- c) Characterize the functions $f : \mathbb{R} \rightarrow \mathbb{R}$ which are continuous with respect to the topology \mathcal{T}_{cof} on the domain and the topology \mathcal{T}_{reg} on the range, in as simple terms as possible.

3. Recall the properties of \mathbb{Q}_p and \mathbb{Z}_p which were established in the last assignment. Define the ring of adeles, \mathbb{A} , as follows. As a set, \mathbb{A} is contained in the Cartesian product of \mathbb{R} and the \mathbb{Q}_p , for all primes p :

$$\mathbb{A} = \{ a \in \mathbb{R} \times \prod_p \mathbb{Q}_p \mid a_p \in \mathbb{Z}_p \text{ for all but finitely many primes } p \}.$$

Equivalently, $\mathbb{A} = \cup_S \mathbb{A}_S$ is the union, extended over all finite sets S of prime numbers, of

$$\mathbb{A}_S = \{ a \in \mathbb{R} \times \prod_p \mathbb{Q}_p \mid a_p \in \mathbb{Z}_p \text{ for all } p \notin S \} \cong \mathbb{R} \times \prod_{p \in S} \mathbb{Q}_p \times \prod_{p \notin S} \mathbb{Z}_p.$$

Equip each $\mathbb{A}_S \cong \mathbb{R} \times \prod_{p \in S} \mathbb{Q}_p \times \prod_{p \notin S} \mathbb{Z}_p$ with the product topology, and decree that a subset $U \subset \mathbb{A}$ is open if and only if $U \cap \mathbb{A}_S$ is open for every S . Define the addition and multiplication of adeles component-by-component.

- a) Show that \mathbb{A} is a locally compact Hausdorff space (a topological space is locally compact if every point has a compact neighborhood).
- b) Show that addition and multiplication, viewed as maps from $\mathbb{A} \times \mathbb{A}$ to \mathbb{A} , are continuous.