

QUIZ • 10 MIN

✓

Congratulations! You passed!

TO PASS 80% or higher

Keep Learning

GRADE
100%

Octave/Matlab Tutorial

Octave/Matlab Tutorial

LATEST SUBMISSION GRADE

100%

✓ Submit your assignment

DUE Sep 30, 2:59 PM SGT

ATTEMPTS 3 every 8 hours

Try again

✓ Receive grade

TO PASS 80% or higher

1. Suppose I first execute the following in Octave/Matlab:

1 / 1 point

1 A = [1 2; 3 4; 5 6];
2 B = [1 2 3; 4 5 6];

Grade

100%

View Feedback

We keep your highest score

Which of the following are then valid commands? Check all that apply. (Hint: A' denotes the transpose of A.)



✓ Correct

2.

Let $A = \begin{bmatrix} 16 & 2 & 3 & 13 \\ 5 & 11 & 10 & 8 \\ 9 & 7 & 6 & 12 \\ 4 & 14 & 15 & 1 \end{bmatrix}$.

1 / 1 point

Which of the following indexing expressions gives $B = \begin{bmatrix} 16 & 2 \\ 5 & 11 \\ 9 & 7 \\ 4 & 14 \end{bmatrix}$? Check all that apply.

✓ Correct

3. Let A be a 10x10 matrix and x be a 10-element vector. Your friend wants to compute the product Ax and writes the following code:

1 / 1 point

1 v = zeros(10, 1);
2 for i = 1:10
3 for j = 1:10
4 v(i) = v(i) + A(i, j) * x(j);
5 end
6 end

How would you vectorize this code to run without any FOR loops? Check all that apply.

✓ Correct

4. Say you have two column vectors v and w , each with 7 elements (i.e., they have dimensions 7x1). Consider the following code:

1 / 1 point

1 z = 0;
2 for i = 1:7
3 z = z + v(i) * w(i)
4 end

Which of the following vectorizations correctly compute z? Check all that apply.

✓ Correct

5. In Octave/Matlab, many functions work on single numbers, vectors, and matrices. For example, the sin function when applied to a matrix will return a new matrix with the sin of each element. But you have to be careful, as certain functions have different behavior. Suppose you have an 7x7 matrix X . You want to compute the log of every element, the square of every element, add 1 to every element, and divide every element by 4. You will store the results in four matrices, A, B, C, D . One way to do so is the following code:

1 / 1 point

1 for i = 1:7
2 for j = 1:7
3 A(i, j) = log(X(i, j));
4 B(i, j) = X(i, j) ^ 2;
5 C(i, j) = X(i, j) + 1;
6 D(i, j) = X(i, j) / 4;
7 end
8 end

Which of the following correctly compute A, B, C , or D ? Check all that apply.

✓ Correct