



Math for the people, by the people.

## contractive maps are uniformly continuous

Canonical name	ContractiveMapsAreUniformlyContinuous
Date of creation	2013-03-22 13:46:28
Last modified on	2013-03-22 13:46:28
Owner	mathcam (2727)
Last modified by	mathcam (2727)
Numerical id	6
Author	mathcam (2727)
Entry type	Theorem
Classification	msc 54A20

**Theorem** A contraction mapping is uniformly continuous.

**Proof** Let  $T : X \rightarrow X$  be a contraction mapping in a metric space  $X$  with metric  $d$ . Thus, for some  $q \in [0, 1)$ , we have for all  $x, y \in X$ ,

$$d(Tx, Ty) \leq qd(x, y).$$

To prove that  $T$  is uniformly continuous, let  $\varepsilon > 0$  be given. There are two cases. If  $q = 0$ , our claim is trivial, since then for all  $x, y \in X$ ,

$$d(Tx, Ty) = 0 < \varepsilon.$$

On the other hand, suppose  $q \in (0, 1)$ . Then for all  $x, y \in X$  with  $d(x, y) < \varepsilon/q$ , we have

$$d(Tx, Ty) \leq qd(x, y) < \varepsilon.$$

In conclusion,  $T$  is uniformly continuous.  $\square$

The result is stated without proof in [?], pp. 221.

## References

- [1] W. Rudin, *Principles of Mathematical Analysis*, McGraw-Hill Inc., 1976.