



planetmath.org

Math for the people, by the people.

uncountable Polish spaces contain Cantor space

Canonical name	UncountablePolishSpacesContainCantorSpace
Date of creation	2013-03-22 18:48:33
Last modified on	2013-03-22 18:48:33
Owner	gel (22282)
Last modified by	gel (22282)
Numerical id	6
Author	gel (22282)
Entry type	Theorem
Classification	msc 54E50
Related topic	PolishSpace

Cantor space is an example of a compact and uncountable Polish space. In fact, every uncountable Polish space contains Cantor space, as stated by the following theorem.

Theorem. *Let X be an uncountable Polish space. Then, it contains a subset S which is homeomorphic to Cantor space.*

For example, the set \mathbb{R} of real numbers contains the <http://planetmath.org/CantorSetCantor> middle thirds set. Note that, being homeomorphic to Cantor space, S must be a compact and hence closed subset of X . The result is trivial in the case of Baire space \mathcal{N} , in which case we may take S to be the set of all $s \in \mathcal{N}$ satisfying $s_n \in \{1, 2\}$ for all n . Then, for any uncountable Polish space X there exists a continuous and one-to-one function $f: \mathcal{N} \rightarrow X$ (see <http://planetmath.org/InjectiveImagesOfBaireSpacehere>). Then f gives a continuous bijection from S to $f(S)$. The <http://planetmath.org/InverseFunctionThe> function theorem implies that f is a homeomorphism between S and $f(S)$ and, therefore, $f(S)$ is homeomorphic to Cantor space.