



planetmath.org

Math for the people, by the people.

examples of locally compact and not locally compact spaces

Canonical name	ExamplesOfLocallyCompactAndNotLocallyCompactSpaces
Date of creation	2013-03-22 12:48:36
Last modified on	2013-03-22 12:48:36
Owner	AxelBoldt (56)
Last modified by	AxelBoldt (56)
Numerical id	18
Author	AxelBoldt (56)
Entry type	Example
Classification	msc 54D45
Related topic	TopologicalSpace

Examples of locally compact spaces include:

- The Euclidean spaces \mathbb{R}^n with the standard topology: their local compactness follows from the Heine-Borel theorem. The <http://planetmath.org/Complexcomplexplane> \mathbb{C} carries the same topology as \mathbb{R}^2 and is therefore also locally compact.
- All topological manifolds are locally compact since locally they look like Euclidean space.
- Any closed or open subset of a locally compact space is locally compact. In fact, a subset of a locally compact Hausdorff space X is locally compact if and only if it is the <http://planetmath.org/SetDifference> of two closed subsets of X (equivalently: the intersection of an open and a closed subset of X).
- The space of <http://planetmath.org/PAdicIntegers> p -adic rationals is homeomorphic to the Cantor set minus one point, and since the Cantor set is compact as a closed bounded subset of \mathbb{R} , we see that the p -adic rationals are locally compact.
- Any discrete space is locally compact, since the singletons can serve as compact neighborhoods.
- The long line is a locally compact topological space.
- If you take any unbounded totally ordered set and equip it with the left order topology (or right order topology), you get a locally compact space. This space, unlike all the others we have looked at, is not Hausdorff.

Examples of spaces which are *not* locally compact include:

- The rational numbers \mathbb{Q} with the standard topology inherited from \mathbb{R} : each of its compact subsets has empty interior.
- All infinite-dimensional normed vector spaces: a normed vector space is finite-dimensional if and only if its closed unit ball is compact.
- The subset $X = \{(0, 0)\} \cup \{(x, y) \mid x > 0\}$ of \mathbb{R}^2 : no compact subset of X contains a neighborhood of $(0, 0)$.