



planetmath.org

Math for the people, by the people.

graph theorems for topological spaces

Canonical name	GraphTheoremsForTopologicalSpaces
Date of creation	2013-03-22 19:15:09
Last modified on	2013-03-22 19:15:09
Owner	joking (16130)
Last modified by	joking (16130)
Numerical id	7
Author	joking (16130)
Entry type	Theorem
Classification	msc 54C05
Classification	msc 26A15

We wish to show the relation between continuous maps and their graphs is closer than it may look. Recall, that if $f : X \rightarrow Y$ is a function between sets, then the set $\Gamma(f) = \{(x, f(x)) \in X \times Y\}$ is called *the graph of f* .

Proposition 1. If $f : X \rightarrow Y$ is a continuous map between topological spaces such that Y is Hausdorff, then the graph $\Gamma(f)$ is a closed subset of $X \times Y$ in product topology.

Proof. Indeed, we will show, that $Z = (X \times Y) \setminus \Gamma(f)$ is open. Let $(x, y) \in Z$. Then $f(x) \neq y$ and thus (since Y is Hausdorff) there exist open subsets $V_1, V_2 \subseteq Y$ such that $f(x) \in V_1$, $y \in V_2$ and $V_1 \cap V_2 = \emptyset$. Since f is continuous, then $U = f^{-1}(V_1)$ is open in X .

Note, that the condition $V_1 \cap V_2 = \emptyset$ implies, that $f(U) \cap V_2 = \emptyset$. Therefore $U \times V_2$ is a subset of Z . On the other hand this subset is open (since it is a product of two open sets) in product topology and $(x, y) \in U \times V_2$. This shows, that every point in Z belongs to Z together with a small neighbourhood, which completes the proof. \square

Unfortunately, the converse of this theorem is not true as we will see later. Nevertheless we can achieve similar result, if we assume a bit more about spaces:

Proposition 2. Let $f : X \rightarrow Y$ be a function, where X, Y are Hausdorff spaces with Y compact. If $\Gamma(f)$ is a closed subset of $X \times Y$ in product topology, then f is continuous.

Proof. Let $F \subseteq Y$ be a closed set. We will show that $f^{-1}(F)$ is also closed. Consider projections

$$\pi_Y : X \times Y \rightarrow Y; \quad \pi_X : X \times Y \rightarrow X.$$

They are both continuous and thus $\pi_Y^{-1}(F)$ is closed in $X \times Y$. Since $\Gamma(f)$ is also closed, then

$$Z = \pi_Y^{-1}(F) \cap \Gamma(f)$$

is closed in $X \times Y$. It is well known, that since Y is compact, then π_X is a closed map (this is easily seen to be equivalent to *the tube lemma*). Furthermore it is easy to see, that $\pi_X(Z) = f^{-1}(F)$ and the proof is complete. \square

Counterexample. Let \mathbb{R} denote the set of reals (with standard topology). Consider function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = 1/x$ and $f(0) = 0$. It is obvious, that f is discontinuous at $x = 0$, but also it can be easily checked, that $\Gamma(f)$ is closed in \mathbb{R}^2 . Note, that \mathbb{R} is not compact.