



planetmath.org

Math for the people, by the people.

example of continuous bijections which are
not homeomorphisms

Canonical name	ExampleOfContinuousBijectionsWhichAreNotHomeomorphisms
Date of creation	2013-03-22 18:54:31
Last modified on	2013-03-22 18:54:31
Owner	joking (16130)
Last modified by	joking (16130)
Numerical id	4
Author	joking (16130)
Entry type	Example
Classification	msc 54C05

Example 1. Assume that X is a topological space, which neither discrete nor antidiscrete. We will show that there are topological spaces Y and Z such that there are continuous bijections $X \rightarrow Y$ and $Z \rightarrow X$ which are not homeomorphisms.

Let $Y = Z = X$ as a sets but topology on Y is antidiscrete and on Z is discrete. Then obviously identity mappings $\text{id} : X \rightarrow Y$ and $\text{id} : Z \rightarrow X$ are continuous, but since X is neither discrete nor antidiscrete, these mappings are not homeomorphisms.

Example 2. Consider the function $f : [0, 1) \rightarrow S^1$ (here S^1 denotes the unit circle in a complex plane) defined by the formula $f(t) = e^{2\pi it}$. It is easy to see that f is a continuous bijection, but f is not a homeomorphism (because $[0, 1)$ is not compact).