



topology of the complex plane

Canonical name	TopologyOfTheComplexPlane
Date of creation	2013-03-22 13:38:40
Last modified on	2013-03-22 13:38:40
Owner	matte (1858)
Last modified by	matte (1858)
Numerical id	8
Author	matte (1858)
Entry type	Definition
Classification	msc 54E35
Classification	msc 30-00
Related topic	IdentityTheorem
Related topic	PlacesOfHolomorphicFunction
Defines	open disk
Defines	accumulation point
Defines	interior point
Defines	open
Defines	closed
Defines	bounded
Defines	compact

The usual topology for the complex plane \mathbb{C} is the topology induced by the metric

$$d(x, y) := |x - y|$$

for $x, y \in \mathbb{C}$. Here, $|\cdot|$ is the <http://planetmath.org/ModulusOfComplexNumbercomplexmodulus>.

If we identify \mathbb{R}^2 and \mathbb{C} , it is clear that the above topology coincides with topology induced by the Euclidean metric on \mathbb{R}^2 .

Some basic topological concepts for \mathbb{C} :

1. The open balls

$$B_r(\zeta) = \{z \in \mathbb{C} : |z - \zeta| < r\}$$

are often called *open disks*.

2. A point ζ is an *accumulation point* of a subset A of \mathbb{C} , if any open disk $B_r(\zeta)$ contains at least one point of A distinct from ζ .
3. A point ζ is an *interior point* of the set A , if there exists an open disk $B_r(\zeta)$ which is contained in A .
4. A set A is *open*, if each of its points is an interior point of A .
5. A set A is *closed*, if all its accumulation points belong to A .
6. A set A is *bounded*, if there is an open disk $B_r(\zeta)$ containing A .
7. A set A is *compact*, if it is closed and bounded.