



planetmath.org

Math for the people, by the people.

global versus local continuity

Canonical name	GlobalVersusLocalContinuity
Date of creation	2013-03-22 19:09:07
Last modified on	2013-03-22 19:09:07
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	4
Author	CWoo (3771)
Entry type	Result
Classification	msc 54C05
Classification	msc 26A15

In this entry, we establish a very basic fact about continuity:

Proposition 1. *A function $f : X \rightarrow Y$ between two topological spaces is continuous iff it is continuous at every point $x \in X$.*

Proof. Suppose first that f is continuous, and $x \in X$. Let $f(x) \in V$ be an open set in Y . We want to find an open set $x \in U$ in X such that $f(U) \subseteq V$. Well, let $U = f^{-1}(V)$. So U is open since f is continuous, and $x \in U$. Furthermore, $f(U) = f(f^{-1}(V)) = V$.

On the other hand, if f is not continuous at $x \in X$. Then there is an open set $f(x) \in V$ in Y such that no open sets $x \in U$ in X have the property

$$f(U) \subseteq V. \tag{1}$$

Let $W = f^{-1}(V)$. If W is open, then W has the property (1) above, a contradiction. Since W is not open, f is not continuous. \square