



Math for the people, by the people.

alternative characterization of ultrafilter

Canonical name	AlternativeCharacterizationOfUltrafilter
Date of creation	2013-03-22 14:42:20
Last modified on	2013-03-22 14:42:20
Owner	yark (2760)
Last modified by	yark (2760)
Numerical id	13
Author	yark (2760)
Entry type	Theorem
Classification	msc 54A20

Let X be a set. A filter \mathcal{F} over X is an ultrafilter if and only if it satisfies the following condition: if $A \coprod B = X$ (see disjoint union), then either $A \in \mathcal{F}$ or $B \in \mathcal{F}$.

This result can be generalized somewhat: a filter \mathcal{F} over X is an ultrafilter if and only if it satisfies the following condition: if $A \cup B = X$ (see union), then either $A \in \mathcal{F}$ or $B \in \mathcal{F}$.

This theorem can be extended to the following two propositions about finite unions:

1. A filter \mathcal{F} over X is an ultrafilter if and only if, whenever A_1, \dots, A_n are subsets of X such that $\coprod_{i=1}^n A_i = X$ then there exists exactly one i such that $A_i \in \mathcal{F}$.
2. A filter \mathcal{F} over X is an ultrafilter if and only if, whenever A_1, \dots, A_n are subsets of X such that $\bigcup_{i=1}^n A_i = X$ then there exists an i such that $A_i \in \mathcal{F}$.