

1. Koshy B, Srinivasan M, Bose A, John SM, et al. 2020. Developmental Trends in Early Childhood and Their Predictors from an Indian Birth Cohort. BMC Public Health. (In Review)
Doi: 10.21203/rs.3.rs-46054/v1
2. Lewis FI, Guga G, Mdoe P, Mduma E, et al. 2020. Introducing a drift and diffusion framework for childhood growth research. Gates Open Research 4:71.
Doi: 10.12688/gatesopenres.13123.1
3. Mahfuz M, Hasan SMT, Alam MA, Das, S, et al. 2020. Aflatoxin exposure was not associated with childhood stunting: results from a birth cohort study in a resource-poor setting of Dhaka, Bangladesh. Pubic Health and Nutrition.
Doi: 10.1017/S1368980020001421
4. Antiporta DA, Ambikapathi R, Bose A, Maciel B, et al. 2020. Micronutrient intake and the probability of nutrient adequacy among children 9–24 months of age: results from the MAL-ED birth cohort study. Public Health and Nutrition.
DOI: <https://doi.org/10.1017/S1368980020000877>
5. Richard SA, McCormick BJJ, Murray-Kolb LE, Bessong P, et al. 2020. Influences on catch-up growth using relative versus absolute metrics: Evidence from the MAL-ED cohort study. BMC Pub Health.
Doi: 10.21203/rs.3.rs-28471/v1
6. Das S, Fahim SM, Alam, MdA, Mahfuz M, et al. 2020. Not water, sanitation and hygiene practice, but timing of stunting is associated with recovery from stunting at 24 months, results from a multi-country birth cohort study. Pub Health Nut.
Doi: 10.1017/S136898002000004X
7. Sanchez JJ, Alam A, Stride CB, Haque A, et al. 2020. Campylobacter infection and household factors are associated with childhood growth in urban Bangladesh, an analysis of the MAL-ED study. PLOS Neg Trop Dis.
Doi: 10.1371/journal.pntd.0008328
8. Giallourou N, Fardus-Reid F, Panic G, Veselkov K, McCormick BJJ, et al. 2020. Metabolic maturation in the first 2 years of life in resource-constrained settings and its association with postnatal growths. Science Advances, vol. 6, no. 15, eaay5969.
Doi: 10.1126/sciadv.aay5969
9. Drago F, Scharf RJ, Maphula A, Nyathi E, et al. 2020. Psychosocial and environmental determinants of child cognitive development in rural South Africa and Tanzania, findings from the MAL-ED cohort. BMC Public Health. 20:505.
Doi: 10.1186/s12889-020-08598-5
10. Koshy B, Srinivasan M, Zachariah SM, Karthikeyan AS, et al. 2020. Body iron and lead status in early childhood and its effects on development and cognition: a longitudinal study from urban Vellore. Public Health Nutr.
Doi: 10.1017/S136898001900462
11. Rogawski McQuade ET, Liu J, Kang G, et al. 2020. Protection from natural immunity against enteric infections and etiology-specific diarrhea in a longitudinal birth cohort. J Infect Dis.
Doi: 10.1093/infdis/jiaa031
12. Alam MA, Richard SA, Fahim SM, et al. 2020. Impact of early-onset persistent stunting on cognitive development at 5 years of age, Results from a multi-country cohort study. PLOS One
Doi: 10.1371/journal.pone.0227839
13. Costa PN, Soares AM, Filho JQ, Junior FS, et al. 2020. Dietary intake from complementary feeding is associated with intestinal barrier function and environmental enteropathy in Brazilian children from the MAL-ED cohort study. Br J Nutr.
Doi: 10.1017/S0007114520000215

14. Colston J, Olortegui MP, Zaitchik B, et al. 2020. Pathogen-Specific Impacts of the 2011–2012 La Niña-Associated Floods on Enteric Infections in the MAL-ED Peru Cohort: A Comparative Interrupted Time Series Analysis. *Environ Res and Pub Health*.
Doi: 10.3390/ijerph17020487
15. Chhabra P, Rouhani S, Browne H, Yori, PP, et al. 2020. Homotypic and heterotypic protection and risk of re-infection following natural norovirus infection in a highly endemic setting. *Clin Infect Dis*.
Doi.org/10.1093/cid/ciaa019
16. Mahfuz M, Murray-Kolb LE, Hasan SMT, et al. 2019. Why Do Children in Slums Suffer from Anemia, Iron, Zinc, and Vitamin A Deficiency? Results from a Birth Cohort Study in Dhaka. *Nutrients*.
Doi: 10.3390/nu11123025
17. Colston JM, Francois R, Pisanic N, et al. 2019. Effects of Child and Maternal Histo-Blood Group Antigen Status on Symptomatic and Asymptomatic Enteric Infections in Early Childhood. *JID* 220:151–62.
Doi: 10.1093/infdis/jiz072
18. Pisanic N, Sarah-Blythe B, Colquechagua FD, et al. 2019. Minimally Invasive Saliva Testing to Monitor Norovirus Infection in Community Settings. *JID* 219:1234–42.
Doi: 10.1093/infdis/jiy638
19. Tofail F, Islam MM, Mahfuz M, Ashraful Alam M., Aktar S, Haque R, et al. 2019. Association of vitamin D nutrition with neuro-developmental outcome of infants of slums in Bangladesh. *PLoS ONE*
Doi: 10.1371/journal.pone.0221805
20. Tanih NF, Bolick DT, Samie A, Nyathi E, et al. 2019. Prevalence of Virulence Genes in Enterococcal *Escherichia coli* Isolates from Young Children from Rural South Africa. *AJTMH*.
Doi:10.4269/ajtmh.19-0192
21. Haque MDA, Platts-Mills JA, Mduma E, Bodhidatta L, et al. 2019. Determinants of *Campylobacter* infection and association with growth and enteric inflammation in children under 2 years of age in low-resource settings. *Nature Scientific Reports*.
doi.org/10.1038/s41598-019-53533-3
22. Rouhani S, Griffin NW, Yori PP, Gehrig JL, et al. 2019. Diarrhea as a potential cause and consequence of reduced gut microbial diversity among undernourished children in Peru. *Clin Infect Dis*.
Doi.org/10.1093/cid/ciz905
23. Rouhani S, Griffin NW, Yori PP, Olortegui BS, et al. 2019 Gut microbiota features associated with *Campylobacter* burden and postnatal linear growth deficits in a Peruvian birth cohort. *Clin Infect Dis*.
Doi.org/10.1093/cid/ciz906
24. Raman AS, Gehrig JL, Venkatesh S, Chang HW, et al. 2019. A sparse covarying unit that describes healthy and impaired human gut microbiota development. *Science*.
Doi.org/10.1126/science.aau4735
25. McCormick BJJ, Murray-Kolb LE, Lee GO, Schulze KJ, et al. 2019. Intestinal permeability and inflammation mediate the association between nutrient density of complementary foods and biochemical measures of micronutrient status in young children: results from the MAL-ED study. *Am. J. Clin. Nutr.*
Doi: 10.1093/ajcn/nqz151
26. Nahar B, Hossain M, Mahfuz M, et al. 2019. Early childhood development and stunting: Findings from the MAL-ED birth cohort study in Bangladesh. *Maternal and Child Nutrition*.
doi.org/10.1111/mcn.12864

27. Ruan-lu L, Pendergast LL, Rasheed M, Tofail F, et al. 2019. Assessing early childhood fluid reasoning in low- and middle-income nations: validity of the Wechsler Preschool and Primary Scale of Intelligence across seven MAL-ED sites. *Journal of Psychoeducational Assessment*.
Doi: 10.1177/0734282919850040
28. McCormick BJJ, Richard SA, Caulfield LE, et al. 2019. Early life child micronutrient status, maternal reasoning, and a nurturing household environment have persistent influences on child cognitive development at age 5 years: results from MAL-ED. *J Nutr.* epub Jun 4.
doi: 10.1093/jn/nxz055.
29. Colston JM, Francois R, Pisanic N, Yori PP, McCormick BJJ, et al. 2019. Effect of child and maternal Histo Blood Group Antigen status on symptomatic and asymptomatic enteric infections in early childhood. *J. Infect. Dis.*
doi: 10.1093/infdis/jiz072
30. Fleece M, Nshama R, Walongo T, Kimathi C, et al. 2019. Longitudinal assessment of antibiotic resistance in fecal *Escherichia coli* in Tanzanian children. *Am J Trop Med and Hygiene*.
Doi.org/10.4269/ajtmh.18-0789
31. Richard SA, McCormick BJJ, Murray-Kolb LE, Lee, GO, et al. 2019. Enteric dysfunction and other factors associated with attained size at five years: MAL-ED birth cohort study findings. *Am J Clin Nutr.*
doi: 10.1093/ajcn/nqz004
32. Rogawski McQuade ET, Clark S, Bayo E, Scharf RJ, et al. 2018. Seasonal food insecurity in Haydom, Tanzania, is associated with low birthweight and acute malnutrition: results from the MAL-ED study. *Am J Trop Med and Hygiene*.
doi: 10.4269/ajtmh.18-0547
33. Pendergast LC, Schaefer BA, Murray-Kolb LE, Svensen E, et al. 2018. Assessing development across cultures, invariance of the Bayley-III scales across seven international MAL-ED sites. *School Psychology Quarterly*. 33: 604–614.
doi: 10.1037/spq0000264
34. Fahim SM, Das S, Gazi A, et al. 2018. Association of intestinal pathogens with faecal markers of environmental enteric dysfunction among slum-dwelling children in the first 2 years of life in Bangladesh. *Trop. Med and Intern Health*. 23(11):1242-1250.
doi: 10.1111/tmi.13141
35. Pan WK, Seidman JC, Ali A, Hoest C, et al. 2019. Oral polio vaccine response in the MAL-ED birth cohort study: Considerations for polio eradication strategies. *Vaccine*. 7;37(2):352-365.
doi: 10.1016/j.vaccine.2018.05.080
36. Mahfuz M, Alam MA, Fahim SM, et al. 2018. Aflatoxin exposure in children living in Mirpur, Dhaka: data from MAL-ED companion study. *J Exposure Sci and Environ Epidemiol*.
doi: 10.1038/s41370-018-0066
37. Platts-Mills, JA, Liu, J, Rogawski, ET, et al. 2018. Use of quantitative molecular diagnostic methods to assess the aetiology, burden, and clinical characteristics of diarrhoea in children in low-resource settings: a reanalysis of the MAL-ED cohort study. *Lancet, Global Health*
doi: 10.1016/S2214-109X(18)30349-8
38. Rogawski ET, Lij, J, Platts-Mills, JA, et al. 2018. Use of quantitative molecular diagnostic methods to investigate the effect of enteropathogen infections on linear growth in children in low-resource settings: longitudinal analysis of results from the MAL-ED cohort study. *Lancet, Global Health*. 6: e1319–28
doi: 10.1016/S2214-109X(18)30351-6

39. Maciel BLL, Moraes ML, Soares AM, Cruz IFS, et al. 2018. Infant feeding practices and determinant variables for early complementary feeding in the first 8 months of life: results from the Brazilian MAL-ED cohort site. *Pub Health Nutr.* 21: 2462–2470.
doi: 10.1017/S136898001800099X
40. Ambikapathi R, Rothstein JD, Yori PP, et al. 2018. Food purchase patterns indicative of household food access insecurity, Children's dietary diversity and intake, and nutritional status using a newly developed and validated tool in the Peruvian Amazon. *Food Security.* June
doi: 10.1007/s12571-018-0815-2
41. Korpe PS, Valencia C, Haque R, Mahfuz M, et.al. 2018. Epidemiology and Risk Factors for Cryptosporidiosis in Children From 8 Low-income Sites: Results From the MAL-ED Study. *Clin. Infect. Dis.*
doi: 10.1093/cid/ciy355
42. Fahim SM, Das S, Sanin KI, et al. 2018. Association of fecal markers of environmental enteric dysfunction with zinc and iron status among children at first two years of life in Bangladesh. *AJTMH.* 99: 489 – 494.
doi: 10.4269/ajtmh.17-0985
43. MAL-ED Network Investigators. 2018. Early childhood cognitive development is affected by interactions among illness, diet, enteropathogens and the home environment: findings from the MAL-ED birth cohort study. *BMJ Glob Health,* 2018; 3:e000752.
doi: 10.1136/bmjgh-2018-000752
44. Raihan MJ, Briskin E, Mahfuz M, et al. 2018. Examining the relationship between blood lead level and stunting, wasting and underweight – A cross-sectional study of children under 2 years-of-age in a Bangladeshi slum. *PLoS ONE* 13(5):e0197856
doi: 10.1371/journal.pone.0197856.
45. Colston JM, Ahmed T, Mahopo C, et al. 2018. Evaluating meteorological data from weather stations, and from satellites and global models for a multi-site epidemiological study. *Env Res.* 165:91-105.
doi: 10.1016/j.envres.2018.02.027
46. Scharf RJ, Rogawski ET, Murray-Kolb L, et al. 2018 Early childhood growth and cognitive outcomes: Findings from the MAL-ED study. *Maternal and Child Nutrition.*
doi: 10.1111/mcn.12584
47. Morseth MS, Henjum S, Schwinger C, et al. 2018. Environmental enteropathy, micronutrient adequacy and length velocity in Nepalese children - the MAL-ED birth cohort study. *Journal of Pediatric Gastroenterology and Nutrition.*
doi: 10.1097/MPG.0000000000001990
48. Sanin KI, Islam MM, Mahfuz M, et al. 2018. Micronutrient adequacy is poor, but not associated with stunting between 12-24 months of age: A cohort study findings from a slum area of Bangladesh. *PLoS ONE* 13(3).
doi: 10.1371/journal.pone.0195072
49. Chen C, Mitchell NJ, Gratz J, Houpt ER, et al. 2018. Exposure to aflatoxin and fumonisin in children at risk for growth impairment in rural Tanzania, *Environment International,* 115:29-37.
doi: 10.1016/j.envint.2018.03.001
50. Ribe IG, Svensen E et al. 2018. Determinants of early child development in rural Tanzania. *Child Adolesc Psychiatry Ment Health,* 12:18.
doi: 10.1186/s13034-018-0224-5
51. Colston JM, Ahmed AMS, Soofi SB, el al. 2018. Seasonality and within-subject clustering of rotavirus infections in an eight-site birth cohort study. *Epidemiology and Infection.*
doi: 10.1017/S0950268818000304

52. Islam MM, Sanin, KI, Mahfuz M, et al. 2018 Risk factors of stunting among children living in an urban slum of Bangladesh: Findings of a prospective cohort study. *BMC Pub Health* 18: 197. doi: 10.1186/s12889-018-5101-x
53. Exum NG, Lee GO, Olórtegui MP, Yori PP, et al. 2018. A longitudinal study of household water, sanitation and hygiene characteristics and environmental enteropathy biomarkers in children less than 24 months in Iquitos Peru. *AJTMH*. doi: 10.4269/ajtmh.17-0464
54. Francois R, Yori PP, Rouhanni S, et al. 2018. The other *Campylobacters*: Not innocent bystanders in endemic diarrhea and dysentery in children in low income settings. *PLOS Neg Trop Dis*. doi: 10.1371/journal.pntd.0006200
55. Hanselman B, Ambikapathi A, Mduma E, et al. 2018. Associations of land, cattle and food security with infant feeding practices among a rural population living in Manyara, Tanzania. *BMC Public Health* 18:159 doi: 10.1186/s12889-018-5074-9
56. Olortegui MP, Rouhani S, Yori PP, et al. 2018. Astrovirus Infection and Diarrhea in 8 Countries. *Pediatrics*. 141(1):e20171326 doi: 10.1542/peds.2017-1326
57. Morseth MS, Torheim LE, Chandyo RK et al. 2017. Severely inadequate micronutrient intake among children 9-24 months in Nepal-The MAL-ED birth cohort study. *Matern Child Nutr*. doi: 10.1111/mcn.12552
58. Morseth MS, Torheim LE, Gebremariam MK, et al. 2017. Tracking of infant and young child feeding practices among 9- to 24-month-old children in Nepal: the MAL-ED Birth Cohort Study. *Public Health Nutr*. Published online 17Oct2017 doi: 10.1017/S1368980017002294.
59. Pendergast LL, Jones P, Scharf R, et al. 2017. Approach temperament across cultures: validity of the infant temperament scale in MAL-ED. *Int. J. of School and Educational Psychol*. Published online 30Oct2017 doi: 10.1080/21683603.2017.1356773
60. MAL-ED Investigators. 2017. The relationship between growth and illness, enteropathogens, and dietary intakes in the first two years of life: findings from the MAL-ED birth cohort study. *BMJ Glob Health* 2017;2:e000370 doi: 10.1136/bmjgh-2017-000370
61. Richard SA, McCormick BJJ, Seidman JC, et al. 2018. Relationships among common illness symptoms and the protective effect of breastfeeding in early childhood in MAL-ED, an eight-country cohort study. *AJTMH*. . doi: 10.4269/ajtmh.17-0457
62. MAL-ED Network Investigators. 2017. Childhood stunting in relation to the pre- and postnatal environment during the first 2 years of life: The MAL-ED longitudinal birth cohort study. *PLOS Medicine*. doi: 10.1371/journal.pmed.1002408
63. Lima AAM, Soares AM, Filho JQS, et al. 2017. Enteroaggregative *E. coli* subclinical infection and co-Infections and impaired child growth in the MAL-ED cohort study. *J Pediatr Gastroenterol Nutr*. 2017 Sep 12. doi: 10.1097/MPG.0000000000001717
64. Rogawski ET, Platts-Mills JA, Seidman JC, et al. 2017. Early antibiotic exposure in low-resource settings is associated with increased weight in the first two years of life. *J Pediatr Gastroenterol Nutr*. 65(3):350-356. doi: 10.1097/MPG.0000000000001640

65. Venkata Raghava Mohan VR, Karthikeyan R, Babji S, et al. 2017. Rotavirus infection and disease in a multisite birth cohort: results from the MAL-ED study. *JID* 216:305–16
doi: 10.1093/infdis/jix199
66. MAL-ED Investigators. 2017. Childhood stunting in relation to the pre- and post-natal environment during the first two years of life: The MAL-ED longitudinal birth cohort study. *PLoS Medicine*. Accepted August 2017
doi: 10.1371/journal.pmed.1002408
67. Rogawski ET, Guerrant RL, Havt A, et al. 2017. Epidemiology of enteroaggregative *Escherichia coli* infections and associated outcomes in the MAL-ED birth cohort. *PLoS Negl Trop Dis*. 11(7):e0005798
doi: 10.1371/journal.pntd.0005798
68. Ahmed AMS, Ahmed T, Long KZ, et al. 2017. Prevalence and risk factors of vitamin D insufficiency and deficiency among 6–24-month-old underweight and normal-weight children living in an urban slum of Bangladesh. *Public Health Nutrition*. 20:1718-28
doi: 10.1017/S1368980015003353.
69. Platts-Mills JA, Taniuchi M, Uddin MdJ, et al. 2017. Association between enteropathogens and malnutrition in children aged 6–23 mo in Bangladesh: a case-control study. *Am J Clin Nutr* 105(5):1132–8.
doi: 10.3945/ajcn.116.138800
70. Havt A, Lima IFN, Medeiros PHQS, et al. 2017. Prevalence and virulence gene profiling of enteroaggregative *Escherichia coli* in malnourished and nourished Brazilian children. *Diagn Microbiol Infect Dis* 89: 98-105
doi: 10.1016/j.diagmicrobio.2017.06.024
71. Exum NG, Kosek MN, Davis MF, et al. 2017. Surface sampling collection and culture methods for *Escherichia coli* in household environments with high fecal contamination. *Int. J. Environ. Res. Public Health* 14(8): 947
doi: 10.3390/ijerph14080947
72. Berendes D, Leon J, Kirby A, et al. 2017. Household sanitation is associated with lower risk of bacterial and protozoal enteric infections, but not viral infections and diarrhea, in a cohort study in a low-income urban neighborhood in Vellore, India. *TMIH*.
doi: 10.1111/tmi.12915.
73. Colston JM, Yori PP, Colantuoni E, et al. 2017. A methodologic framework for modeling and assessing biomarkers of environmental enteropathy as predictors of growth in infants: an example from a Peruvian birth cohort. *Am J Clin Nutr* 106:245–55.
doi: 10.3945/ajcn.116.151886
74. Ambikapathi R, Kosek M, Yori PP, et al. 2017. El Niño Southern Oscillation affects food consumption, intake, and dietary diversity in the Peruvian Amazon. *The FASEB Journal*. 31(1) Supplement 455.2.
75. Lee GO, McCormick BJJ, Seidman JC, et al. 2017. Infant nutritional status, feeding practices, enteropathogen exposure, socioeconomic status, and illness are associated with gut barrier function as assessed by the lactulose mannitol test in the MAL-ED birth cohort. *Am J Trop Med Hyg*. 97(1):281-290.
doi: 10.4269/ajtmh.16-0830
76. Ambikapathi R, Kosek M, Yori PP, et al. 2017. El Niño Southern Oscillation affects girls' nutrient intakes and adequacies in the Peruvian Amazon. *The FASEB Journal*. 31(1) Supplement 639.43.
77. Jones PC, Pendergast LL, Schaefer BA, et al. 2017. Measuring home environments across cultures: invariance of the HOME scale across eight international sites from the MAL-ED study. *J Sch Psychol*. 64:109-127.

- doi: 10.1016/j.jsp.2017.06.001
78. Alam MA, Mahfuz M Islam MM, et al. 2017. Contextual factors for stunting among children of age 6 to 24 months in an under-privileged community of Dhaka, Bangladesh. *Indian Pediatr.* 54:373-376. pii: S097475591600060
 79. Kosek MN, Lee GO, Guerrant RL, et al. 2017. Age and sex normalization of intestinal measures of enteropathy in MAL-ED. *J Pediatr Gastroenterol and Nutr.*
doi: 10.1097/MPG.0000000000001610
 80. Mushaphi LF, Mahopo TC, Nesamvuni CN, et al. 2017. Recommendations for infant feeding policy and programs in Dzimauli region, South Africa: results from the MAL-ED birth cohort. *Food and Nutrition Bulletin.*
doi: 10.1177/0379572117696662
 81. Moyo CS, Francis J, Bessong PO. 2017. Perceptions of community-based field workers on the effect of a longitudinal biomedical research project on their sustainable livelihoods. *BMC Public Health.* 17(1):267.
doi: 10.1186/s12889-017-4138-6
 82. DeFrancesco AS, Tanih NF, Samie A, et al. 2017. Antibiotic resistance patterns and beta-lactamase identification in *Escherichia coli* isolated from young children in rural Limpopo Province, South Africa: The MAL-ED cohort. *S Afr Med J.* 107(3):205-214.
doi:10.7196/SAMJ.2017.v107i3.12111
Guest Editorial: Bessong PO and Guerrant RL. 2017. Improving our understanding of antibiotic resistance: The relevance of surveillance at the population level. *S Afr Med J.* 107(3):167.
doi: 10.7196/SAMJ.2017.v107i3.12357
 83. Mitchell NJ, Hsu HH, Chandyo RK, et al. 2017. Aflatoxin exposure during the first 36 months of life was not associated with impaired growth in Nepalese children: An extension of the MAL-ED study. *PLoS One.* 12(2):e0172124.
doi: 10.1371/journal.pone.0172124
 84. The MAL-ED Network Investigators. 2017. Causal pathways from enteropathogens to environmental enteropathy: findings from the MAL-ED birth cohort study. *Ebiomedicine.* 18:109-117.
doi: 10.1016/j.ebiom.2017.02.024
 85. Lang DR. Developmental effects of early life exposure to enteric pathogens and other environmental factors. Old Herborn University Monograph #30, The Magnificent Microbiome - future aspects. Editors: Peter J. Heidt, Thomas C.G. Bosch, Tore Midtvedt, and Volker Rusch. Old Herborn University Foundation, Herborn, Germany: (2017) pp 115-132. https://www.old-herborn-university.de/wp-content/uploads/publications/books/OHUni_book_30.pdf
 86. Lee GO, Olortequi MP, Salas MS, et al. 2017. Environmental enteropathy is associated with cardiometabolic risk factors in Peruvian children. *J Dev Orig Health Dis.* 8(3):337-348.
doi: 10.1017/S2040174417000071
 87. Lima AA, Leite AM, Di Moura A, et al. 2017. Determinant Variables, Enteric pathogen burden, gut function, and immune-related inflammatory biomarkers associated with childhood malnutrition: a prospective case-control study in northeastern Brazil. *Pediatr Infect Dis J.*
doi: 10.1097/INF.0000000000001569
 88. Rogawski ET, Bartelt LA, Platts-Mills JA, et al. 2017. Determinants and impact of *Giardia* infection in the first 2 years of life in the MAL-ED birth cohort. *J Pediatric Infect Dis Soc.* 6(2):153-160.
doi: 10.1093/jpids/piw082
 89. Rogawski ET, Platts-Mills JA, Seidman JC, et al. 2017. Use of antibiotics in children younger than two years in eight countries: a prospective cohort study. *Bull World Health Organ.*

- doi: 10.2471/BLT.16.176123
Erratum in: *Bull World Health Organ.* 2017. 95(2):164.
90. Schwinger C, Fadnes LT, Shrestha SK, et al. 2017. Predicting undernutrition at age 2 years with early attained weight and length compared with weight and length velocity. *J Pediatr.* 182:127-132.e1. doi: 10.1016/j.jpeds.2016.11.013
 91. Mitchell NJ, Riley RT, Egner PA, et al. 2017. Chronic aflatoxin exposure in children living in Bhaktapur, Nepal: Extension of the MAL-ED study. *J Expo Sci Environ Epidemiol.* 27(1):106-111. doi: 10.1038/jes.2015.87
 92. McCormick BJJ, Lee GO, Seidman JC et al. 2017. Dynamics and trends in fecal biomarkers of gut function in children from 1–24 months in the MAL-ED study. *Am J Trop Med Hyg.* 96(2):465-472. doi: 10.4269/ajtmh.16-0496
 93. Rogawski EA, Platts-Mills JA, Seidman JC, et al. 2017. Diverse antibiotic use practices in the first two years of life across 8 sites in the MAL-ED birth cohort. *Bull World Health Organ* 95(1):49-61. doi: 10.2471/BLT.16.176123
 94. DeBoer MD, Scharf RJ, Leite AM, et al. 2017. Systemic inflammation, growth factors, and linear growth in the setting of infection and malnutrition. *Nutrition.* 33:248-253. doi: 10.1016/j.nut.2016.06.013
 95. Hoest C, Seidman JC, Lee G et al. 2016. Vaccine coverage and adherence to EPI schedules in eight resource poor settings in the MAL-ED cohort study. *Vaccine.* 35(3):443-451. doi: 10.1016/j.vaccine.2016.11.075
 96. Hossain MI, Haque R, Mondal D et al. 2016. Undernutrition, vitamin A and iron deficiency are associated with impaired intestinal mucosal permeability in young Bangladeshi children assessed by lactulose/mannitol test. *PLoS One* 11(12):e0164447. doi: 10.1371/journal.pone.0164447
 97. Rouhani S, Peñataro Yori P, Paredes Olortegui M, et al. 2016. Norovirus infection and acquired immunity in 8 countries: results from the MAL-ED study. *Clin Infect Dis.* 62(10):1210-7. doi: 10.1093/cid/ciw072
 98. Exum NG, Olortegui MP, Yori PP, et al. 2016. Floors and toilets: association of floors and sanitation practices with fecal contamination in Peruvian Amazon peri-urban households. *Environ Sci Technol.* 50(14):7373-81. doi: 10.1021/acs.est.6b01283
 99. Amour C, Gratz J, Mduma E, et al. 2016. Epidemiology and impact of *Campylobacter* infection in children in 8 low-resource settings: results from the MAL-ED study. *Clin Infect Dis.* 63(9):1171–1179. doi: 10.1093/cid/ciw542
 100. Guerrant RL, Leite AM, et al. 2016. Biomarkers of environmental enteropathy, inflammation, stunting, and impaired growth in children in Northeast Brazil. *PLoS One.* 11(9):e0158772. doi:10.1371/journal.pone.0158772.
 101. Lee GO, Richard SA, Kang G, et al. 2016. A comparison of diarrheal severity scores in the MAL-ED multisite community-based cohort study. *J Pediatr Gastroenterol Nutr.* 63(5):466-473. doi: 10.1097/MPG.0000000000001286
 102. Kosek, MN, Mduma E, et al. 2016. Plasma tryptophan and the kynurenine–tryptophan ratio are associated with the acquisition of statural growth deficits and oral vaccine underperformance in populations with environmental enteropathy. *Am. J. Trop. Med. Hyg.* 95(4):928–937. doi:10.4269/ajtmh.16-0037.
 103. Ambikapathi R, Kosek M, Lee GO, et al. 2016. How multiple episodes of exclusive breastfeeding impact estimates of exclusive breastfeeding duration: report from the eight-site MAL-ED birth

- cohort study. *Maternal and Child Nutrition*. 12:740-756.
doi: 10.1111/mcn.12352
104. Arndt MB, Rachardson BA, Ahmed T, et al. 2016. Fecal markers of environmental enteropathy and subsequent growth in Bangladeshi children. *Am J Trop Med Hyg*. 95(3):694-701.
doi: 10.4269/ajtmh.16-0098
 105. Prata, M, A Havt, et al. 2016. Comparisons between myeloperoxidase, lactoferrin, calprotectin and lipocalin-2, as fecal biomarkers of intestinal inflammation in malnourished children. *J. Translational Sci*. 2(2):134–139.
doi: 10.15761/JTS.1000130.
 106. McCormick BJJ and Lang D. 2016. Diarrheal disease and enteric infections in LMIC communities: how big is the problem? *Tropical Diseases, Travel Medicine and Vaccines*. 2:11.
doi: 10.1186/s40794-016-0028-7.
 107. Syed S, Ali A, and Duggan C. 2016. Environmental enteric dysfunction in children (Review). *J Pediatr Gastroenterol Nutr*. 63 (1):6-14.
doi: 10.1097/MPG.0000000000001147
 108. Choudhury N, Bromage S, Alam MA, et al. 2016. Intervention study shows suboptimal growth among children receiving a food supplement for five months in a slum in Bangladesh. *Acta Paediatrica*.
doi: 10.1111/apa.13528
 109. Mahfuz M, Alam MA, Islam MM, et al. 2016. Effect of micronutrient powder supplementation for two and four months on hemoglobin level of children 6–23 months old in a slum in Dhaka: a community based observational study. *BMC Nutrition*. 2:21.
doi: 10.1186/s40795-016-0061-y
 110. Ahmed AMS, Soares Magalhaes RJ, Long KZ, et al. 2016. Association of vitamin D status with incidence of enterotoxigenic, enteropathogenic and enteroaggregative *Escherichia coli* diarrhoea in children of urban Bangladesh. *TMIH*. 21(8): 973-984.
doi: 10.1111/tmi.12731
 111. Ahmed AMS, Magalhaes RJS, Ahmed T, et al. 2016. Vitamin-D status is not a confounder of the relationship between zinc and diarrhoea: a study in 6–24-month-old underweight and normal-weight children of urban Bangladesh. *Eur J of Clin Nutr*. 70:620-628.
doi: 10.1038/ejcn.2016.7
 112. Ahmed AMS, Ahmed T, Magalhaes RJS, Long KZ, et al. 2016. Association between serum vitamin D, retinol and zinc status, and acute respiratory infections in 6-24 months old underweight and normal weight children living in an urban slum in Bangladesh. *Epidemiology and Infection*. 144: 3494-3506.
doi: 10.1017/S0950268816001771
 113. Mayneris-Perxachs J, Lima AAM, Guerrant RL, et al. 2016. Urinary N-methylnicotinamide and β -aminoisobutyric acid predict catch-up growth in undernourished Brazilian children. *Nature Scientific Report*. 6:19780.
doi: 10.1038/srep19780.
 114. Liu J, Gratz J, Amour C, et al. 2016. Optimization of quantitative PCR methods for enteropathogen detection. *PLOS ONE*
doi: 10.1371/journal.pone.0158199
 115. Oria, RB et al. 2016. Early-life enteric infections: relation between chronic systemic inflammation and poor cognition in children. *Nutr Rev*. 74(6):374-86.
doi: 10.1093/nutrit/nuw008.

116. Platts-Mills JA, Babji S, Bodhidatta L, et al. 2015. Pathogen-specific burdens of community diarrhoea in developing countries: a multisite birth cohort study (MALED). *Lancet Glob Health*. 3(9):e564-75.
doi: 10.1016/S2214-109X(15)00151-5
117. Bauck, A., Patil, C., Mduma, et al. 2015. Nutrient sources in the diets of rural Tanzanian children 9-24 months and local pathways to nutrient adequacy. *The FASEB Journal*, 29(1 Suppl):391-2.
118. Chandyo, R, Maciel, B, Ambikapathi, R, Patil, et al. 2015. Quality of the early infant diet across the eight low and middle income countries from the malnutrition and enteric disease (MAL-ED) study. *The FASEB Journal*, 29(1 Suppl):901-28.
119. Patil CL, Turab A, Ambikapathi R, et al. 2015. Early interruption of exclusive breastfeeding: results from the eight-country MAL-ED study. *J Health Popul Nutr*. 34:10.
doi: 10.1186/s41043-015-0004-2.
120. Lang D. 2015. Opportunities to assess factors contributing to the development of the intestinal microbiota in infants living in developing countries. *Microb Ecol Health Dis*. 29;26:28316.
doi: 10.3402/mehd.v26.28316.
121. Checkley W, White AC Jr, Jaganath D, et al. 2015. A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for cryptosporidium. *Lancet Infect Dis*. 15(1):85-94.
doi.org/10.1016/S1473-3099(14)70772-8.
122. The MAL-ED Network Investigators. 2014. The Malnutrition and Enteric Disease Study (MAL-ED): understanding the consequences for child health and development. *Clin Infect Dis*. 59suppl 4: S193-S330.

Individual papers listed below:

- a) The MAL-ED Network Investigators. The MAL-ED Study: A multinational and multidisciplinary approach to understand the relationship between enteric pathogens, malnutrition, gut physiology, physical growth, cognitive development, and immune responses in infants and children up to 2 years of age in resource-poor environments
doi:10.1093/cid/ciu653
- b) Keusch GT, Denno DM, Black RE, et al. Environmental Enteric Dysfunction: pathogenesis, diagnosis, and clinical consequences
doi:10.1093/cid/ciu485
- c) Denno DM, VanBuskirk K, Nelson ZC, et al. Use of the lactulose to mannitol ratio to evaluate childhood environmental enteric dysfunction: a systematic review
doi:10.1093/cid/ciu541
- d) Richard SA, Barrett LJ, Guerrant RL, et al. Disease surveillance methods used in the 8-site MAL-ED cohort study
doi:10.1093/cid/ciu435
- e) Houpt E, Gratz J, Kosek M, et al. Microbiologic methods utilized in the MAL-ED cohort study
doi:10.1093/cid/ciu413
- f) Platts-Mills JA, McCormick BJJ, Kosek M, et al. Methods of analysis of enteropathogen infection in the MAL-ED cohort study
doi:10.1093/cid/ciu408
- g) Kosek M, Guerrant RL, Kang G, et al. Assessment of environmental enteropathy in the MAL-ED cohort study: theoretical and analytic framework
doi:10.1093/cid/ciu457
- h) Caulfield LE, Bose A, Chandyo RK, et al. Infant feeding practices, dietary adequacy, and micronutrient status measures in the MAL-ED study
doi:10.1093/cid/ciu421

- i) Richard SA, McCormick BJJ, Miller MA, et al. Modeling environmental influences on child growth in the MAL-ED cohort study: opportunities and challenges
doi:10.1093/cid/ciu436
 - j) Murray-Kolb LE, Rasmussen ZA, Scharf RJ, et al. The MAL-ED cohort study: methods and lessons learned when assessing early child development and caregiving mediators in infants and young children in 8 low- and middle-income countries
doi:10.1093/cid/ciu437
 - k) Hoest C, Seidman JC, Pan W, et al. Evaluating associations between vaccine response and malnutrition, gut function, and enteric infections in the MAL-ED cohort study: methods and challenges
doi:10.1093/cid/ciu611
 - l) Ahmed T, Mahfuz M, Islam MM, et al. The MAL-ED cohort study in Mirpur, Bangladesh
doi:10.1093/cid/ciu458
 - m) Lima AAM, Oriá RB, Soares AM, et al. Geography, population, demography, socioeconomic, anthropometry, and environmental status in the MAL-ED cohort and case-control study sites in Fortaleza, Ceará, Brazil
doi:10.1093/cid/ciu438
 - n) John SM, Thomas RJ, Kaki S, et al. Establishment of the MAL-ED birth cohort study site in Vellore, Southern India
doi:10.1093/cid/ciu390
 - o) Shrestha PS, Shrestha SK, Bodhidatta L, et al. Bhaktapur, Nepal: The MAL-ED birth cohort study in Nepal
doi:10.1093/cid/ciu459
 - p) Turab A, Soofi SB, Ahmed I, et al. Demographic, socioeconomic, and health characteristics of the MAL-ED network study site in rural Pakistan
doi:10.1093/cid/ciu391
 - q) Yori PP, Lee G, Olórtegui MP, et al. Santa Clara de Nanay: The MAL-ED cohort in Peru
doi:10.1093/cid/ciu460
 - r) Bessong PO, Nyathi E, Mahopo TC, et al. Development of the Dzimauli community in Vhembe District, Limpopo Province of South Africa, for the MAL-ED Cohort Study
doi:10.1093/cid/ciu418
 - s) Mduma ER, Gratz J, Patil C, et al. The Etiology, risk factors, and interactions of enteric infections and malnutrition and the consequences for child health and development study (MAL-ED): description of the Tanzanian site
doi:10.1093/cid/ciu439
123. Mohan VR, Sharma S, Ramanujam K, et al. 2014. Effects of elevated blood lead levels in preschool children in urban Vellore. *Indian Pediatr.* 51(8):621-5.
PMID: 25128994
124. Platts-Mills JA, Gratz J, Mduma E, et al. 2014. Association between stool enteropathogen quantity and disease in Tanzanian children using TaqMan Array Cards: A nested case-control study. *Am J Trop Med Hyg.* 90(1):133-8.
doi: 10.4269/ajtmh.13-0439
125. Lee G, Paredes Olortegui M, Rengifo Pinedo S, et al. 2014. Infant feeding practices in the Peruvian Amazon: implications for programs to improve feeding. *Rev Panam Salud Publica.* 36(3):150-7.
PMID: 25418764
126. Platts-Mills JA, Liu J, Gratz J, et al. 2014. Detection of *Campylobacter* in stool and determination of significance by culture, enzyme immunoassay, and PCR in developing countries. *J Clin Microbiol.*

- 52:1074-80.
doi: 10.1128/JCM.02935-13
127. Platts-Mills JA and Kosek M. 2014. Update on the burden of *Campylobacter* in developing countries. *Curr Opin Infect Dis.* 27(5):444-50.
doi: 10.1097/QCO.0000000000000091
 128. Liu J, Kabir F, Manneh J, et al. 2014. Development and assessment of molecular diagnostic tests for 15 enteropathogens causing childhood diarrhoea: a multicentre study. *Lancet Infect. Dis.* 14(8):716-24.
doi.org/10.1016/S1473-3099(14)70808-4
 129. Lee GO, Kosek P, Lima AA, et al. 2014. The Lactulose: Mannitol diagnostic test by HPLC and LC-MS/MS platforms: considerations for field studies of intestinal barrier function and environmental enteropathy. *J Pediatr Gastroenterol Nutr.* 59(4).
doi: 10.1097/MPG.0000000000000459
 130. Pendergast LL, Scharf RJ, Rasmussen ZA, et al. 2014. Postpartum depressive symptoms across time and place: structural invariance of the Self-Reporting Questionnaire among women from the international, multi-site MAL-ED study. *J Affect Disord.* 167:178-86.
doi.org/10.1016/j.jad.2014.05.039
 131. Psaki SR, Seidman JC, Miller M, et al. 2014. Measuring socioeconomic status in multicountry studies: results from the eight-country MAL-ED study. *PHM.* 12(8).
doi:10.1186/1478-7954-12-8
 132. McCormick, BJJ. 2013. Frequent symptomatic or asymptomatic infections may have long-term consequences on growth and cognitive development. Series of the Old Herborn University Seminar Monograph. 27. Persisting Consequences of Intestinal Infection.
 133. Peterson KM, Buss J, Easley R, et al. 2013. REG1B as a predictor of childhood stunting in Bangladesh and Peru. *Am. J. Clin. Nutr.* 97:1129–33.
doi: 10.3945/ajcn.112.048306
 134. Liu J, Gratz J, Amour C, et al. 2013. A laboratory developed Taqman array card for simultaneous detection of nineteen enteropathogens. *J. Clin. Microbiol.* 51(2):472-480.
doi: 10.1128/JCM.02658-12
 135. Kosek M, Haque R, Lima A, et al. 2013. Fecal markers of intestinal inflammation and permeability associated with the subsequent acquisition of linear growth deficits in infants. *Am. J. Trop. Med. Hyg.* 88:390-396.
doi:10.4269/ajtmh.2012.12-0549
 136. Lewis FI and McCormick BJJ. 2012. Revealing the complexity of health determinants in resource-poor settings. *Am. J. Epidemiol.* 176(11):1051-1059.
doi: 10.1093/aje/kws183
 137. Psaki S, Bhutta ZA, Ahmed T, et al. 2012. Household food access and child malnutrition: results from the eight-country MAL-ED study. *Popul Health Metr.* 10(1):24.
doi: 10.1186/1478-7954-10-24
 138. Alonso WJ and McCormick BJJ. 2012. EPIPOI: A user-friendly analytical tool for the extraction and visualization of temporal parameters from epidemiological time series. *BMC Public Health* 12:982.
doi:10.1186/1471-2458-12-982
 139. Alonso WJ, Acuna-Soto R, Giglio R, et al. 2012. Spatio-temporal patterns of diarrhoeal mortality in Mexico. *Epidemiol Infect.* 140:91-99.
doi: 10.1017/S0950268811000562
 140. McCormick BJJ, Alonso WJ, and Miller MA. 2011. An exploration of spatial patterns of seasonal diarrhoeal morbidity in Thailand. *Epidemiol Infect.* 140(7):1236-43.
doi: 10.1017/S0950268811001919

141. Lang D. 2010. The MAL-ED Project: Deciphering the relationships among normal gut flora, enteric infection and malnutrition and their association with immune response to vaccines. Old Herborn University Seminar Monograph #24. Development of Strategies to Overcome Barriers to Effective Mucosal Immunization of Infants in Developing Countries.