

70015 Mathematics for Machine Learning: Exercises

Mark van der Wilk, Yingzhen Li*
{m.vdwilk,yingzhen.li}@imperial.ac.uk

October 4, 2022

Contents

1	Notation of probabilities	1
2	Warm-up Exercises	2
2.1	Probability Theory	2
2.2	Linear Algebra	4
3	Warm-up Exercises Answers	5
3.1	Warm-up Exercises	5
3.2	Linear Algebra	7

1 Notation of probabilities

In this course we will use the notation for probabilities that is common in machine learning. The main advantage is that this notation is shorter, although it does leave certain things implicit. We include this to reduce confusion.

Consider a probability space $(\Omega, \mathcal{E}, \mathbb{P})$ with sample space Ω (all possible outcomes of a random procedure), event space \mathcal{E} (the set of all sets of outcomes that we assign a probability to), and probability function $\mathbb{P} : \mathcal{E} \rightarrow [0, 1]$ (a function that assigns a probability to an event), with a random variable $X : \Omega \rightarrow \mathbb{R}^D$.

- With $\mathbb{P}(E)$ we denote the probability of an event $E \in \mathcal{E}$, where E is a set of outcomes.
- Following the usual convention, we use the same notation when considering random variables, e.g. $\mathbb{P}(X < 2)$ is short for $\mathbb{P}(\{s \in \Omega : X(s) < 2\})$ (see §6.1 in 50008 *Probability & Statistics*).
- We usually work directly with random variables, and specify all properties using a probability mass function (pmf) or probability density function (pdf). For a specific outcome of the random variable α , we write:

$$\mathbb{P}(X = \alpha) = p_X(\alpha) \quad \text{for a pmf } p_X(\cdot), \quad (1)$$

$$\mathbb{P}(X \in [a, b]) = \int_a^b p_X(\alpha) d\alpha \quad \text{for a pdf } p_X(\cdot) \text{ with } \alpha \in \mathbb{R}, \quad (2)$$

$$\mathbb{P}(X \in A) = \int_A p_X(\alpha) d\alpha \quad \text{for a pdf } p_X(\cdot) \text{ with } \alpha \in \mathbb{R}^D. \quad (3)$$

- Sometimes we may write vectors in boldface, i.e. $\mathbf{x} \in \mathbb{R}^D$. We won't always though, so keep track of how we define variables!
- We generally denote outcomes of random variables without referring explicitly to the random variable itself. For example, when we refer to an outcome \mathbf{x} , we implicitly know there is a random variable that can take this value. We usually denote this as the capital, for example here X .

*Many thanks to teaching assistants Carles Balsells Rodas, and Alex Spies for their solutions and improvements to the document.

- Sometimes we abuse notation, and drop the random variable when denoting distributions when the argument of the function identifies it, e.g. $p(\mathbf{x}) = p_X(\mathbf{x})$.
- If we want to be explicit about the random variable that we are evaluating the density/mass of, I will write e.g. $p_{X,Y}(\mathbf{x}, \mathbf{y}) = p_{X|Y}(\mathbf{x}|\mathbf{y})p_Y(\mathbf{y})$.
- Expectations can be denoted in two ways:

$$\mathbb{E}_X[f(X)] \quad \text{to emphasise that } X \text{ is random, if it is clear what its distribution is,} \quad (4)$$

$$\mathbb{E}_{p(\mathbf{x})}[f(\mathbf{x})] \quad \text{to emphasise that we will be integrating over the distribution } p(\mathbf{x}). \quad (5)$$

In both cases this corresponds to the integral $\int p(\mathbf{x})f(\mathbf{x})d\mathbf{x}$.

- Often, densities and pmfs can be discussed in exactly the same way, if we think of the density of a discrete RV as a sum of delta functions. I.e. $p(\mathbf{x}) = \sum_o \delta(\mathbf{x} - \mathbf{x}_o)p_o$, where $\{\mathbf{x}_o\}$ is the set of discrete possible outcomes that X can take, and p_o are their corresponding probabilities. This allows us to write an expectation as an integral, regardless of whether the RV is continuous or discrete, because for discrete RVs we get:

$$\mathbb{E}_{p(\mathbf{x})}[f(\mathbf{x})] = \int p(\mathbf{x})f(\mathbf{x})d\mathbf{x} = \int \sum_o \delta(\mathbf{x} - \mathbf{x}_o)p_o f(\mathbf{x})d\mathbf{x} = \sum_o f(\mathbf{x}_o)p_o. \quad (6)$$

(A delta function has the property that $\int_A \delta(\mathbf{x})d\mathbf{x}$ is 1 if $0 \in A$, and 0 otherwise. Linearity of integrals still holds. It can often be seen as the limit of a Gaussian distribution with zero variance.)

2 Warm-up Exercises

To start, here are some exercises which test knowledge which is assumed in the course.

2.1 Probability Theory

We assume that you are familiar with probability theory up to the Computing 2nd year 50008 *Probability & Statistics* course. Here are some questions to serve as a refresher. Students who are not familiar with this background should refer to the notes of 50008 *Probability & Statistics* or relevant chapters of [Deisenroth et al., 2020]. **We recommend you look at these questions when/before the course starts.** If you need a refresher, or if you do not know the notation, refer to the 50008 *Probability & Statistics* notes, or discuss with a TA.

Question 1 (Set Theory and Probability). Using the three axioms of probability show that

- Write down the sample space of a dice. In your notation, use the set A to denote the event of a 3 or 4 occurring. What is the complement of A , denoted $\neg A$?
- For a problem about lengths, we have a sample space $\Omega = [0, 1]$. For $A = (0.3, 0.4]$, what is $\neg A$?
- $\mathbb{P}(\neg A) = 1 - \mathbb{P}(A)$
- $\mathbb{P}(\emptyset) = 0$, where \emptyset is the empty set
- $0 \leq \mathbb{P}(A) \leq 1$
- $A \subseteq B \implies \mathbb{P}(A) \leq \mathbb{P}(B)$
- Hint:* Consider the following definition. $B \setminus A = \{x \in B : x \notin A\}$
- $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- (*) if $\{A_i\}_{i=1}^\infty \subseteq \Omega$ and $A_i \subseteq A_{i+1} \forall i$ then:

$$\mathbb{P}\left(\bigcup_{i=1}^\infty A_i\right) = \lim_{i \rightarrow \infty} \mathbb{P}(A_i)$$

Hint: Use axiom 3. *: The emphasis of this course isn't on these kinds of details, even though this should be doable with 1st-year calculus.

- i. For two mutually exclusive events A, B , what is $\mathbb{P}(A \cup B)$?

See Deisenroth et al. [2020, §6.1.2] for a general overview, and §4, §§5.1-5.4 of 50008 *Probability & Statistics* for more details.

Question 2 (Independent events). **Independent events don't come up as much as independent random variables, so it's ok to just follow this answer, rather than spending lots of time on it.** When tossing two coins (where we care about the order), we have a sample space $\Omega = \{HH, HT, TH, TT\}$.

- What outcomes are contained in the event that corresponds to the the first coin being heads? We denote the event E_{1H} , and others similarly.
- If you assume that all outcomes have equal probability, show that E_{1H} and E_{2T} are independent.
- If you assume that E_{1H} and E_{2H} are independent and 0.5 each, show that all outcomes must have equal probability.

See §5.3.3 in 50008 *Probability & Statistics*.

Question 3 (Random Variables). Consider throwing two fair dice.

- What is the sample space for all outcomes that you can get from throwing two dice? We specify the probability of each outcome to be the same.
- Define two random variables A, B which map the outcome to the face value on each die respectively. Find the probability mass function for A from the probability on outcomes. The answer will work from the definition of a random variable, but you will probably intuitively get the right answer as well.
- Show that A and B are independent.
- Define the random variable $C = A + B$. Derive the probability mass function of C .

See §6 of 50008 *Probability & Statistics*.

Question 4 (Continuous Random Variables). Consider the random variable X with a probability density $p(x) = C \cdot x$ when $x \in [0, 1]$ and 0 elsewhere.

- Calculate C .
- Calculate $\mathbb{P}(0.3 \leq X \leq 0.75)$.
- Calculate $\mathbb{P}(X \in [0.3, 0.75] \cup [0.8, 0.9])$.
- Calculate $\mathbb{E}_X[X]$, $\mathbb{E}_X[X^2]$, $\mathbb{V}_X[X]$.

Check your answers by performing numerical integration, e.g. in Python.

See §6.3, §7 of 50008 *Probability & Statistics* or Deisenroth et al. [2020, §6.2.2].

Question 5 (Joint Discrete Random Variables). Consider two random variables A, C , where A is the outcome of one die, and C gives the sum of A and the sum of another die B .

- From intuition, write a table of $\mathbb{P}(C = c | A = a)$, which we use to denote the probability of C taking the value c , if we know that A has taken the value a .
- Write a table of $\mathbb{P}(C = c, A = a)$. To help you think it through, consider a tree of outcomes that can occur. This helps illustrate independence between outcomes, which helps you figure out when you can multiply probabilities.
- From the values in the table $\mathbb{P}(C = c, A = a)$ find $\mathbb{P}(2 \leq C \leq 4)$ and $\mathbb{P}(2 \leq C \leq 4, 2 \leq A \leq 4)$.

We will cover conditional probability more later, but for now just think it through.

Question 6 (Multivariate Integration). Consider two continuous random variables X, Y with joint density $p(x, y) = C \cdot (x^2 + xy)$.

- Find C .

- b. Find $\mathbb{P}(0.3 \leq X \leq 0.5)$.
- c. Find $\mathbb{P}(X < Y)$. Perform the integration twice in both orders, once integrating over x first, once by integrating over y first.
- d. **Bonus:** Convince yourself that you know how to do this for $p(x, y, z) = C \cdot (x^2 + xyz)$ as well.

Check your answers by performing numerical integration, e.g. in Python.

Question 7 (Statistics). Recall the following statistical terminology.

- a. What is a statistic?
- b. What is an estimator?
- c. What is a consistent estimator?

2.2 Linear Algebra

Question 8 (Dot product). Compute $\mathbf{x}^\top \mathbf{y}$ where $\mathbf{x} = (1, -2, 5, -1)^\top$ and $\mathbf{y} = (0, 4, -3, 7)^\top$.

Question 9 (Matrix product). Compute $\mathbf{y} = A\mathbf{x}$ as well as the ℓ_2 norm of \mathbf{x} and \mathbf{y} , where

$$A = \begin{pmatrix} -1 & 4 & 7 & 2 \\ 3 & -2 & -1 & 0 \\ 5 & 3 & 0 & -1 \end{pmatrix}, \quad \mathbf{x} = (-3, 2, 1, 3)^\top.$$

Question 10 (Basis). Which of the following set of vectors are basis for \mathbb{R}^2 ?

- 1. $\{(1, 1), (1, 0)\}$
- 2. $\{(2, 4), (3, -1)\}$
- 3. $\{(1, -1), (0, 2), (2, 1)\}$
- 4. $\{(2, -1), (-2, 1)\}$
- 5. $\{(0, 3)\}$

Question 11 (Span of vectors). Which of the following points are within the span of $\{(-1, 0, 2), (3, 1, 0)\}$?

- 1. $(0, 1, 1)$
- 2. $(1, 1, 4)$
- 3. $(2, 1, 1)$
- 4. $(-3, 4, 2)$
- 5. $(0, 0, 0)$

Question 12 (Rotation matrix in \mathbb{R}^2). What is the 2×2 matrix that rotates all the non-zero vectors in \mathbb{R}^2 by 45° counter-clockwise?

Question 13 (Linear equations). Given the following system of linear equations:

$$\begin{aligned} x + 2y &= 2 \\ 3x + 2y + 4z &= 5 \\ -2x + y - 2z &= -1 \end{aligned}$$

Answer the following questions:

- a. Writing this system in a matrix form $A\mathbf{x} = \mathbf{b}$ with $\mathbf{x} = (x, y, z)^\top$. What are A and \mathbf{b} ?
- b. Solve this system, or show that the solution does not exist.
- c. What is the rank of A ?

References

Marc Peter Deisenroth, A. Aldo Faisal, and Cheng Soon Ong (2020). *Mathematics for machine learning*. Cambridge University Press.