Detector induced assymetry in CP violation measurements

Eugenia Spedicato, Lina Maria Ortiz Parra, Jonah Blank

March 6, 2020

Comments - efficiencies

- very small errors in $D = \frac{\epsilon_+ \epsilon_-}{\epsilon_+ + \epsilon_-}$ $\rightarrow D = 0$ out of 5σ -range
- *D* is much smaller in for the *UP*-polarity
- smaller error for UP due to higher statistics
- no difference in the efficiencies between UP and DOWN within scope of the error
- in the MC: $\epsilon_{D^*}=0$ (Dst_reconstructed always 0) in our computation: $\epsilon_{D^*}=\epsilon_{\pi,s}\cdot\epsilon_{D^0}$

Comments - plots


- lacksquare structure of $\epsilon(\phi)$ probably due to rectangular detector shape
- peak in $\epsilon_{D^*}(\theta)$ within scope of error

Total


Efficiencies

Polarity	ϵ_{π}	$\epsilon_{\mathcal{K}}$	$\epsilon_{\pi,s}$	ϵ_{D^0}	ϵ_{D^*}
UP	86.65 ± 0.01	84.63 ± 0.01	76.65 ± 0.02	$\textbf{73.34} \pm \textbf{0.02}$	56.31 ± 0.02
DOWN	86.68 ± 0.01	84.67 ± 0.01	76.66 ± 0.02	73.39 ± 0.02	56.35 ± 0.02


π -efficiency




K-efficiency




soft π -efficiency



D^0 -efficiency



D*-efficiency




Charge: +

Efficiencies


Polarity	ϵ_{π}	$\epsilon_{\mathcal{K}}$	$\epsilon_{\pi,s}$	ϵ_{D^0}	ϵ_{D^*}
UP DOWN	$86.66 \pm 0.02 \\ 86.70 \pm 0.02$	$\begin{array}{c} 85.02 \pm 0.02 \\ 85.07 \pm 0.02 \end{array}$	$76.37 \pm 0.02 \\ 76.98 \pm 0.02$	$73.01 \pm 0.03 \\ 73.06 \pm 0.03$	$55.86 \pm 0.03 \\ 56.33 \pm 0.03$
UP	π	K	soft π	D^0	D*
			2 352 910 3 081 050		
DOW	N π	K	soft π	D^0	<i>D</i> *
$N_{ m reco}$			0 2374360 0 3084220		




π -efficiency




K-efficiency




soft π -efficiency



D^0 -efficiency



D*-efficiency




Charge: -

Efficiencies


Polarity	ϵ_{π}	$\epsilon_{\mathcal{K}}$	$\epsilon_{\pi,s}$	ϵ_{D^0}	ϵ_{D^*}
UP DOWN	$\begin{array}{c} 86.65 \pm 0.02 \\ 86.66 \pm 0.02 \end{array}$	$\begin{array}{c} 84.25 \pm 0.02 \\ 84.27 \pm 0.02 \end{array}$	$76.93 \pm 0.02 \\ 76.34 \pm 0.02$	$73.67 \pm 0.03 \\ 73.72 \pm 0.03$	$56.76 \pm 0.03 \\ 56.36 \pm 0.02$
UP	π	K	soft π	D^0	D*
			2 371 060 3 082 060		
DOW	N π	K	soft π	D^0	<i>D</i> *
$N_{ m reco}$	•		0 2356980 0 3087370		




π -efficiency




K-efficiency




soft π -efficiency



D^0 -efficiency



D*-efficiency



Deviation

Table: The deviation
$$\frac{\epsilon_+ - \epsilon_-}{\epsilon_+ + \epsilon_-}/10^{-3}$$

Polarity	π	K	$soft\pi$	D^0	D^*
UP DOWN	$0.1 \pm 0.2 \\ 0.2 \pm 0.2$		-3.7 ± 0.2 4.2 ± 0.2		


Table: The deviation
$$\frac{N_+ - N_-}{N_+ + N_-}/10^{-3}$$

Polarity	π	K	$soft\pi$	D^0	<i>D</i> *
UP -	-0.1 ± 0.4	4.7 ± 0.4	-3.8 ± 0.5	-4.7 ± 0.5	-8.2 ± 0.5
DOWN -	-0.3 ± 0.4	5.2 ± 0.4	3.7 ± 0.5	-5.0 ± 0.5	-0.8 ± 0.5


- calculation via N_{reco} doesn't work, $N_{\text{tot},+/-}$ different \rightarrow no normalization
- interesting: $D_{soft \pi} \& D_{D^0}$ cancel partially in DOWN, but add up in UP




Comparison of different charges with UP polarity - D^*




Comparison of different charges with DOWN polarity - D^*





Comparison - $\overline{D^*p_T}$







Comparison - $D^*\phi$







Comparison - $D^*\theta$






Comparison - $D^*\eta$







D^* deviation dependencies - UP polarity




D* deviation dependencies - DOWN polarity



D^* deviation - ϕ





- left UP, right DOWN
- clear dependency in ϕ , inverted $UP \leftrightarrow DOWN$
- doesn't seem to have dependency on other topological variables
 - \rightarrow form of the detector is biggest source of induced CPV

$\overline{D^*}$ deviation UP+DOWN

