

Eva Conquet, Arpat Ozgul, Daniel T. Blumstein, Kenneth B. Armitage, Madan K. Oli, Julien G. A. Martin, Tim H. Clutton-Brock, Maria Paniw. 2021.
Demographic consequences of changes in environmental periodicity. *Ecology*.

Data and R scripts to reproduce the analysis of the effect of changes in environmental periodicity on the dewy pine (*Drosophyllum lusitanicum*) population.

Author(s)

Eva Conquet
Department of Evolutionary Biology and Environmental Studies, University of Zurich
Winterthurerstrasse 190, Zurich 8057, Switzerland
eva.conquet@uzh.ch

Arpat Ozgul
Department of Evolutionary Biology and Environmental Studies, University of Zurich
Winterthurerstrasse 190, Zurich 8057, Switzerland
arpat.ozgul@ieu.uzh.ch

Daniel T. Blumstein
Department of Ecology and Evolutionary Biology, University of California Los Angeles
Los Angeles, CA 90095, USA
The Rocky Mountain Biological Laboratory
Crested Butte, CO 81224, USA
marmots@ucla.edu

Kenneth B. Armitage
Department of Ecology and Evolutionary Biology, The University of Kansas
Lawrence, KS 66045-7534, USA
marmots@ku.edu

Madan K. Oli
Department of Wildlife Ecology and Conservation, University of Florida
Gainesville, FL 32611, USA
olim@ufl.edu

Julien G. A. Martin
Department of Biology, University of Ottawa
Ottawa K1N 9A7, Canada
School of Biological Sciences, University of Aberdeen

Aberdeen AB24 2TZ, UK
julien.martin@uottawa.ca

Tim H. Clutton-Brock
Department of Zoology, University of Cambridge
Downing Street, Cambridge CB2 3EJ, UK
Kalahari Research Trust
Kuruman River Reserve, Northern Cape, South Africa
Mammal Research Institute, University of Pretoria
Pretoria, South Africa
thcb@cam.ac.uk

Maria Paniw
Department of Evolutionary Biology and Environmental Studies, University of Zurich
Zurich 8057, Switzerland
Department of Conservation and Global Change, Doñana Biological Station (EBD-CSIC)
Seville 41092, Spain
m.paniw@gmail.com

File list

```
DewyPinesTSF1_to_3_Data.csv  
DewyPinesTSF3plus_Data.csv  
01_DewyPines_VitalRatesEstimation_TSF1_to_3.R  
02_DewyPines_VitalRatesEstimation_TSF3plus.R  
03_DewyPines_Constants.R  
04_DewyPines_MPMs.R  
05_DewyPines_Simulations.R  
06_DewyPines_Results.R  
07_DewyPines_Sensitivities_Megamatrix_PostFireStates.R
```

Description

DewyPinesTSF1_to_3_Data.csv – This dataset contains individual capture-recapture data collected on a population of dewy pines (*Drosophyllum lusitanicum*) at three sites (a human-disturbed site C and two naturally fire-disturbed sites E and F) between 2011 and 2019 in southern Spain. For each individual at each capture, the data contains information on the site, transect, and sub-quadrat where the individual was found (site, transect, and subQuadrat), the year and time since fire (TSF; one, two, and three corresponding to the first, second, and third and above years after a fire occurred, respectively) and the stage (stage; SD = seedling, J = juvenile, SR = small reproductive individual, LR = large reproductive individual). In addition, the data contains information on whether an individual survived to the next year (surv; 1 if it survived, 0 if not), whether a juvenile transitioned to large reproductive (transitionJ; 1 if it transitioned to LR, 0 if not),

whether a small and large reproductive individual grew or shrank, respectively (transitionSR and transitionLR; 1 if it transitioned to the other reproductive stage, 0 if not), whether an individual flowered (fl; 1 if it flowered, 0 if not), and the number of flowering stalks and flowers per stalk (fs and fps). The dataset also contains values for the above-ground density (density; in individuals/m²) and its squared value (density2). This dataset is used to estimate vital rates in the dewy pine population in the deterministic post-fire habitat states TSF₁ (time since fire), TSF₂, and TSF₃, and assess the effect of changes in vital-rate periodicity on the dewy pine population dynamics.

DewyPinesTSF3plus_Data.csv – This dataset contains individual capture-recapture data collected on a population of dewy pines (*Drosophyllum lusitanicum*) at various little or highly grazed sites between 2011 and 2019 in southern Spain. For each individual at each capture, the data contains information on the site, transect and sub-quadrat where the individual was found (site, transect, and subQuadrat) and the type of grazing at that site (LS; LG = little grazed, HG = highly grazed), the year and time since fire (TSF; one, two, and three corresponding to the first, second, and third and above years after a fire occurred, respectively), and the stage (stage; SD = seedling, J = juvenile, SR = small reproductive individual, LR = large reproductive individual). In addition, the data contains information on whether an individual survived to the next year (surv; 1 if it survived, 0 if not), whether a juvenile transitioned to large reproductive (transitionJ; 1 if it transitioned to LR, 0 if not), whether a small and large reproductive individual grew or shrank, respectively (transitionSR and transitionLR; 1 if it transitioned to the other reproductive stage, 0 if not), whether an individual flowered (fl; 1 if it flowered, 0 if not), and the number of flowering stalks and flowers per stalk (fs and fps). The dataset also contains values for the above-ground density (density; in individuals/m²) and its squared value (density2). This dataset is used to estimate vital rates in the dewy pine population in the last stochastic post-fire habitat state TSF_{>3} (time since fire), and assess the effect of changes in vital-rate periodicity on the dewy pine population dynamics.

01_DewyPines_VitalRatesEstimation_TSF1_to_3.R – This script uses the individual capture-recapture data on the dewy pine population for TSF₁, TSF₂, and TSF₃ (DewyPinesTSF1_to_3_Data.csv) to estimate survival, transition, and reproductive rates (flowering probability, number of stalks, and number flower per stalk). We estimated vital rates using binomial (for survival, transition, and flowering probability rates) and Poisson generalized linear models (GLMs; for the number of flowering stalks and the number of flowers per stalk). We included time since fire, density, and squared density as fixed effects. We selected the best fixed effect using the AICc. We checked for over- and under-dispersion in the number of flowering stalks and the number of flowers per stalk models and fitted a quasi-Poisson model to account for it. At the end of the script, vital rates are stored in a data frame and all models are saved as .RData objects.

02_DewyPines_VitalRatesEstimation_TSF3plus.R – This script uses the individual capture-recapture data on the dewy pine population for TSF_{>3} (DewyPinesTSF3plus_Data.csv) to estimate survival, transition, and reproductive rates (flowering probability, number of stalks, and number flower per stalk). We estimated vital rates using binomial (for survival, transition, and flowering probability rates) and

Poisson generalized linear mixed models (GLMMs; for the number of flowering stalks and the number of flowers per stalk). We included time since fire, density, and squared density as fixed effects, and year as a random effect on the intercept. We selected the best random effect by looking at the part of the variance explained by a year random effect compared to no random effect, using the *r.squaredGLMM* function of the *MuMIn* R package. We selected the best fixed effect using the AICc. We checked for over- and under-dispersion in the number of flowering stalks and the number of flowers per stalk models and fitted a quasi-Poisson model to account for it. At the end of the script, vital rates are stored in a data frame and all models are saved as `.RData` objects.

`03_DewyPines_Constants.R` – This script provides the values for constant vital rates for the seed bank dynamics in the human-disturbed (*seed.bankSiteC*) and naturally fire-disturbed sites (*seed.bankSitesEF*), the number of seeds per flower (*seeds.per.flower*), and the transition probability to seedling or juvenile from the seed bank in TSF₀ (*sb.transition.TSF0*). The constants are stored as `.csv` files or `.RData` objects.

`04_DewyPines_MPMs.R` – This script provides the functions to build periodic matrix population models (MPMs) for the dewy pine population, using vital rates estimated in `01_DewyPines_VitalRatesEstimation_TSF1_to_3.R` and `02_DewyPines_VitalRatesEstimation_TSF3plus.R`. The script contains one function for each post-fire habitat state (TSF₀ to TSF₃) for the human-disturbed site (site C) and the naturally fire-disturbed sites (sites E and F). Depending on the TSF and site, the functions can take the density and the year as arguments and return the corresponding MPM.

`05_DewyPines_Simulations.R` – This script provides the *stoch.sim.droso* function to perform simulations under a control (no changes in vital-rate periodicity in the naturally fire-disturbed population), and a perturbed scenario where vital rates from the human-disturbed population are introduced in one of the five post-fire habitat states. In the fifth post-fire habitat state (TSF₃), various proportions of the number of years spent in that state can be perturbed. The function takes the number of simulations (*nsimul*; default is 500), the number of years in each simulation (*nyears*; default is 100), the initial population vector (*n0*), the fire-return probability (*fire.prob*; default is 0.033), the density scenario (*density*; TRUE if density dependence should be taken into account, FALSE if simulations should be performed at average population density), and the proportions of years spent in TSF₃ to be perturbed (*state5.perturbed.years.prop*). For each simulation, a sequence of post-fire habitats (*states*) is created using a Markov chain *envS* in which transitions from the first to the fifth state are deterministic and the transition between the fifth and the first state depends on the fire-return probability. The function returns a list of annual growth rates for each simulation, a vector containing a 1 if the simulation led to population quasi-extinction, a list of yearly population densities for each simulation, a list of post-fire habitat sequences, and a list of population vectors. The rest of the code performs all the simulations used in the manuscript and saves the results as a `.RData` object.

`06_DewyPines_Results.R` – This script processes the results of the vital-rate periodicity perturbation simulations for the dewy pine population. It computes and plots the

mean stochastic log lambda, variance across 100 log lambda, and quasi-extinction probability in each scenario across 500 simulations. The results are saved as a `.csv` file.

`07_DewyPines_Elasticities_Megamatrix_PostFireStates.R` – This script computes and plots the elasticities of the population growth rate to each post-fire state using the megamatrix approach. We parameterized a 25 X 25 block-diagonal matrix **A** with each of five post-fire state-specific matrix population models on the diagonal and zeros elsewhere. We then built the megamatrix $\mathbf{M} = (\mathbf{C} \otimes \mathbf{I}) \times \mathbf{A}$, with **C** a matrix containing probabilities of transition between post-fire habitat states and **I** a 5 X 5 identity matrix. We then computed the elasticity of the population growth rate to each post-fire state using the *elasticity* function of the *popbio* package. The results are saved as a `.csv` file.