
COVID-19:



Everything You Need to Know About the booster shot



54.8%

Of the worlds population have already recieved at least one Covid Vaccine

All countries need to review their strategies now.

All nations are called to take the fight against this virus seriously. They should communicate well with their populations, and pull all the stops when it comes to looking for ways to mitigate the spr

Dr. Michael J. Ryan, WHO Informal Advisory Group, March 2020



Who can get a booster?

When can one get a booster?

Which boosters are there?

Anyone above the Age of 18



< 6months after 1st shot

Pfizer, J&J ,Moderna,Astrazeneca





The "panic-then-forget" cycle must be stopped. Instead, a "whole-of-society" approach must be adopted. It is imperative that the the International Health Regulations (IHR) be implemented in countries of all income levels.

Every country should do everything to follow through on prioritizing, funding, and implementing preparedness interventions.

COVID-19:







Which booster should i take?

There is ideally no specific vaccine meant for take, however, the World Health Organization has some approved vaccines that are recommended for take internationaly, these are the vaccines that are appoved by the local government and can be accesed by citizens of these specifc countries. In terms of health risk analysis, there have been reported cased of blood clots after vaccination, however, these cases have only happened in about 1 per 10 milion and research around this topic is still ongoing. The WHO however urges persons to access these vaccines and take them as soon as they can because they have been tested and measured for quality administration Sasidharan et al (2020)



REFERENCES

https://www.who.int/emergencies/diseases/novel-coronavirus-2019/media-resources/of https://ourworldindata.org/covid-vaccinations

https://ourworldindata.org/covid-vaccinations

Sasidharan, S., Singh, D. H., Vijay, S., & Manalikuzhiyil, B. (2020). COVID-19: Pan (info) demic. Turkish Journal of Anaesthesiology and Reanimation, 48(6), 438.-