

# Quantum Mechanics A (Physics 212A) Fall 2018

## Worksheet 4 – Solutions

### Problems

#### 1. Interferometry

We can consider the path taken of a photon as (approximately) a two-state quantum system spanned by  $|u\rangle$ ,  $|d\rangle$  for whether it went up or down respectively.

Consider the following two interferometers:




Figure 1: A balanced Mach-Zender interferometer (i) and another with the final beam splitter removed (ii)

The elements BS are beam splitters which implement the Hadamard gate on the incoming beam:  $H = \frac{1}{\sqrt{2}}[(|u\rangle + |d\rangle)\langle u| + (|u\rangle - |d\rangle)\langle d|]$

The elements M are mirrors which transform  $|\text{In}\rangle \rightarrow |\text{Out}\rangle = -|\text{In}\rangle$

The element PS is a phase shifter which transform  $|\text{In}\rangle \rightarrow |\text{Out}\rangle = e^{i\phi}|\text{In}\rangle$

The element D is a detector which measures photons going in.

For each device, determine the probability for detecting a photon as a function of  $\phi$ .

Let's do setup (ii) first.

After the first beam splitter the state is  $|1\rangle = \frac{1}{\sqrt{2}}(|u\rangle + |d\rangle)$

The pair of mirrors introduces an overall minus sign  $|2\rangle = -|1\rangle$  which is irrelevant

The phase shifter then acts on the up travelling beam  $|3\rangle = \frac{1}{\sqrt{2}}(e^{i\phi}|u\rangle + |d\rangle)$

But then the only part of the beam entering the detector D is the up component. Therefore  $P(\phi) = |\langle u|3\rangle|^2 = \frac{1}{2}$ . Pretty boring.

For the true MZ interferometer (i) we apply the Hadamard:

$$|4\rangle = H|3\rangle = \frac{1}{2}(e^{i\phi} + 1)|u\rangle + \frac{1}{2}(e^{i\phi} - 1)|d\rangle$$

Then again it's measuring 'up' component so  $P(\phi) = |\langle u|4\rangle|^2 = \frac{1}{4}|e^{i\phi} + 1|^2 = \frac{1}{2}(1 + \cos \phi)$

## 2. Time-energy uncertainty

Consider an observable  $\mathbf{A}$  which has no explicit dependence on time. Define the variance of  $\mathbf{A}$  in a state  $\psi$  to be

$$\Delta A \equiv \sqrt{\langle (\mathbf{A} - \langle \mathbf{A} \rangle_\psi)^2 \rangle_\psi}$$

as usual, with  $\langle \mathbf{A} \rangle_\psi \equiv \langle \psi | \mathbf{A} | \psi \rangle$ . Let

$$\Delta T \equiv \frac{\Delta A}{|\partial_t \langle \mathbf{A} \rangle_\psi|}.$$

This is a measure of the time required for  $\langle \mathbf{A} \rangle$  to change significantly, *i.e.* by an amount comparable to its variance  $\Delta A$ . Show that

$$\Delta E \Delta T \geq \frac{\hbar}{2},$$

where  $\Delta E$  is the variance of  $\mathbf{H}$ , the Hamiltonian.

Let  $H_0 \equiv \mathbf{H} - \langle \mathbf{H} \rangle_\psi \equiv \mathbf{H} - E$  and  $A_0 \equiv \mathbf{A} - \langle \mathbf{A} \rangle_\psi$ . Apply Cauchy-Schwarz to  $|\alpha\rangle \equiv A_0|\psi\rangle$  and  $|\beta\rangle \equiv H_0|\psi\rangle$ . This gives

$$(\Delta A)^2 (\Delta E)^2 \geq \frac{1}{4} |\langle \psi | [\mathbf{A}, \mathbf{H}] | \psi \rangle|^2 = \frac{1}{4} |\mathbf{i}\hbar \partial_t \langle \mathbf{A} \rangle_\psi|^2.$$

where in the last step we've used the Heisenberg equation of motion.

## 3. Entropy and thermodynamics

Consider a quantum system with hamiltonian  $\mathbf{H}$  and Hilbert space  $\mathcal{H} = \text{span}\{|n\rangle, n = 0, 1, 2, \dots\}$ . Its behavior at finite temperature can be described using the *thermal density matrix*

$$\rho_\beta \equiv \frac{1}{Z} e^{-\beta \mathbf{H}}$$

where  $\beta \equiv \frac{1}{T}$  specifies the temperature and  $Z$  is a normalization factor. (We can think about this as the reduced density matrix resulting from coupling the system to a heat bath and tracing out the Hilbert space of the (inaccessible) heat bath.)

- (a) Find a formal expression for  $Z$  by demanding that  $\rho_\beta$  is normalized appropriately. ( $Z$  is called the *partition function*.)

$$\text{Recall that } \text{Tr} [\rho_\beta] = 1 = \frac{1}{Z} \text{Tr} [e^{-\beta \mathbf{H}}] \rightarrow Z = \text{Tr} [e^{-\beta \mathbf{H}}] = \sum_i e^{-\beta E_i}$$

- (b) Show that the von Neumann entropy of  $\rho_\beta$  can be written as

$$S_\beta = E/T + \log Z \tag{1}$$

where  $E \equiv \langle \hat{H} \rangle_\rho = \text{Tr} \rho \hat{H}$  is the expectation value for the energy. The expression above for  $S_\beta$  is the thermal entropy.

Recall the definition of the von Neumann entropy:  $S[\rho_\beta] = -\text{Tr} [\rho_\beta \log \rho_\beta]$

$$\log \rho_\beta = \log \frac{e^{-\beta \mathbf{H}}}{Z} = \log e^{-\beta \mathbf{H}} - \log Z = -(\beta \mathbf{H} + \log Z)$$

$$\rightarrow S = \text{Tr} \left[ \frac{e^{-\beta \mathbf{H}}}{Z} (\beta \mathbf{H} + \log Z) \right] = \text{Tr} [\rho_\beta \beta \mathbf{H}] + \text{Tr} [\rho_\beta \log Z] = \beta \langle \mathbf{H} \rangle + \log Z \text{Tr} [\rho_\beta]$$

Therefore  $S = \frac{E}{T} + \log Z$  by  $\text{Tr} [\rho_\beta] = 1$  and  $\beta = \frac{1}{T}$

(c) Evaluate  $Z$  and  $E$  for the case where the system is a simple harmonic oscillator

$$\mathbf{H} = \hbar\omega \left( \mathbf{n} + \frac{1}{2} \right)$$

with  $\mathbf{n}|n\rangle = n|n\rangle$ .

$$Z = \sum_n e^{-\beta E_n} = \sum_i e^{-\omega\beta(n+\frac{1}{2})} = e^{-\frac{\omega\beta}{2}} \sum_{n=0}^{\infty} e^{-\omega\beta n}$$

This sum is a geometric series:  $\sum_n x^{-n} = \frac{x}{x-1}$  which implies:  $Z = e^{-\frac{\beta\omega}{2}} \frac{e^{\beta\omega}}{e^{\beta\omega}-1}$

One way to evaluate  $E$  is to put our statistical mechanics knowledge to work:

$$\langle E \rangle = -\frac{\partial \log Z}{\partial \beta} = -\frac{1}{Z} \frac{\partial Z}{\partial \beta} = \frac{\omega}{2} \left( \frac{1+e^{\beta\omega}}{e^{\beta\omega}-1} \right)$$