

A tight-binding model for p electrons with spin-orbit interaction

Anikin Evgeny, 128

August 16, 2016

In the following paper we will consider a tight-binding model for p zone with spin-orbit interaction.

Before we start, let us remind the Clebsch-Gordan coefficients for the case $l = 1$, $s = \frac{1}{2}$. Let $a_{j,m}$ annihilate the state with angular momentum j and the projection m ($j \in \{3/2, 1/2\}$), and $b_{m,s}$ — the state with orbital momentum projection m ($j = 1$) and spin projection s .

$$\begin{aligned}
 a_{\frac{3}{2}, \frac{3}{2}} &= b_{1, \frac{1}{2}} \\
 a_{\frac{3}{2}, \frac{1}{2}} &= \sqrt{\frac{1}{3}} b_{1, -\frac{1}{2}} + \sqrt{\frac{2}{3}} b_{0, \frac{1}{2}} \\
 a_{\frac{3}{2}, -\frac{1}{2}} &= \sqrt{\frac{2}{3}} b_{0, -\frac{1}{2}} + \sqrt{\frac{1}{3}} b_{-1, \frac{1}{2}} \\
 a_{\frac{3}{2}, -\frac{3}{2}} &= b_{-1, -\frac{1}{2}} \\
 a_{\frac{1}{2}, \frac{1}{2}} &= \sqrt{\frac{2}{3}} b_{1, -\frac{1}{2}} - \sqrt{\frac{1}{3}} b_{0, \frac{1}{2}} \\
 a_{\frac{1}{2}, -\frac{1}{2}} &= -\sqrt{\frac{1}{3}} b_{0, -\frac{1}{2}} + \sqrt{\frac{2}{3}} b_{-1, \frac{1}{2}}
 \end{aligned} \tag{1}$$

After expressing the b operators using p_x , p_y and p_z , we immediately obtain

$$\begin{aligned}
 a_{\frac{3}{2}, \frac{3}{2}} &= \sqrt{\frac{1}{2}} \left(p_{x, \frac{1}{2}} - i p_{y, \frac{1}{2}} \right) \\
 a_{\frac{3}{2}, \frac{1}{2}} &= \sqrt{\frac{1}{6}} \left(p_{x, -\frac{1}{2}} - i p_{y, -\frac{1}{2}} \right) + \sqrt{\frac{2}{3}} p_{z, \frac{1}{2}} \\
 a_{\frac{3}{2}, -\frac{1}{2}} &= \sqrt{\frac{2}{3}} p_{z, -\frac{1}{2}} + \sqrt{\frac{1}{6}} \left(p_{x, \frac{1}{2}} + i p_{y, \frac{1}{2}} \right) \\
 a_{\frac{3}{2}, -\frac{3}{2}} &= \sqrt{\frac{1}{2}} \left(p_{x, -\frac{1}{2}} + i p_{y, -\frac{1}{2}} \right)
 \end{aligned} \tag{2}$$

$$\begin{aligned}
a_{\frac{1}{2}, \frac{1}{2}} &= \sqrt{\frac{1}{3}} \left(p_{x, -\frac{1}{2}} - ip_{y, -\frac{1}{2}} \right) - \sqrt{\frac{1}{3}} p_{z, \frac{1}{2}} \\
a_{\frac{1}{2}, -\frac{1}{2}} &= -\sqrt{\frac{1}{3}} p_{z, -\frac{1}{2}} + \sqrt{\frac{1}{3}} \left(p_{x, \frac{1}{2}} + ip_{y, \frac{1}{2}} \right)
\end{aligned} \tag{3}$$

As (2), (3) define a unitary transformation, p can be easily expressed via a .
The Hamiltonian is

$$\begin{aligned}
H_{\text{full}} &= -\Delta E_{SO} a_{\frac{1}{2}, -\frac{1}{2}}^\dagger a_{\frac{1}{2}, -\frac{1}{2}} + \\
&\quad + 2(t_{\parallel} \cos p_x + t_{\perp} \cos p_y) p_{x, \frac{1}{2}}^\dagger p_{x, \frac{1}{2}} + \\
&\quad + 2(t_{\perp} \cos p_x + t_{\parallel} \cos p_y) p_{y, \frac{1}{2}}^\dagger p_{y, \frac{1}{2}} + \\
&\quad + 2t_3(\cos p_x + \cos p_y) p_{z, -\frac{1}{2}}^\dagger p_{z, -\frac{1}{2}} \tag{4}
\end{aligned}$$

It contains the atomic spin-orbit part and the part which depends on the interaction between neighbour atoms.

After simple calculation we obtain the Hamiltonian matrix in the basis of p operators:

$$\begin{aligned}
H &= -\frac{E_{SO}}{3} \begin{pmatrix} 1 & i & -1 \\ -i & 1 & i \\ -1 & -i & 1 \end{pmatrix} + \\
&\quad + \begin{pmatrix} t_{\parallel} \cos p_x + t_{\perp} \cos p_y & 0 & 0 \\ 0 & t_{\perp} \cos p_x + t_{\parallel} \cos p_y & 0 \\ 0 & 0 & t_3(\cos p_x + \cos p_y) \end{pmatrix} \tag{5}
\end{aligned}$$

The energy levels for the case $\Delta E_{SO} = 1$, $t_{\parallel} = 0.3$, $t_3 = t_{\perp} = 0.15$ are shown on the figure.




Figure 1: Energy levels




Figure 2: The difference of two upper energy levels

In some cases it would be (maybe) more convenient to write the Hamiltonian in the basis of a operators. For solving that problem, the following observation would be useful.

Let $|A\rangle$ and $|B\rangle$ be the spinless states with angular momentum projections m and m' . Let $|A\rangle$ also be localised in $(0,0)$, and $|B(\phi)\rangle$ — in $(r \cos \phi, r \sin \phi)$. Then the following equation holds:

$$\langle A|B(\phi)\rangle = e^{-i(m-m')\phi} \langle A|B(0)\rangle \quad (6)$$

That allows to write any tight-binding Hamiltonian directly in terms of a states.

First Let the angle be zero, and let the hopping integrals for the b operators be described by matrix

$$T_b = \begin{pmatrix} t_1 & 0 & t_2 \\ 0 & t_3 & 0 \\ t_2 & 0 & t_1 \end{pmatrix} \quad (7)$$

The t_i can be expressed via t_\perp, t_\parallel .

$$\begin{aligned} t_1 &= \frac{1}{2} (t_\parallel + t_\perp) \\ t_2 &= \frac{1}{2} (t_\parallel - t_\perp) \end{aligned} \quad (8)$$

Then for the states described by full momentum (a states) we obtain

$$T_x = \begin{pmatrix} t_1 & \frac{1}{\sqrt{3}}t_2 & \sqrt{\frac{2}{3}}t_2 \\ \frac{1}{\sqrt{3}}t_2 & \frac{1}{3}(t_1 + 2t_3) & \frac{\sqrt{2}}{3}(t_1 - t_3) \\ \sqrt{\frac{2}{3}}t_2 & \frac{\sqrt{2}}{3}(t_1 - t_3) & \frac{1}{3}(2t_1 + t_3) \end{pmatrix} \quad (9)$$

For the arbitrary angle, as follows from 6,

$$\begin{aligned}
T_\phi &= \text{diag}(e^{-\frac{3i\phi}{2}}, e^{\frac{i\phi}{2}}, e^{\frac{i\phi}{2}}) \times T_x \times \text{diag}(e^{\frac{3i\phi}{2}}, e^{-\frac{i\phi}{2}}, e^{-\frac{i\phi}{2}}) = \\
&= \begin{pmatrix} t_1 & \frac{1}{\sqrt{3}}t_2e^{-2i\phi} & \sqrt{\frac{2}{3}}t_2e^{-2i\phi} \\ \frac{1}{\sqrt{3}}t_2e^{2i\phi} & \frac{1}{3}(t_1 + 2t_3) & \frac{\sqrt{2}}{3}(t_1 - t_3) \\ \sqrt{\frac{2}{3}}t_2e^{2i\phi} & \frac{\sqrt{2}}{3}(t_1 - t_3) & \frac{1}{3}(2t_1 + t_3) \end{pmatrix} \quad (10)
\end{aligned}$$

The Hamiltonian is now

$$\begin{aligned}
H &= \text{diag}(0, 0, -\Delta E_{SO}) + 2 \cos p_x T_0 + 2 \cos p_y T_{\frac{\pi}{2}} + \\
&\quad + 2 \cos(p_x + p_y) \tilde{T}_{\frac{\pi}{4}} + 2 \cos(p_x - p_y) \tilde{T}_{-\frac{\pi}{4}} + \dots \quad (11)
\end{aligned}$$

We will treat the first term as the unperturbed system and all other as a perturbation. The unperturbed Hamiltonian has a pair of degenerate levels. The non-trivial topology can only exist due to "entanglement" of these levels. As we treat the T matrices as a perturbation, we will find the eigenfunctions and eigenvalues of the restricted perturbation matrix V .

$$\begin{aligned}
V &= 2 \begin{pmatrix} t_1(\cos p_x + \cos p_y) + 2\tilde{t}_1 \cos p_x \cos p_y & \frac{1}{\sqrt{3}}t_2(\cos p_x - \cos p_y) + \frac{2i}{\sqrt{3}}\tilde{t}_2 \sin p_x \sin p_y \\ \frac{1}{\sqrt{3}}t_2(\cos p_x - \cos p_y) + \frac{2i}{\sqrt{3}}\tilde{t}_2 \sin p_x \sin p_y & \frac{1}{3}(t_1 + 2t_3)(\cos p_x + \cos p_y) + \frac{2}{3}(t_1 + 2\tilde{t}_3) \cos p_x \cos p_y \end{pmatrix} = \\
&= \begin{pmatrix} a & b \\ b^* & c \end{pmatrix} \quad (12)
\end{aligned}$$