

# Predicting Carbon Emission at Fine Time Granularity Using Autoformer

Shuyi Wei<sup>1</sup> (✉), Xin You<sup>2</sup>, and Yaonan Jiang<sup>1</sup>

<sup>1</sup> College of Computer Science & Technology, Hangzhou Dianzi University,  
Hangzhou, China

[weis16@uci.edu](mailto:weis16@uci.edu); [21052115@hdu.edu.cn](mailto:21052115@hdu.edu.cn)

<sup>2</sup> Zhuoyue Honors College, Hangzhou Dianzi University, Hangzhou, China  
[21051606@hdu.edu.cn](mailto:21051606@hdu.edu.cn)

**Abstract.** Carbon dioxide (CO<sub>2</sub>) is a major contributor to global warming. Accurate and high-resolution prediction of CO<sub>2</sub> emissions is critical to achieving carbon neutrality around the world. Previous methods used traditional statistical models or machine learning models, which can only predict the annual CO<sub>2</sub> emissions. With the development of deep learning models, such as GRU, they have been used for prediction. However, there is a lack of predictions for fine time granularity for CO<sub>2</sub> emissions. Thus, in this paper, we propose a carbon emission prediction model at fine time granularity. Autoformer is used to improve the prediction accuracy. A carbon emission dataset that contains 3 years of China's over 300 cities on a daily basis is used for prediction. The results show that compared with other methods, our model can achieve the highest prediction accuracy. Our model provides high-quality, fine-grained CO<sub>2</sub> emissions data to support global emission monitoring across various urban content.

**Keywords:** Carbon dioxide prediction · Autoformer · Fine time granularity

## 1 Introduction

### 1.1 Background

Human activities, particularly the burning of fossil fuels for energy, deforestation, and industrial processes, release large amounts of CO<sub>2</sub> into the atmosphere [1]. As one of the major greenhouse gases, CO<sub>2</sub> contributes significantly to global warming, resulting in profound effects on the human environment and poses a substantial threat to the long-term development of the world economy [2, 3]. Reducing greenhouse gas emissions and developing a low-carbon economy have become the consensus of all countries in the world [4].

The current CO<sub>2</sub> emission prediction methods can be divided into three types. They are statistical models, traditional machine learning methods, and deep learning models. The statistical models are not suitable for predicting sequences of CO<sub>2</sub> emissions with complex nonlinear relationships and small sample adaptation.

Traditional machine learning methods require feature selection to establish CO<sub>2</sub> prediction models. Feature selection relies on a significant amount of socioeconomic and energy-related statistical data. The challenge arises when needed data are unavailable, and the feature selection is unpredictable, making it difficult to make accurate predictions [5–8].

With the development of deep learning, models based on Recurrent Neural Networks (RNNs) have been employed for predicting CO<sub>2</sub> emissions. Despite their effectiveness in time-series prediction, RNNs architectures face limitations in capturing long-range dependencies and exhibiting computational inefficiency due to their sequential execution mechanism [9–11].

However, previous studies usually forecasted the emissions on an annual basis based on the statistical yearbook data at the national or province level, which can only provide a rough overview of the changes in carbon emissions. Hence, detailed and spatially explicit estimates of emissions are urgently needed.

By incorporating multi-source data such as electricity generation, regional energy consumption statistics, gridded population density, night lights, urban form, and GDP data, researchers have constructed daily updated CO<sub>2</sub> emission datasets with higher spatial and temporal resolution [12–14]. These datasets offer the opportunity for a more fine-grained spatiotemporal prediction of CO<sub>2</sub> emissions.

## 1.2 Main Contributions

The contributions of this paper mainly have three aspects. First, we conduct a framework containing each step of our work. Among them, in contrast to previous studies, we use a high-quality city-level CO<sub>2</sub> emission dataset on a daily basis around China, which supports our goal of making fine-grained CO<sub>2</sub> emission prediction. Also, a new methodology is proposed to predict future CO<sub>2</sub> emission.

Second, we bring Autoformer into our model with added time features to make more accurate results. This new model has a better performance than others in terms of the value of evaluation metrics, and also the results shown in plots.

Third, the time series is added to selected deep learning models in order to yield an accurate result with temporal feature. Also, the performance of this method is demonstrated by comparing with other models about the metrics RMSE and MAE, including traditional ARIMA methods, two variants of RNN-based models (RNN and GRU), and two variants of Transformer based models (Transformer and Informer).

The remainder of this paper is organized as follows: Sect. 2 provides a literature review on CO<sub>2</sub> emission datasets and prediction methods. Section 3 introduces our work ranging from data collection to methodologies we used in detail. In Sect. 4, the result is accentuated by visualizing and comparing models' performance. The conclusions and future research are summarized in Sect. 5.

## 2 Literature Review

This section is divided into two parts: CO<sub>2</sub> emission datasets and prediction models. Each part introduces the main results up to now, which are the foundation of our research.

### 2.1 CO<sub>2</sub> Emission Datasets

In this part, abundant datasets about CO<sub>2</sub> emissions are searched. The following describes the datasets from coarse-grained to fine-grained with their incongruity with this topic.

The most famous environmental research institution is called World Resources Institute (WRI). In its database, many kinds of datasets about CO<sub>2</sub> emission were found. For instance, ‘Global Emissions of CO<sub>2</sub> From Fossil Fuels: 1900–2004’ dataset has the long-range data for many years but lack of more detailed geographical features for it only at national level [15].

Also, some data from international organization are discovered. EEA Greenhouse gas data reported by European Union recently shows the greenhouse gas emission including CO<sub>2</sub> in all nations of European Union. This dataset, however, does not have the detailed time statistics based on month [16].

Many datasets have the common flaw, especially in detailed time and space description. In order to look for fine-grained data in terms of geography, it is beneficial to visit national institution.

U.S. Energy Information Administration (EIA) presented ‘CO<sub>2</sub> emission aggregates by fuel, state, and sector’ dataset. Although it measured carbon emissions in different sectors, it presented a sketchy view of USA CO<sub>2</sub> emission because it was collected in terms of years and provinces [17].

The organize China Emission Accounts and Datasets (CEADs) sorted out the CO<sub>2</sub> emission among Chinese provinces between 1997 and 2021. As same as the USA data introduced above, this dataset also does not have detailed statistics in time and space [18].

Another website also provides CO<sub>2</sub> emission of China, however, it offers more fine-grained data in grid. Unfortunately, just few areas were selected such as Yangtze River Delta economic belt and Pearl River Delta economic belt. This is a hindrance for us to show the whole world’s CO<sub>2</sub> emission as much as possible [19].

### 2.2 CO<sub>2</sub> Emission Prediction Models

In this part, we review the literature on CO<sub>2</sub> emission prediction methods. The methods are classified into three types. They are statistical models, traditional machine learning methods, and deep learning models.

**Statistical Models** Statistical models, including regression analysis and time series modeling, are commonly used in CO<sub>2</sub> emission prediction [26].

Lotfalipour et al. [20] predicted the CO<sub>2</sub> emissions using the grey model and the AutoRegressive Integrated Moving Average (ARIMA) model, and found that the forecasting performance of the grey model is better. Kour et al. [21] used ARIMA model to predict CO<sub>2</sub> emissions in South Africa. Hosseini et al. [22] used two regression models, multiple linear regression and multiple polynomial regression to predict Iran's CO<sub>2</sub> emissions in 2030.

Zhao et al. and Yang et al. [23, 24] used the STIRPAT model to predict CO<sub>2</sub> emissions, and built different scenarios to investigate the trajectories and peak times of CO<sub>2</sub> emissions in the future under different policy impacts.

Ding et al. [25] proposed a grey multivariable model by making three improvements, including modifying the calculation formula of the background values and optimizing the time response function based on the least-error law, to quantify future Chinese CO<sub>2</sub> emissions from fuel combustion. They further proposed an information-based grey model that integrates the damping accumulative generating operator, data smoothing index, and particle swarm optimization, which can be used to predict the carbon emissions of China's 30 provinces [26].

In general, the regression analysis and ARIMA models are suitable for predicting linear relationships. The STIRPAT model can achieve higher accuracy if the input dataset satisfies preconditions. The grey model is suitable for small sample data for short time series. These statistical models are not suitable for predicting sequences of CO<sub>2</sub> emissions with complex nonlinear relationships.

**Traditional Machine Learning Methods** The traditional machine learning methods used models such as Support Vector Regression (SVR), Random Forest (RF), gradient-boosted trees, and shallow neural networks to predict CO<sub>2</sub> emissions, which aims to better predict non-linear relationships within the time series data.

Li et al. [5] predicted city-level CO<sub>2</sub> emissions in China based on a set of open-access data. They first used two feature selection techniques, Recursive Feature Elimination, and Boruta, to extract important features for modeling CO<sub>2</sub> emissions. Then, they trained four machine learning models, namely SVR, RF, Gradient Boosting Machine (GBM), and extreme Gradient Boosting (XGBoost), to predict CO<sub>2</sub> emissions, finding that the XGBoost model achieved the highest estimation accuracy.

Song et al. [6] employed the correlation heatmap and Boruta methods to extract critical variables for modeling. Then, they used the stacking integration framework to integrate several single models, including LightGBM, CatBoost, and NGBoost, to improve the prediction accuracy and model generalization.

Khajavi et al. [7] compared response surface methodology, RF, and SVR model to predict CO<sub>2</sub> emission produced from road transport sector in China. They found that RF model with the Slime Mould Algorithm had the best prediction accuracy.

Traditional machine learning methods require feature selection to establish CO<sub>2</sub> prediction models. Feature selection relies on a significant amount of socioeconomic and energy-related statistical data. Challenge arises when needed data are unavailable and the feature selection is unpredictable, making it difficult to do accurate predictions.

Ren et al. [8] first calculated the carbon emissions and carbon emission intensity of Guangdong province, China from 1995 to 2019. Then, they constructed a fastlearning network improved by chicken swarm optimization to predict carbon emissions from

2020 to 2060 and explored the peak and neutralization of carbon emissions in nine scenarios. Although this shallow network does not need to select features, its performance is not better than deep learning models.

**Deep Learning Models** With the bloom of deep learning, deep neural networks have been proposed to leverage multiple hidden layers to automatically extract intricate data features. Among them, Recurrent Neural Network (RNN) is a type of neural network architecture designed to handle sequential data, which has achieved remarkable success in time series prediction [9]. LSTM (Long Short-Term Memory) and Gated Recurrent Unit (GRU) are two variants of the RNN that address the vanishing gradient problem and capture long-term dependencies in sequential data [10, 11].

Huang et al. [27] applied the LSTM to predict CO<sub>2</sub> emissions and then found LSTM had a better prediction performance than back propagation neural network and Gaussian process regression.

Zuo et al. [28] proposed an LSTM-STIRPAT model to analyze the CO<sub>2</sub> emission peak of 30 Chinese provinces in China. Initially, the LSTM model is used to predict the CO<sub>2</sub> emissions. According to the prediction results, the provinces are divided into some with peak value and others without peak value. Finally, the empirical analysis of the STIRPAT model was adopted to identify the main drivers of CO<sub>2</sub> emissions.

Wen et al. [29] proposed a hybrid model that integrates ARIMA and LSTM to forecast future carbon emission trends in China. This approach utilizes the strengths of both ARIMA and LSTM to capture linear and nonlinear relationships within the carbon emission data. The final prediction results are obtained by combining the inverse errors of ARIMA and LSTM with the goal of improving prediction accuracy.


## 3 Methodology

### 3.1 Framework

Figure 1 shows the research framework of the CO<sub>2</sub> prediction method. Initially, daily CO<sub>2</sub> emission data at the city level in China are collected. Then, the dataset is pre-processed, including invalid data deletion and data aggregation. After that, the Autoformer model is adopted to predict the city-level CO<sub>2</sub> emission data on a daily basis. Finally, we compare Autoformer with the ARIMA, the variants of the RNN-based models, and the other variants of the Transformer-based models to show the better performance of it. Moreover, the prediction results are visualized through several graphs to demonstrate an intuitive comparison result.

### 3.2 Data Collection and Preprocessing

Carbon Monitor is a near-real-time daily CO<sub>2</sub> emission dataset that shows the variations in CO<sub>2</sub> emissions from fossil fuel combustion and cement production at the national level [12]. Daily CO<sub>2</sub> emissions are estimated from a wide range of activity data, including electricity generation data, production data of energy-intensive industrial products,



**Fig. 1.** Model comparison & Result visualization.

ground transportation mobility data, maritime and air transportation activity data, and fuel consumption data.

GRACED is the near real-time global gridded daily CO<sub>2</sub> emissions dataset from fossil fuel and cement production [13]. Gridded fossil emissions are calculated for different sectors based on the daily national CO<sub>2</sub> emissions from Carbon Monitor, the spatial patterns of the point source emission dataset, the emission database for global atmospheric research, and the spatiotemporal patterns of satellite NO<sub>2</sub> retrievals [12].

Based on the Carbon Monitor and GRACED datasets, Huo et al. [12–14] constructed the is a near-real-time and city-level CO<sub>2</sub> emission dataset called Carbon Monitor Cities (CM-Cities). CM-Cities provides daily emission estimates from 2019 to 2021 for 1500 cities in 46 countries and disaggregates into five sectors: power generation, residential, industry, ground transportation, and aviation. When delineating city boundaries, CM-Cities establishes two types of spatial scopes: Functional Urban Areas (FUA) and Global Administrative Areas (GADM), which provide information to distinguish community-wide emissions and administrative emissions.

We use the GADM dataset of China from the CM-Cities datasets, which employs the GADM level-2 administrative areas for prefecture-level cities in China [14].


The data needs to be preprocessed. The cities that have empty values in sectors during the study period from 2019 to 2021 are ignored. The missing values are mainly in the power sector during the whole year of 2021. We obtain 300 cities with complete CO<sub>2</sub> emission data, and sum up the values of CO<sub>2</sub> emission in five sectors to obtain the total values of CO<sub>2</sub> emission in one day. Figure 2 shows the 300 cities on the map.

### 3.3 Our Model

Our model is built on the foundation of Autoformer since its families have shown great modeling ability for long-range dependencies and interactions in sequential data and thus are appealing to time series modeling [33].



**Fig. 2.** 300 cities in China.



**Fig. 3.** Model's structure.

Vaswani et al. [30] proposed the vanilla Transformer with the encoder-decoder structure. Both the encoder and decoder are composed of multiple identical blocks. Each encoder block consists of a multi-head self-attention module and a position-wise feed-forward network while each decoder block inserts cross-attention models between the multi-head self-attention module and the position-wise feed-forward network.

Many variants of Transformer have been proposed to address special challenges in time series modeling, especially the Autoformer model.

Wu et al. [31] studied the long-term forecasting problem of time series and proposed Autoformer, which introduces a decomposition transformer architecture and replaces the attention module with an Auto-Correlation mechanism.

Based on Autoformer, we added a time feature. Figure 3 shows our model's structure. Generally, we concatenate the CO2 emission sequence with the time feature as a combined input metric. However, the time series needs to be processed into a metric. The detailed method is that, the time series is divided into three dimensions: year, month and date, each dimension is encoded into the range between  $-0.5$  and  $0.5$ , putting them together into a metric.

### 3.4 Evaluation Metrics

The evaluation metrics used in this study are MSE, MAE, and  $R^2$ , where  $n$  denotes the length of test sets and  $y_i$  and  $\hat{y}_i$  denote the test set's real and predicted values, respectively. The lower the values of the first two measures, the better the fit.

- a) RMSE (Root Mean Square Error): It reflects the mean error of prediction with outlier sensitivity. The calculation of RMSE is given in Eq. 1:

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n \left( y_i - \hat{y}_i \right)^2} \quad (1)$$

- b) MAE (Mean Absolute Error): It reflects the mean error of prediction with outlier insensitivity. The calculation of MAE is given in Eq. 2:

$$MAE = \frac{\sum_{i=1}^n |y_i - \hat{y}_i|}{n} \quad (2)$$

## 4 Result


### 4.1 Experimental Details

We have compared Autoformer with other baselines in both traditional machine learning and deep learning fields, including ARIMA, GRU, RNN, Informer and Transformer [9, 11, 21, 30, 32]. For GRU, Informer, Transformer, and ours, we choose the same sequence length and prediction length, which is 1 step prediction for every 7 steps sequence. Also, for these models, we divide the datasets into three sections – training, validation and testing – the ratio between them is 7:1:2. While for RNN, we choose the same prediction length (1 step), but change the sequence to 6 steps, which shows a better performance. As for ARIMA and RNN, to meet the needs of the models, we only divide them into two sections – training and testing, and the ratio is 8:2.


### 4.2 Model Parameter Regulation

**ARIMA** To determine the optimal parameters of the ARIMA model, the Autocorrelation Function (ACF) and Partial Autocorrelation Function (PACF) are used to observe the dataset. ACF and PACF can measure the correlation between a time series and its lag. Therefore, the characteristics of the dataset can be observed by analyzing the results of the ACF and PACF of the dataset. Figure 4 shows the ACF and PACF results obtained from the time series of a randomly selected city (Aksu) among all 300 cities. Through analyzing Fig. 4, the number of autoregressive terms( $p$ ) and moving average terms( $q$ )





**Fig. 4.** ACF and PACF results.



**Fig. 5.** MAE in test set.

are determined between 1 and 5, and the differential order( $r$ ) is determined between 0 and 2. After multiple experiments with different parameters, the optimal parameters of ( $p, q, r$ ) is (2, 1, 1).

**GRU** During the parameter tuning process of GRU, we set the epoch number to 100, and the output results of models with different virous parameters have a common characteristic: the prediction error of the test set will increase significantly after the epoch number is larger than 40, as shown in Figs. 5 and 6. Therefore, 40 is the final epoch number we selected. The optimal parameters of GRU are batch\_size of 32 and gru\_unit of 64.

**Transformer, Informer and Ours** Transformer, Informer and ours use the same model parameter regulation method, comparing the prediction results of models with different label\_len. The final optimal label\_len of Transformer and Informer is 2, and it is 3 of ours.




Fig. 6. RMSE in test set.

Table 1. RMSE and MAE in different models.


	RMSE	MAE
ARIMA	10.14261225	7.651988617
RNN	6.90614478	4.64482946
GRU	11.507702124	5.187694
Transformer	8.231267646	5.40434041
Informer	8.200191370	5.27341824
<b>Autoformer</b>	<b>6.856545364</b>	<b>4.51108246</b>

4.3 Result Comparison and Visualization


RMSE and MAE results of all models are reported in Table 1, and our model has a relatively accurate performance. To show the results more vividly, we chose Shanghai as a representative to do the visualization, and the visual result is shown in Figs. 7, 8, and 9. Within the figures, the first line of the charts shows the prediction results and the real data, the second line of the charts shows the error value of the prediction.

From Fig. 7, we can clearly see that ARIMA exhibits a significant error, showing an over-smoothing prediction and having no ability in seasonal and abrupt data prediction. While in Figs. 8 and 9 show that the other models can predict the trends, whereas ours performs the best in predicting the peak and valley values and has the best ability to predict the fluctuation data. It can be found by looking at the error line. Additionally, the models from GRU, RNN to Transformer, and Informer display an increasing error in long-term predicting, while our model barely shows. Our model has the smallest error value and the fitted value are quite close to real value at the end of the time line.

In order to best present the error distribution, we draw scatter charts to show the deviation value of the models in Fig. 10. Comparing the scatter charts, it is easy to find that our model has the fewest error outliers, with R2 0.44 and RMSE 28.10, which means that it has the best reliability in prediction.



**Fig. 7.** Visualization for ARIMA.



**Fig. 8.** Visualization for GRU and RNN.

Besides, we create a comparative map (Fig. 11) of 300 cities to show the overall situation of our model's performance, finding that basically all the cities are well forecast.

#### 4.4 Synergistic and Integrated

In our experiment, we divide the datasets into three sections, training, validation and testing sections, and the ratio between them is 7:1:2. In order to compare all the models at their best level, we regulate their parameters to ensure each of them is in their best performances, using RMSE and MAE as criterion. Then we do visualization to present our results more understandable and directly. Also, we do the comparison to verify our model is the best either in data or through visualization.




Fig. 9. Visualization for Transformer, Informer and Autoformer.





Fig. 10. Errors in different models.



**Fig. 11.** Map comparing true and predicted values on June 1st, 2021.

## 5 Conclusion

Accurate and precise prediction of CO<sub>2</sub> emissions in cities on a daily basis provides the most up-to-date and fine-grained overview of the change in CO<sub>2</sub> emissions. In this paper, we use the Autoformer framework to predict the long-term daily dynamic change process of CO<sub>2</sub> emissions, which has a high prediction accuracy.

A city-level daily CO<sub>2</sub> emission dataset covering 3 years of CO<sub>2</sub> emission in 300 cities in China is used to demonstrate the effectiveness of our method. The method can be applied to other cities around the world.

Our prediction model focuses on predicting total CO<sub>2</sub> emissions in cities without distinguishing between specific sectors. In the future, we plan to propose new models that support sectoral CO<sub>2</sub> emissions in more detail. Besides, in this study, we construct the prediction model for each city separately. The carbon emission among cities may have potential spatial relationships. In the future, we plan to use the relationship between the spatial distribution characteristics of CO<sub>2</sub> emissions and the economic activities among cities to support the spatiotemporal prediction of CO<sub>2</sub> emissions.

**Acknowledgement.** Shuyi Wei and Xin You contributed equally to this work and should be considered co-first authors.

## References

1. Mehmood, I., Bari, A., Irshad, S., et al.: Carbon cycle in response to global warming. *Environ. Clim. Plant Veg. Growth*, 1–15 (2020)
2. Al-Ghussain, L.: Global warming: review on driving forces and mitigation. *Environ. Prog. Sustain. Energy*. **38**(1), 13–21 (2019)

3. Dong, K., Hochman, G., Timilsina, G.R.: Do drivers of CO<sub>2</sub> emission growth alter overtime and by the stage of economic development? *Energy Policy*. **140**, 111420 (2020)
4. Nyambuu, U., Semmler, W.: Climate change and the transition to a low carbon economy—carbon targets and the carbon budget. *Econ. Model.* **84**, 367–376 (2020)
5. Li, Y., Sun, Y.: Modeling and predicting city-level CO<sub>2</sub> emissions using open access data and machine learning. *Environ. Sci. Pollut. Res.* **28**, 19260–19271 (2021)
6. Song, Z., Wu, Y., Hua, C., et al.: A multi-factor prediction model for carbon productivity based on stacking integration method. In: 2022 5th International Conference on Pattern Recognition and Artificial Intelligence (PRAI), pp. 83–93. IEEE (2022)
7. Khajavi, H., Rastgoo, A.: Predicting the carbon dioxide emission caused by road transport using a Random Forest (RF) model combined by Meta-Heuristic Algorithms. *Sustain. Cities Soc.* **93**, 104503 (2023)
8. Ren, F., Long, D.: Carbon emission forecasting and scenario analysis in Guangdong Province based on optimized Fast Learning Network. *J. Clean. Prod.* **317**, 128408 (2021)
9. Cho, K., Van Merriënboer, B., Gulcehre, C., et al.: Learning phrase representations using RNN encoder-decoder for statistical machine translation. In: Conference on Empirical Methods in Natural Language Processing (EMNLP) (2014)
10. Yu, Y., Si, X., Hu, C., et al.: A review of recurrent neural networks: LSTM cells and network architectures. *Neural Comput.* **31**(7), 1235–1270 (2019)
11. Cho, K., Van Merriënboer, B., Bahdanau, D., et al.: On the properties of neural machine translation: Encoder-decoder approaches. arXiv preprint arXiv:1409.1259. (2014)
12. Liu, Z., Ciais, P., Deng, Z., et al.: Carbon Monitor, a near-real-time daily dataset of global CO<sub>2</sub> emission from fossil fuel and cement production. *Sci. Data.* **7**(1), 392 (2020)
13. Dou, X., Wang, Y., Ciais, P., et al.: Near-real-time global gridded daily CO<sub>2</sub> emissions. *The Innovations.* **3**(1) (2022)
14. Huo, D., Huang, X., Dou, X., et al.: Carbon Monitor Cities near-real-time daily estimates of CO<sub>2</sub> emissions from 1500 cities worldwide. *Sci. Data.* **9**(1), 533 (2022)
15. Search Results | World Resources Institute. World Resources Institute, [www.wri.org/search?keys=CO2&f%5B0%5D=content\\_type%3Adata](https://www.wri.org/search?keys=CO2&f%5B0%5D=content_type%3Adata)
16. data.europa.eu. [data.europa.eu/data/datasets/data\\_national-emissions-reported-to-the-unfccc-and-to-the-eu-greenhouse-gas-monitoring-mechanism-14?locale=en](https://data.europa.eu/data/datasets/data_national-emissions-reported-to-the-unfccc-and-to-the-eu-greenhouse-gas-monitoring-mechanism-14?locale=en)
17. U.S. Energy Information Administration. [www.eia.gov/opendata/browser/co2-emissions/co2-emissions-aggregates?frequency=annual&data=value;&sortColumn=period;&sortDirection=desc](https://www.eia.gov/opendata/browser/co2-emissions/co2-emissions-aggregates?frequency=annual&data=value;&sortColumn=period;&sortDirection=desc)
18. China Emission Accounts and Datasets. [www.ceads.net/user/search.php?kwtype=0&pagelang=en&searchtype=titlekeyword&typeid=45&q=co2](https://www.ceads.net/user/search.php?kwtype=0&pagelang=en&searchtype=titlekeyword&typeid=45&q=co2). Accessed 12 May 2024
19. MEICGreenhouseGases – MEICModel. [meicmodel.org.cn/?page\\_id=2345](https://meicmodel.org.cn/?page_id=2345)
20. Lotfalipour, M.R., Falahi, M.A., Bastam, M.: Prediction of CO<sub>2</sub> emissions in Iran using grey and ARIMA models. *Int. J. Energy Econ. Policy.* **3**(3), 229–237 (2013)
21. Kour, M.: Modelling and forecasting of carbon-dioxide emissions in South Africa by using ARIMA model. *Int. J. Environ. Sci. Technol.* **20**(10), 11267–11274 (2023)
22. Hosseini, S.M., Saifoddin, A., Shirmohammadi, R., et al.: Forecasting of CO<sub>2</sub> emissions in Iran based on time series and regression analysis. *Energy Rep.* **5**, 619–631 (2019)
23. Zhao, L., Zhao, T., Yuan, R.: Scenario simulations for the peak of provincial household CO<sub>2</sub> emissions in China based on the STIRPAT model. *Sci. Total Environ.* **809**, 151098 (2022)
24. Yang, F., Shi, L., Gao, L.: Probing CO<sub>2</sub> emission in Chengdu based on STRIPAT model and Tapio decoupling. *Sustain. Cities Soc.* **89**, 104309 (2023)
25. Ding, S., Dang, Y.G., Li, X.M., et al.: Forecasting Chinese CO<sub>2</sub> emissions from fuel combustion using a novel grey multivariable model. *J. Clean. Prod.* **162**, 1527–1538 (2017)

26. Ding, S., Zhang, H.: Forecasting Chinese provincial CO<sub>2</sub> emissions: a universal and robust new-information-based grey model. *Energy Econ.* **121**, 106685 (2023)
27. Huang, Y., Shen, L., Liu, H.: Grey relational analysis, principal component analysis and forecasting of carbon emissions based on long short-term memory in China. *J. Clean. Prod.* **209**, 415–423 (2019)
28. Zuo, Z., Guo, H., Cheng, J.: An LSTM-STRIPAT model analysis of China's 2030 CO<sub>2</sub> emissions peak. *Carbon Manag.* **11**(6), 577–592 (2020)
29. Wen, T., Liu, Y., he Bai, Y., et al.: Modeling and forecasting CO<sub>2</sub> emissions in China and its regions using a novel ARIMA-LSTM model. *Heliyon.* **9**(11) (2023)
30. Vaswani, A., Shazeer, N., Parmar, N., et al.: Attention is all you need. *Adv. Neural Inf. Proces. Syst.* **30** (2017)
31. Wu, H., Xu, J., Wang, J., et al.: Autoformer: Decomposition transformers with auto-correlation for long-term series forecasting. *Adv. Neural Inf. Proces. Syst.* **34**, 22419–22430 (2021)
32. Zhou H, Zhang S, Peng J, et al. Informer: Beyond efficient transformer for long sequence time-series forecasting. *Proceedings of the AAAI conference on artificial intelligence.* 2021, 35(12): 11106–11115
33. Wen, Q., Zhou, T., Zhang, C., et al.: Transformers in time series: A survey. *The 32nd International Joint Conference on Artificial Intelligence (IJCAI 2023).* 6778–6786 (2023). <https://arxiv.org/abs/2202.07125>