## Tutorial 2

# $\begin{array}{c} {\rm Shaun\ Schreiber} \\ 16715128 \end{array}$

#### 12 February 2014


## Question 1

$$Q = \{s,q1,q2,r1,r2\}$$
 
$$\sum_{} = \{a,b\}$$
 
$$\delta =$$

$$\begin{array}{c|cccc} & a & b \\ \hline s & q1 & q2 \\ q1 & q1 & q2 \\ q2 & q1 & q2 \\ r1 & r2 & r1 \\ r2 & r2 & r1 \\ \end{array}$$

$$\begin{array}{l} q1=s \\ F=\{q1,r1\} \end{array}$$

#### Question 2a






Figure 1: Constains ab as a substring. DFA =  $\{\{q1, q2, q3\}, \{a, b\}, \{(q1, b) -> q1, (q1, a) -> q2, (q2, a) -> q2, (q2, b) -> q3, (q3, b) -> q3, (q3, a) -> q3\}, q1, \{q3\}\}$ 

 $\begin{array}{l} {\rm Figure~2:~Constains~ba~as} \\ {\rm a~substring} \\ {\rm DFA} = \{\{q1,\,q2,\,q3\},\,\{a,\,b\},\,\{(q1,\,a)\,-\,>\,q1,\,(q1,\,b)\,-\,>\,q2,\,(q2,\,b)\,-\,>\,q2,\,(q2,\,a)\,-\,>\,q3,\,(q3,\,a)\,-\,>\,q3,\,(q3,\,b)\,-\,>\,q3\},\,q1,\,\{q3\}\} \end{array}$ 




Figure 3: Constains neither the substrings ab nor ba

### Question 2b




Figure 4: Only allows the string a

Figure 5: Only allows the string b




Figure 6: Allowing any strings except a and b

#### Question 3a




Figure 7: Has the length of at least 3 and its third symbol is a 0

# Question 3b




Figure 8: Contains at least two 0s and at most one 1