MTH 316 Notes

Evan Fox

July 12, 2022

Chapter 1

Cosets and Lagrange's theorem

1.1 Cosets

Definition 1.1.1 (Cosets)

Let H be a subgroup of G then for each $a \in G$ we define the left coset of H as

$$aH = \{ah \in G | h \in H\}$$

We will find this definition usefull in our study of groups. First we prove basic propertys of cosets.

Theorem 1.1.2 (Properties of Cosets)

- 1. aH = H iff $a \in H$.
- 2. aH = bH iff $a \in bH$
- 3. o(aH) = o(H).
- 4. aH = bH or $aH \cap bH = \emptyset$.

Proof. (1) Assume aH = H, then since $e \in H$, $ae = a \in aH$ thus $a \in H$. Conversly assume that $a \in H$, then for $ah \in aH$ we have by closure $ah \in H$. Simillarly if $h \in H$ then $ha^{-1} \in H$ and so $h \in aH$. For (2) The proof is similar. For (3) we want to find a bijection from aH to H. Define $\phi : H \to aH$ as $\phi(h) \mapsto ah$. Then one may easily show this is a bijection. (4) Assume $aH \neq bH$, then for the sake of contradiction assume that $aH \cap bH \neq \emptyset$. We fix x in the intersection and note that $x = ah_1$ and $x = bh_2$. Then $b = ah_1h_2^{-1} \in aH$, but then by (2) aH = bH; a contradiction.

Note that the last two parts of the theorem show that the cosets of a subgroup H partition the group G, later we will find out when it makes sense to define an operation on these equivalence classes.

1.2 Lagrange's Theorem

This is often called the A, B, Cs in finite group theory and one of the biggest results in MTH 316.

Theorem 1.2.1 (Lagrange) Let $H \leq G$ then o(H)|o(G).

Proof. Let $H \leq G$. Using the result from the last section we already know that the cosets of H partition G. Now consider the set of left cosets of H in G.

$$\{g_1H, g_2H, \ldots g_rH\}$$

Since $a \in aH$ we know that each element of G is in at least one coset, then since the cosets are disjoint,

$$ro(H) = o(G)$$