

Understanding Analysis Chapter 1

Evan Fox (efox20@uri.edu)

December 25, 2021

The first part of chapter one just went over some prelims/reviews. Here are my solutions to selected exercises.

Question 1.

(a) Prove that the $\sqrt{3} \notin \mathbb{Q}$

First, I prove that $3|x^2 \implies 3|x$ as I will need this fact in my proof of the irrationality of $\sqrt{3}$.

Proof. We use the contrapositive, so assume that $3 \nmid x$. Then there exists $k \in \mathbb{Z}$ such that $x = 3k + 1$ or $x = 3k + 2$. Then note that if $x = 3k + 1$, then

$$x = 3k + 1$$

$$x^2 = 9k^2 + 6k + 1 = 3(3k^2 + 2k) + 1$$

and in the other case we have

$$x = 3k + 2$$

$$x^2 = 9k^2 + 12k + 4 = 3(3k^2 + 4k + 1) + 1$$

So, in both cases we see that $3 \nmid x^2$ as desired. \square

Now I prove that $\sqrt{3} \notin \mathbb{Q}$

Proof. For the sake of contradiction, assume that $\sqrt{3} \in \mathbb{Q}$. Then we may fix $m, n \in \mathbb{Q}$ such that $\sqrt{3} = \frac{m}{n}$. We then have that,

$$3 = \left(\frac{m}{n}\right)^2$$

by the fundamental theorem of arithmetic, we may write m^2 and n^2 in terms of their prime factors and cancel any factor(s) that they have in common, i.e., we reduce $\frac{m^2}{n^2}$ such that they have no common factors. Since m and n have no common factors we note that they cannot both be divisible by 3. First observe that

$$3n^2 = m^2 \tag{1}$$

and hence we have $3|m^2$ which implies $3|m$; so we fix $k \in \mathbb{Z}$ such that $m = 3k$ and then we substitute this expression for m back into (1).

$$3n^2 = (3k)^2 = 9k^2 \tag{2}$$

$$n^2 = 3k^2 \tag{3}$$

and we have that 3 divides n^2 and by extension 3 divides n . Thus we have a contradiction as desired. \square

- (b) Does a similar argument work to prove that $\sqrt{6} \notin \mathbb{Q}$? Where does the proof breakdown for $\sqrt{4}$?

Yes, whether or not this method works for \sqrt{x} is related to the prime factorization of x , since the prime factors of 6 both have an exponent of 1, $6|m^2 \implies 6|m$ will hold. This is because if 2 and 3 are prime factors of m^2 then they will have to be prime factors of m , otherwise how would they have been prime factors of m^2 ? The point is squaring a number doesn't add new prime factors; it just multiplies the exponent of each prime factor by 2. When we try to apply this argument to $\sqrt{4}$ the problem is that $4|m^2 \implies 4|m$ doesn't hold since the exponent of the prime factor of 4 is 2. To give an example note that $4|36 = 6^2 = 2^2 3^2$ but $4 \nmid 6 = 3(2)$.

Question 2.

Prove that there is no rational number satisfying $2^r = 3$.

Proof. We use contradiction, so assume that there exists $r \in \mathbb{Q}$ such that $2^r = 3$. Then since r is rational by assumption we may fix $m, n \in \mathbb{Q}$ such that $r = \frac{m}{n}$. Substituting in for r gives

$$2^{\frac{m}{n}} = 3$$

then we raise both sides to the n^{th} power and get

$$2^m = 3^n$$

which contradicts the uniqueness of the fundamental theorem of arithmetic. \square

Question 3.

The triangle inequality is given by $|a + b| \leq |a| + |b|$.

- (a) Verify the triangle inequality in the special case that a and b have the same sign.

Proof. Let $a, b \in \mathbb{R}$ and assume that a and b have the same sign. Then if a and b are both negative we see that $|a + b| = a + b$ and if a and b are both positive then $|a + b| = a + b$. Then note that regardless of the signs of a and b , $|a| + |b| = a + b$. It is then clear that the inequality holds. \square

- (b) Give a general proof the the triangle inequality.

Proof. First we prove that $(a+b)^2 \leq (|a|+|b|)^2$ by observing $ab \leq |ab|$. Then we multiply both sides by 2 and obtain $2ab \leq 2|ab|$, we then add $a^2 + b^2$ to both sides and get $a^2 + 2ab + b^2 \leq |a|^2 + 2|ab| + |b|^2$. Factoring this gives $(a+b)^2 \leq (|a|+|b|)^2$. We can now use this to prove the triangle inequality by taking the square root of both sides which yeilds,

$$|a + b| \leq ||a| + |b||$$

but since $|a| + |b| \geq 0$ we have

$$|a + b| \leq |a| + |b|$$

as desired. \square

- (c) Use the triangle inequality to prove that $|a - b| \leq |a - c| + |c - d| + |d - b|$.

Proof. Let $x = (a - c)$ and let $y = (c - d) + (d - b)$, the the triangle inequality tells us that

$$\begin{aligned} |x + y| &\leq |x| + |y| \\ |a - b| &\leq |a - c| + |(c - d) + (d - b)| \end{aligned}$$

Now we can apply the triangle inequality again to the second term on the right hand side of the last equation. So we let $z = (c - d)$ and let $w = (d - b)$, by the triangle inequality we the have $|c - b| \leq |c - d| + |d - b|$. Substituting back in gives,

$$\begin{aligned} |a - b| &\leq |a - c| + |(c - d) + (d - b)| = |a - c| + |c - d| + |d - b| \\ |a - b| &\leq |a - c| + |c - d| + |d - b| \end{aligned}$$

as desired.

□

Question 4.

Given a function f and a subset of its domain A , let $f[A]$ denote the range of f over A , i.e., $f[A] = \{f(x) | x \in A\}$

- (a) let $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x^2$. Let $A = [0, 2]$ and let $B = [1, 4]$. Does $f[A \cap B] = f[A] \cap f[B]$? What about $f[A \cup B] = f[A] \cup f[B]$?

Proof. Both parts of this question are true. First I prove $f[A \cap B] = f[A] \cap f[B]$. First note that $A \cap B = [1, 2]$ then we may compute the range of f on this domain and we get that $\text{ran } f = [1, 4]$.

To prove that $\text{ran } f = [1, 4]$ first we prove that $\text{ran } f \subseteq [1, 4]$ so let $y \in \text{ran } f$. Then we fix $x \in \text{dom } f$ such that $y = f(x) = x^2$. Then since f is increasing on the interval $[1, 2]$ it is clear that $y \in [1, 4]$. Now to prove that $[1, 4] \subseteq \text{ran } f$ let $y \in [1, 4]$ be arbitrary. Then we must find a $x \in \text{dom } f$ such that $f(x) = y$. Choosing $x = \sqrt{y}$ gives the desired result. Since $f(\sqrt{y}) = (\sqrt{y})^2 = y$. Since \sqrt{y} is well defined on $[0, 4]$ we conclude that $\text{ran } f = [1, 4]$. Hence $f[A \cap B] = [1, 4]$. □

- (b) Give an example where $f[A \cap B] = f[A] \cap f[B]$ doesn't hold.

Example: let $A = [-1, -2]$ and $B = [1, 2]$.

- (c) Let $A, B \subseteq \mathbb{R}$ and prove that for an arbitrary function $g : \mathbb{R} \rightarrow \mathbb{R}$, $g[A \cap B] \subseteq g[A] \cap g[B]$.

Proof. Let $y \in g[A \cap B]$ be arbitrary. Then there exists an $x \in A \cap B$ such that $y = g(x)$. Since $x \in A \cap B$ we know that $x \in A$ and $x \in B$. We also have that $y = g(x)$. Hence $y \in g[A]$ and $y \in g[B]$. Then we have that $y \in g[A] \cap g[B]$. \square

*side note: if I add the condition that g be injective, I think that you could prove $g[A \cap B] = g[A] \cap g[B]$. A counter example existed for $f(x) = x^2$ only because it is not injective (I think).

- (d) Form a conjecture about $g[A \cup B]$ and $g[A] \cup g[B]$ and prove it.

Conjecture: $g[A \cup B] = g[A] \cup g[B]$.

Proof. Let $y \in g[A \cup B]$ then we may fix $x \in A \cup B$ such that $y = g(x)$. Since we have that $x \in A$ or $x \in B$, we know that either $y \in g[A]$ or $y \in g[B]$. It is then clear that y is in the union. To prove the other direction assume that $y \in g[A] \cup g[B]$, then either $y \in g[A]$ or $y \in g[B]$. If $y \in g[A]$ then we may fix $x \in A$ such that $y = g(x)$. Then it follows that $x \in A \cup B$; So then $y \in g[A \cup B]$. If it is not the case that $y \in g[A]$ then $y \in g[B]$ must be true and the argument is the same. \square

Question 5.

s