

Algebra

Evan Fox

July 21, 2022

Chapter 1

Rings

1.1 Integral domains and division rings

Definition 1.1.1

Let R be a set and define $+$: $R \times R \rightarrow R$ denoted $(a, b) \mapsto a + b$ and \cdot : $R \times R \rightarrow R$ denoted $(a, b) \mapsto ab$ and assume

1. $(R, +)$ is an abelian group
2. (R, \cdot) is associative and closed
3. $a(b + c) = ab + ac$ and $(b + c)a = ba + ca$

Then R is a ring.

If the multiplication on R is commutative then we say that R is a commutative ring. Note in general there need not exist $1 \in R$ satisfying $1 \cdot a = a$ for all $a \in R$, since (R, \cdot) is not necessarily a group. An informal way of defining a ring is just to consider a set where addition, subtraction, and multiplication make sense.

Remark 1.1.2

Let R be a ring, for $a, b \in R$ we have

1. $0b = 0$
2. $(-a) \cdot b = -(a \cdot b)$
3. $(-a) \cdot (-b) = ab$

Proof. For the first statement note

$$0b = (0 + 0)b = 0b = 0b + 0b$$

since R is a group under addition we may make full use of the cancelation property which gives $0 = 0b$. Next observe,

$$(-a)b + ab = ((-a) + a)b = 0b = 0$$

so that ab is the additive inverse of $(-a)b$. The proof of the last result is simillar. □

The definition of a ring is very general, some classic examples are the integers (mod n), matrices, and continous functions. Showing these are rings is easy.

Remark 1.1.3

Let R be a ring and S a non-empty set. Recall R^S denotes the set of functions $f : S \rightarrow R$. Give a addition and multiplication that turns R^S into a ring such that if $|S| = 1$ we have $R^S \sim R$.

Proof. exercise. □

Now we will introduce a very important type of ring called an integral domain. First we must consider the concept of a zero divisor.

Definition 1.1.4

Let R be a ring and $a \in R$ with $a \neq 0$. If there exists non-zero $b \in R$ such that $ab = 0$, then a is a zero divisor.

Definition 1.1.5

Integral Domain A ring R is an integral domain if

1. R is communitive and $1 \in R$.
2. R has no zero divisiors.

An obvious example of an integral domain is the integers. One can show that that the deffinition of an integral domain is equivalent to a communitive ring where multiplicative cancelation holds.

Proposition 1.1.6

Let R be a communitive ring, then R is an integral domain iff $ab = ac \implies b = c$ for all $a, b, c \in R$ with a non-zero.

Proof. For the first direction assume that R is an integral domain and let $a \neq 0, b, c \in R$ satisfy $ab = ac$. Then

$$ab - ac = 0 \implies a(b - c) = 0$$

and since a is non zero and R has no zero divisors we have $b - c = 0$ so $b = c$.

Conversly, assume that the cancellation property holds in R . We need to prove that $ab = 0$ implies $a = 0$ or $b = 0$; so it is sufficient to assume $ab = 0$ and that $a \neq 0$ and show $b = 0$. We know that

$$ab = a0 = 0$$

and the cancellation property shows that $b = 0$. □

Definition 1.1.7

A ring R (not necessarily commutative) is a division ring if $(R^*, 0)$ is a group

If R is a commutative division ring then we say that R is a field. Common examples of fields are \mathbb{Q}, \mathbb{R} , and \mathbb{C} . A more exotic example is the quadratic field $\mathbb{Q}(\sqrt{D}) := \{a+b\sqrt{D} | a, b \in \mathbb{Q}\}$ where D is a non-square rational number. The proof that this is a field is left as an exercise.

Theorem 1.1.8

A finite integral domain is a field.

Proof. Let R be a finite integral domain and let x_1, \dots, x_n be a list of all elements. Fix $a \neq 0 \in R$ and consider the list

$$S = ax_1, \dots, ax_n$$

Since R is an integral domain if there exists $i, j \in \{1, \dots, n\}$ such that $ax_i = ax_j$ we have

$$a(x_i - x_j) = 0 \implies x_i = x_j.$$

Hence every element in S is distinct, then since the length of S is the same as the cardinality R we see that every element of R appears in S . Clearly a appears in this list, hence there must exist x_{j_0} such that $ax_{j_0} = a$. Now note that for any $r \in R$ we can write $r = ax_k$ and

$$rx_{j_0} = ax_{j_0}x_k = ax_k = r$$

so x_{j_0} is the unit of R . Now we must find the multiplicative inverse of a , since $x_{j_0} \in R$ it appears in S . Hence there exist $x_{i_0} \in R$ such that $ax_{i_0} = x_{j_0}$; it follows that every non zero element of R has a multiplicative inverse. Thus R is a field □

The above theorem is clearly false in the infinite case (consider the integers), however an interesting note is where the above proof breaks down. In the finite case the function $\cdot_a : R \rightarrow R$ defined by $r \mapsto a \cdot r$ is an injective mapping from R to R and is then necessarily a bijection. However for sets with infinite cardinality, you can have a injective map from a set to itself that is not surjective; and this is where the logic of the above breaks down.

A field is a very important structure in mathematics and we will have much more to say about them later.

1.2 Ideals and Quotients

In group theory we had subgroups, to extend this to rings we simply require clousure of multiplication.

Definition 1.2.1

Let R be a ring and suppose $S \subseteq R$. We say S is a subring of R if

1. *S is a subgroup of R when considered under addition*
2. *S is closed under multiplication*

We may want to test wether or not a given subset is a subring, here we show a way of doing that (we have basicaly just stolen the subgroup test.)

Theorem 1.2.2

Subring test $S \subseteq R$ is a subring if

1. *$a, b \in S \implies a - b \in S$ and $ab \in S$*

Proof. exercise □

We will find that the concept of subrings is not so important. What we really want to study are the ideals of a ring. In group theory we found the concept of a normal subgroup to be extremly usefull, so it is natural to ask if we can produce a similar concept for rings such that we may bring over many of our previous results. If we consider R as a group it is clear that every subgroup is normal (since R is abelian), but what should we require of multiplication? We will soon see that the next deffinition is the correct one.

Definition 1.2.3

Ideals of a ring Let R be a ring. A (two-sided) ideal of R is a subring U such that for all $r \in R$ and all $u \in U$ $ru \in U$ and $ur \in U$.

Remark 1.2.4

Technically we may speak of left ideals and right ideals, were U would only be closed under multiplication on the left or right, this is why we must add the condition of a two sided ideal. In these notes we will only consider the notion of two sided ideals. Note in a commutative ring a left and right ideal coincide and there is no difference.

Now given an ideal U we know that the cosets $a + U$ for all $a \in R$ partition R and we know how to turn R/U into a group. Now we need to define a multiplication on R/U to turn it into a ring. Obviously we would like

$$(a + U)(b + U) = (ab) + U$$

lets show that our definition of ideal ensures that this will work

Proposition 1.2.5

R

U with the above operations is a ring

Proof. we must show that the definition of multiplication is well defined. Let $a, a', b, b' \in R$ and assume $a + U = a' + U$ and $b + U = b' + U$. We need to show that

$$(ab) + U = (a'b') + U$$

Note $a \in a + U$ so $a = a' + u_s$ and the same argument shows $b = b' + u_t$ for $u_s, u_t \in U$. Now for $x \in (ab) + U$ we have

$$\begin{aligned} x &= ab + u_1 = (a' + u_2)(b' + u_3) + u_1 \\ &= a'b' + a'u_3 + u_2b' + u_2u_3 = a'b' + u_4 \in (a'b') + U. \end{aligned}$$

Which shows $(ab) + U \subseteq (a'b') + U$. The converse is similar and this completes the proof. □

Hence multiplication works as expected. The above proof also makes clear why we defined ideals the way we did. Now we move on to consider two more types of ideals, prime ideals and maximal ideals.

Definition 1.2.6

Prime Ideals Let I be an ideal of a ring R , we say I is a prime ideal if all $a, b \in R$ we have $ab \in I$ implies $a \in I$ or $b \in I$.

Prime ideals get their name from the ideals $p\mathbb{Z} \subseteq \mathbb{Z}$. An easy exercise is to show that for all prime p , $p\mathbb{Z}$ is prime.

Definition 1.2.7

Maximal Ideals Let M be an ideal. We say that M is maximal if $M \subseteq U \subseteq R$ implies $M = U$ or $M = R$.

An ideal is maximal when it is impossible to fit another ideal between it and the entire ring.

We immediately see two results

Theorem 1.2.8

I is a prime ideal of a ring R iff R/I is an integral domain

Proof. Suppose that I is a prime ideal then $(a + I)(b + I) = I = (ab) + I$ hence a or b is in I so either $(a + I) = I$ or $(b + I) = I$. Conversely let R/I be a ring and I an ideal. Now suppose R/I

R/I is an integral domain. Now assume $ab \in I$. Then $I = (ab) + I = (a + I)(b + I)$ and since R/I

R/I is an integral domain one of $a + I$ or $b + I$ must equal I and it follows from this that either $a \in I$ or $b \in I$. \square

For maximal ideals we see an even stronger result

Theorem 1.2.9

M is a maximal ideal if and only if R/M is a field.