

THETA RHYTHM IS WIDESPREAD IN VOCAL PRODUCTION ACROSS THE ANIMAL REALM.

THÉOPHANE PIETTE^{*1}, CHUNDRA CATHCART², CHIARA BABRIERI³, DIDIER
GRANDJEAN⁴, ÉLOÏSE DÉAUX¹, ANNE-LISE GIRAUD¹

^{*}Corresponding Author: theophane.piette@unige.ch

¹Department of Basic Neuroscience, University of Geneva, Geneva, Switzerland

²Center for the Interdisciplinary Study of Language Evolution, University of Zurich,
Zurich, Switzerland

³ Department of Evolutionary Biology and Environmental Studies, University
of Zurich, Zurich, Switzerland

⁴Swiss Center for Affective Sciences, University of Geneva, Geneva,
Switzerland

1. Introduction

Speech production rate, conserved between 2 to 8 syllables per second, plays a key role in comprehension. Recent studies that highlighted the presence of the same rhythm in vocal productions and mouth movements of non-human primates, suggest that it may be inherited from mastication (Risueno-Segovia et al, 2020). However, other factors could constrain vocal rhythm. Notably, syllabic rate also matches the frequency of theta brain oscillations (2 -7 Hz). While oscillations are rhythmic electrical activity spontaneously generated by neural tissues, they can also be stimulus-driven and research has demonstrated that theta oscillations entrain to speech during listening (Giraud et al, 2014). As brain oscillations frequencies are conserved in animals, syllabic rate may instead be constrained by the recipient's neuronal activity (Buzaski et al, 2013). To determine what factor(s) may have contributed to the emergence of the theta vocal rhythm, we collected vocalizations for 90 species of animals. We used phylogenetic regression to assess whether weight, mastication or environmental constraints explain the variation in animal vocal rhythms while controlling for phylogenetic relationships.

2. Methods

2.1 *Vocal, physical and environmental data*

We gathered, when they were available, recordings for one species per infra-order of tetrapods as well as some fishes and insects (N=92, mammal = 24, bird = 58, amphibian = 4, reptile =1, insect = 4 and fish= 1). To obtain vocal production

rhythm, we then extracted sequences, consisting of one subject vocalizing for at least 2s, with less than 1s of silence between two sounds. For each sequence, the envelope, was computed, and analyzed using a wavelet transform to extract the peak rhythmic changes of amplitude (Tilsen et al 2014). Bodyweight data were collected (N=92), habitat type was defined as either closed, semi-closed, open, wetland or water, and mastication status as yes or no. We obtained data on the context of production, but these were only available for 35% of the recordings. Therefore, we assumed that such effect would relate to flexible rhythm adjustment within an already physically constrained rhythmic range (Martin et al, 2022).

2.2 Statistical analyses

Given the extent of the current dataset, we are still in the process of building the corresponding phylogenetic tree. Thus using log-transformed weight, mastication, environments, and their interactions as fixed effects, and random slopes of log weight by groups and random intervals for orders, we built a linear mixed model investigating the variation of the median log-transformed rhythm frequency.

3. Results

We can observe in our preliminary results that most of our species (74.4%) have a median frequency located between 2Hz to 7Hz. When testing our linear mixed model, we noticed that our random effects did not explain the variance of rhythm in our species. We also found that none of our predictors significantly influenced vocal rhythm (Most impacting factor: Environment, $F=1.0570$, $p\text{-value}=0.3748$).

4. Conclusion

Theta production rhythm seems widespread across animal vocal productions, independently of their physical characteristics, mastication status or environment. As it is present in most species, even the ones that do not masticate, it appears unlikely that this rhythm is constrained by a limit of the natural movement of the articulators. Furthermore, the lack of effect of weight and environmental constraints tends to point towards a more common and endogenous constraint for vocalization rhythm. As brain oscillations and neurons characteristics, in general, are well conserved across species, it could be that the rhythm of the vocalizations has adapted to them as a form of 'neuronal exploitation', to ensure the efficient reception by the listener. As the range of oscillations in animals' brains is not documented for most species, further work on phylogeny and the direct link between vocalizations and brain oscillations in animals will be needed to strengthen this hypothesis or lead the way for alternative explanations.

Acknowledgements

The NCCR Evolving Language, Swiss National Science Foundation Agreement Nr. #51NF40_180888, funded this work. The authors are grateful to the CSIRO Australian National Wildlife Collection, (<https://ror.org/059mabc80>), Fonoteca Zoologica (<https://www.fonozoo.com/>), the Museum für Naturkunde of Berlin (<https://www.museumfuernaturkunde.berlin>) and the Xeno-Canto Foundation for Nature Sounds (<https://xeno-canto.org>) for their assistance in undertaking this research. The authors thank all the researchers from the NCCR Evolving Language as well as Adrien Meguerditchian, Pascal Belin and Steffen R.Hage for their insights on the project. The authors would like to thank Aaron Bauer, Andrew Spencer, Alex Kwet, Alex Rohtla, Camila Ferrara, Daniel Blumstein, Élodie Briefer, Émilie Genty, Frank Lambert, Isabelle Charrier, Gerald Carter, Hans Schneider, Marc D. Hauser, Nikola Falk, Peter Boesman and Robert Seyfarth for providing us with recordings for this study. The authors also thank two anonymous reviewers for their helpful comments on an earlier version of the manuscript.

References

- Buzsáki, G., Logothetis, N., & Singer, W. (2013). Scaling brain size, keeping timing: evolutionary preservation of brain rhythms. *Neuron*, 80(3), 751-764. <https://doi.org/10.1016/j.neuron.2013.10.002>
- Fujii, S., & Wan, C. Y. (2014). The role of rhythm in speech and language rehabilitation: the sep hypothesis. *Frontiers in Human Neuroscience*, 8. <https://doi.org/10.3389/fnhum.2014.00777>
- Giraud, A.-L., & Poeppel, D. (2012). Cortical oscillations and speech processing: emerging computational principles and operations. *Nature Neuroscience*, 15(4), 511-517. <https://doi.org/10.1038/nn.3063>
- Hoyo, J. del, Elliott, A., Sargatal, J., & Cabot, J. (Éds.). (1992). *Handbook of the birds of the world*. Lynx Edicions.
- Martin, M., Gridley, T., Elwen, S. H., & Charrier, I. (2022). Feel the beat: Cape fur seal males encode their arousal state in their bark rate. *The Science of Nature*, 109(1), 5. <https://doi.org/10.1007/s00114-021-01778-2>
- Noujaim, S. F., Lucca, E., Muñoz, V., Persaud, D., Berenfeld, O., Meijler, F. L., & Jalife, J. (2004). From mouse to whale: a universal scaling relation for the pr interval of the electrocardiogram of mammals. *Circulation*, 110(18), 2802-2808. <https://doi.org/10.1161/01.CIR.0000146785.15995.67>
- Ravosa, M. J., Ross, C. F., Williams, S. H., & Costley, D. B. (2010). Allometry of masticatory loading parameters in mammals. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology*, 293(4), 557-571. <https://doi.org/10.1002/ar.21133>

- Risueno-Segovia, C., & Hage, S. R. (2020). Theta synchronization of phonatory and articulatory systems in marmoset monkey vocal production. *Current Biology*, 30(21), 4276-4283.e3. <https://doi.org/10.1016/j.cub.2020.08.019>
- Tilsen, S., & Johnson, K. (2008). Low-frequency Fourier analysis of speech rhythm. *The Journal of the Acoustical Society of America*, 124(2), EL34-EL39. <https://doi.org/10.1121/1.2947626>
- Wilson, D. E., Mittermeier, R. A., & Cavallini, P. (Éds.). (2009). *Handbook of the mammals of the world*. Lynx Edicions: Conservation International: IUCN.
- Worthington, J., Young, I. S., & Altringham, J. D. (s. d.). Short communication the relationship between body mass and ventilation rate in mammals. 4.