

Thesis Introduction

Ewan Pinnington

September 30, 2016

1 Notation

List of symbols and meanings consistent throughout thesis.

2 The global carbon cycle

Carbon is one of the most abundant elements, making up around half of all living matter dry mass on Earth. The global carbon cycle describes the movement of carbon through the Earth system. In the Earth system large amounts of carbon are present in the oceans, atmosphere, land surface and crust. These stores of carbon are referred to as reservoirs or pools. The amount of carbon on Earth can be considered constant, as under terrestrial conditions it is not common to have nuclear transmutation. Therefore terrestrial processes using carbon must transfer it between the global carbon pools, this is referred to as a flux. In pre-industrial times these fluxes of carbon between different pools have remained approximately constant over long time scales REF.

The greenhouse gas effect describes the process by which radiatively active gases (CO₂, water vapour, ozone, etc.) in the Earth's atmosphere contribute to the warming of the planet by absorbing long-wave radiation emitted from the Earth's surface and reradiating this absorbed energy in all directions, causing more warming below [Mitchell, 1989]. The natural greenhouse gas effect raises the global mean surface temperature by 30K, making the Earth habitable for the many lifeforms upon it. The increase in atmospheric greenhouse gases since the industrial revolution due to anthropogenic activities has amplified the greenhouse effect and caused global warming. CO₂ has been found to be the most important human-contributed compound to this warming [Falkowski et al., 2000]. In figure 1 we show a simplified schematic of the global carbon cycle taken from the Intergovernmental Panel on Climate Change (IPCC), in this schematic we can see the large rise in atmospheric CO₂ since the industrial revolution with an increase of 240 Pg C up to 2011.

As atmospheric CO₂ levels have risen, natural sinks of CO₂ (fluxes out of the atmosphere) have intensified with both the land surface and oceans absorbing more CO₂ from the atmosphere. This can be seen in figure 1, with the net ocean flux of CO₂ to the atmosphere decreasing from an estimated +0.7 Pg C yr⁻¹ to -2.3 Pg C yr⁻¹ and the land surface flux of CO₂ to the atmosphere decreasing from -1.7 Pg C yr⁻¹ to -2.6 Pg C yr⁻¹. More recent estimates from Le Quéré et al. [2015] indicate these sinks have further intensified with the ocean sink estimated to be 2.9 ± 0.5 Pg C yr⁻¹ and the land surface sink 4.1 ± 0.9 Pg C yr⁻¹ for the year 2014. The intensification of the land carbon sink is thought to be partly due to a combination of forest regrowth as well as rising CO₂ and increased nitrogen deposition having a fertilisation effect [Ciais et al., 2014]. It has also been shown that the land surface sink has been enhanced by an increase in diffuse photosynthetically active radiation as a result of increased cloud cover associated with increased anthropogenic emissions [Mercado et al., 2009]. It is vitally important to understand the response of these sinks of CO₂ to

climate change in the future. If either the oceans or land surface were to stop absorbing this same percentage of CO₂ from the atmosphere we would see even more dramatic increases in atmospheric CO₂ levels and thus a much greater rate of global warming. Some estimates even show the land surface changing from a sink of CO₂ to a source of CO₂ under certain future emission scenarios [Sitch et al., 2008].

Land surface carbon uptake is the least understood process in the global carbon cycle [Ciais et al., 2014], this can be seen in the uncertainties given in Figure 1. In current estimates of the global carbon budget land surface carbon uptake is estimated by taking the residual of all other calculated sources and sinks of carbon, so that

$$S_{LAND} = E_{FF} + E_{LUC} - (G_{ATM} + S_{OCEAN}) \quad (1)$$

where S_{LAND} is the global residual land sink of CO₂, E_{FF} is the CO₂ emissions from fossil fuels, E_{LUC} is the CO₂ emissions from land use change (mainly deforestation), G_{ATM} is the atmospheric CO₂ growth rate and S_{OCEAN} is the mean ocean CO₂ sink [Le Quéré et al., 2015]. In Figure 2 we can see the growth in this residual land sink with increased emissions, it can also be seen that there is a high variability in this sink, largely due to year to year variations in precipitation, surface temperature, radiation and volcanic eruptions.

Land use change is the next most uncertain component in the global carbon cycle and the second largest anthropogenic source of CO₂. It is not well understood how much CO₂ is removed from the atmosphere by regrowth of previously deforested land (either by felling or fire), although it is thought that regrowth forests could be stronger carbon sinks than old growth forests due to more rapid biomass accumulation under succession [Pan et al., 2011]. Better understanding the response of the land surface to disturbance will help constrain future carbon budgets.

Good paper to highlight the fact that the percentage of CO₂ absorbed by the land surface has remained approximately constant with rising atmospheric CO₂ levels??? Maybe just use IPCC fig 6.8?

In figure 6.1 and 6.8: Partitioning of fluxes important and hard (shown by error on estimates in fig 6.1). Land surface carbon uptake least understood mechanism in the global carbon cycle, ref IPCC. Will uptake remain the same under climate change.

Human emissions of CO₂ have perturbed the global C cycle and caused a large continual increase in atmospheric CO₂ levels.

Look at papers recommended on Flux Course, some good ones to reference???

IPCC figure 6.1 and 6.8: Partitioning of fluxes important and hard (shown by error on estimates in fig 6.1). Land surface carbon uptake least understood mechanism in the global carbon cycle, ref IPCC. Will uptake remain the same under climate change.

Booth et al. [2012] Have shown that global warming is highly sensitive to land carbon cycle processes and highlighted the need to improve understanding of land surface carbon uptake and its response to climate change.




Figure 1: Global carbon cycle simplified schematic [Ciais et al., 2014]. Black numbers and arrows represent reservoir mass and exchange fluxes estimated for the time prior to the industrial era (~ 1750). Red numbers and arrows represent annual fluxes average over the 2000-2009 time period. Red numbers in the reservoirs indicate the cumulative change of carbon over the industrial period (1750-2011).




Figure 2: Annual anthropogenic CO₂ emissions and their partitioning among the atmosphere, land and ocean from 1750 to 2011 [Ciais et al., 2014].

3 The role of models




Figure 3: Modelled land sink [Ciais et al., 2014].

IPCC figure 6.16 and section 6.3.2.6.6: Contribution of models to understanding the terrestrial carbon cycle. Reference every DALEC paper.

4 Eddy covariance and other observations

Baldocchi paper: Many observations of forest carbon flux made worldwide.

5 Data assimilation

Role of DA in NWP improving forecast skill.

References

- B. B. B. Booth, C. D. Jones, M. Collins, I. J. Totterdell, P. M. Cox, S. Sitch, C. Huntingford, R. A. Betts, G. R. Harris, and J. Lloyd. High sensitivity of future global warming to land carbon cycle processes. *Environmental Research Letters*, 7(2):024002, 2012.
- P. Ciais, C. Sabine, G. Bala, L. Bopp, V. Brovkin, J. Canadell, A. Chhabra, R. DeFries, J. Galloway, M. Heimann, et al. Carbon and other biogeochemical cycles. In *Climate change 2013: the*

- physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pages 465–570. Cambridge University Press, 2014.
- P. Falkowski, R. J. Scholes, E. Boyle, J. Canadell, D. Canfield, J. Elser, N. Gruber, K. Hibbard, P. Högberg, S. Linder, F. T. Mackenzie, B. Moore III, T. Pedersen, Y. Rosenthal, S. Seitzinger, V. Smetacek, and W. Steffen. The global carbon cycle: A test of our knowledge of earth as a system. *Science*, 290(5490):291–296, 2000. ISSN 0036-8075. doi: 10.1126/science.290.5490.291.
- C. Le Quéré, R. Moriarty, R. M. Andrew, J. G. Canadell, S. Sitch, J. I. Korsbakken, P. Friedlingstein, G. P. Peters, R. J. Andres, T. Boden, et al. Global carbon budget 2015. *Earth System Science Data*, 7(2):349–396, 2015.
- L. M. Mercado, N. Bellouin, S. Sitch, O. Boucher, C. Huntingford, M. Wild, and P. M. Cox. Impact of changes in diffuse radiation on the global land carbon sink. *Nature*, 458(7241):1014–1017, 04 2009. URL <http://dx.doi.org/10.1038/nature07949>.
- J. F. Mitchell. The “greenhouse” effect and climate change. *Reviews of Geophysics*, 27(1):115–139, 1989.
- Y. Pan, R. A. Birdsey, J. Fang, R. Houghton, P. E. Kauppi, W. A. Kurz, O. L. Phillips, A. Shvidenko, S. L. Lewis, J. G. Canadell, et al. A large and persistent carbon sink in the world’s forests. *Science*, 333(6045):988–993, 2011.
- S. Sitch, C. Huntingford, N. Gedney, P. Levy, M. Lomas, S. Piao, R. Betts, P. Ciais, P. Cox, P. Friedlingstein, et al. Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five dynamic global vegetation models (dgvms). *Global Change Biology*, 14(9):2015–2039, 2008.