List of Papers

Ewan Pinnington

Today

First paper 'Breathing of the terestrial biosphere' [1]

- Eddy covariance measures the covariance between fluctuations in verticle velocity and the mixing ratio of trace gases of interest. Seasonal C flux.
- -ve values of F_n represent a loss of CO_2 from the atmos. and a gain by the surface. F_n net ecosystem exchange, F_a Gross canopy assimilation, F_r ecosystem respiration.
- Hard to find natural stands of vegetation that meet the right standards for good eddy covariance measurement. Problems with night time measurement. EC not able to measure fluxes of complex terrain.
- Disturbance (fire, drought) needs to be incorporated into models. Annual variability in carbon fluxes.
- Mid aged forests (50-100 years) are best carbon sinks.
- Network now contains more variaties of forest and vegetation so PD has shifted towards zero (fig. 4).

Second paper 'An Improved analysis of forest carbon dynamics using data assimilation' [18]

- Measurements from young pine stand in ponderossa oregon (warm dry summers wet cool winters).
- DA combining mments of C stocks made over times and m/ments of C flux made directly (EC), along with model.
- Ensemble Kalman filter being used.
- Results show significantly more accurate analysis when using DA than model alone.
- Monte-Carlo technique?
- dynamics of soil C and WD pools an area of uncertainty.
- No energy balance? lack of models to describe photosynthesis and plant growth.

Thesis Laura 'Correlated observation errors in data assimilation' [15]

- Chapter 2: Different methods of DA and approximations. Error covariances and correlations, Desroziers method?
- Chapter 3: Matrix approximations to include some correlations for R and avoid the need to explicitly calculate R^{-1} (computationally taxing). Calculate trace of analysis error covariance matrix, S_a , to understand how useful an observation set is to the DA scheme (smaller trace $S_a \implies$ better reduction of error variance). May have to calculate S_a^* if R known to be specified incorrectly. Look at vector and matrix norms.

The REFLEX project: comparing different algorithms and implementations for the inversion of a terrestrial ecosystem model against eddy covariance data [10]

• Data given to participants to use with DALEC and DALEC-D and their choice of MDF algorithm. Tristan's results not used?

Effect of correlated observation error on parameters, predictions, and uncertainty [17]

• Hydrology, Considering a denitrification model with diagonal and then full weight R matrix.

A regularization of the carbon cycle data-fusion problem [4]

• Sensitivity testing of DALEC model with 4D-VAR.

Diagnosis of obs, background and analysis-error statistics in obs space [5]

• Looking at $d_o^b = y^o - H(x^b)$ and $d_o^a = y^o - H(x^a)$ for finding R and B with covariances.

Dynamic Data Assimilation: a least squares approach [11]

- Try the least squares non linear second order joby, looks good, could help instead of using B just minimise for the observations and R pg 154 and pg 401
- Model adjoint Linear 4DVAR observation operators good fundamental DA reference.

SiPNET paper Braswell 2005 [2]

• More complicated Carbon balance model to look at and understand!

Estimating parameters of a forest ecosystem C model with measurements of stocks and fluxes as joint constraints [12]

- Reason Tristan wrote this project, can do better?
- Parameter estimation for DALEC using MC procedures.
- Woody biomass increment and to a lesser degree soil respiration measurements give marked reductions in uncertainties in parameter estimates and model predictions as they are 'orthogonal' constraints to NEE measurements.

• None of the data good at constraining fine root or soil C pool dynamics, need new measurements?

Observation error correlations in IASI radiance data [16]

• Satellite takes same obs. every time step, Flux tower variable? Mainly temporal correlations possible!

Data assimilation with correlated observation errors: experiments with a 1-D shallow water model [14]

• Use an idealised perfect model and approximate forms of R to include correlations.

Where does the carbon go? A model—data intercomparison of vegetation carbon allocation and turnover processes at two temperate forest free-air CO2 enrichment sites [3]

• Use temperate forest free-air CO2 enrichment (FACE) to test. How better to control allocation parameters, allocation should be controlled by functions rather than parameters.

Information content of infrared satellite sounding measurements with respect to CO₂ [6]

• Use information content measures to look at satellite measurements

Estimation of entropy reduction and degrees of freedom for signal for large variational analysis systems [8]

• Information content measure for operational data assimilation schemes

Measures of observation impact in Gaussian data assimilation [9]

• Information content meausre for Gaussian data assimilation schemes, nice proofs and explanation!

A Practical Method to Estimate Information Content in the Context of 4D-Var Data Assimilation. I: Methodology [13]

• Information content meausre for 4Dvar, nice proofs and explanation! Degrees of freedom for signal good job!

Estimation of entropy reduction and degrees of freedom for signal for large variational analysis systems [7]

• FIND AND READ! DOFS and other information measures, uses the form I am used to $n - tr(B^{-1}A)$

References

- [1] Dennis Baldocchi. Turner review no. 15. 'breathing' of the terrestrial biosphere: lessons learned from a global network of carbon dioxide flux measurement systems. Australian Journal of Botany, 56(1):1–26, 2008.
- [2] Bobby H Braswell, William J Sacks, Ernst Linder, and David S Schimel. Estimating diurnal to annual ecosystem parameters by synthesis of a carbon flux model with eddy covariance net ecosystem exchange observations. *Global Change Biology*, 11(2):335–355, 2005.
- [3] Martin G De Kauwe, Belinda E Medlyn, Sönke Zaehle, Anthony P Walker, Michael C Dietze, Ying-Ping Wang, Yiqi Luo, Atul K Jain, Bassil El-Masri, Thomas Hickler, et al. Where does the carbon go? a model—data intercomparison of vegetation carbon allocation and turnover processes at two temperate forest free-air co2 enrichment sites. *New Phytologist*, 2014.
- [4] Sylvain Delahaies, Ian Roulstone, and Nancy Nichols. A regularization of the carbon cycle data-fusion problem. In *EGU General Assembly Conference Abstracts*, volume 15, page 4087, 2013.
- [5] Gérald Desroziers, Loic Berre, Bernard Chapnik, and Paul Poli. Diagnosis of observation, background and analysis-error statistics in observation space. Quarterly Journal of the Royal Meteorological Society, 131(613):3385–3396, 2005.
- [6] RJ Engelen and GL Stephens. Information content of infrared satellite sounding measurements with respect to co2. *Journal of Applied Meteorology*, 43(2):373–378, 2004.
- [7] Michael Fisher. Estimation of entropy reduction and degrees of freedom for signal for large variational analysis systems. European Centre for Medium-Range Weather Forecasts, 2003.
- [8] Michael Fisher and European Centre for Medium Range Weather Forecasts. Estimation of entropy reduction and degrees of freedom for signal for large variational analysis systems. Reading, England: European Centre for Medium-Range Weather Forecasts, 2003. Cover title.
- [9] Alison Fowler and Peter Jan van Leeuwen. Measures of observation impact in gaussian data assimilation. 2011.
- [10] Andrew Fox, Mathew Williams, Andrew D Richardson, David Cameron, Jeffrey H Gove, Tristan Quaife, Daniel Ricciuto, Markus Reichstein, Enrico Tomelleri, Cathy M Trudinger, et al. The reflex project: comparing different algorithms and implementations for the inversion of a terrestrial ecosystem model against eddy covariance data. Agricultural and Forest Meteorology, 149(10):1597–1615, 2009.
- [11] John M Lewis, Sivaramakrishnan Lakshmivarahan, and Sudarshan Dhall. *Dynamic data as-similation: a least squares approach*, volume 13. Cambridge University Press, 2006.
- [12] Andrew D Richardson, Mathew Williams, David Y Hollinger, David JP Moore, D Bryan Dail, Eric A Davidson, Neal A Scott, Robert S Evans, Holly Hughes, John T Lee, et al. Estimating parameters of a forest ecosystem c model with measurements of stocks and fluxes as joint constraints. *Oecologia*, 164(1):25–40, 2010.

- [13] Adrian Sandu, Kumaresh Singh, Mohamed Jardak, Kevin Bowman, and Meemong Lee. A practical method to estimate information content in the context of 4d-var data assimilation. i: Methodology. 2012.
- [14] Laura M Stewart, Sarah Dance, and Nancy K Nichols. Data assimilation with correlated observation errors: experiments with a 1-d shallow water model. *Tellus A*, 65, 2013.
- [15] Laura M Stewart, SL Dance, and NK Nichols. Correlated observation errors in data assimilation. *International journal for numerical methods in fluids*, 56(8):1521–1527, 2008.
- [16] LM Stewart, J Cameron, SL Dance, S English, J Eyre, and NK Nichols. Observation error correlations in iasi radiance data. *Mathematics report series*, 1, 2009.
- [17] Claire R Tiedeman and Christopher T Green. Effect of correlated observation error on parameters, predictions, and uncertainty. Water Resources Research, 2013.
- [18] Mathew Williams, Paul A Schwarz, Beverly E Law, James Irvine, and Meredith R Kurpius. An improved analysis of forest carbon dynamics using data assimilation. *Global Change Biology*, 11(1):89–105, 2005.