

Data

Ewan Pinnington

January 22, 2017

1 Introduction

As part of this PhD an extended period of time has been spent at the Alice Holt Research Station (Hampshire, UK) working with Forest Research (The research arm of the UK Forestry Commission). After initially completing one year of an ongoing field campaign to measure stem respiration using an infra-red gas analyser, a measurement campaign was designed to produce a set of observations for use in this PhD project. This involved the establishment and sampling of three transects throughout the Straits Inclosure (part of the Alice Holt forest). The establishment of these transect and measurements are outlined in this chapter.

2 Alice Holt research site

The Alice Holt Forest is a research forest area managed by the UK Forestry Commission located in Hampshire, SE England. Forest Research have been operating a CO₂ flux measurement tower in a portion of the forest, the Straits Inclosure, continuously since 1998. The Straits Inclosure is a 90 ha area of deciduous broadleaved plantation woodland located on a surface water gley soil and was initially planted with oak in the 1820s [Schlich and Perrée, 1905] and then replanted in the 1930s. The majority of the canopy trees are oak (*Quercus robur* L.), with an understory of hazel (*Corylus avellana* L.) and hawthorn (*Crataegus monogyna* Jacq.) [Pitman and Broadmeadow, 2001], but there is a small area of conifers (*Pinus nigra* ssp. *laricio* (Maire) and *P. sylvestris* L.) within the tower measurement footprint area depending on wind direction. An aerial photograph of the site is shown in Figure 1. The Straits Inclosure is a flat area at an altitude of approximately 80m,

surrounded by mixed lowland woods and both arable and pasture agricultural land. In Wilkinson et al. [2012] an analysis of stand-scale 30 minute average net CO₂ fluxes (NEE) from 1998-2011 for the Straits Inclosure found a mean annual NEE of $-486 \text{ g C m}^{-2} \text{ yr}^{-1}$ and demonstrated the forest was a substantial sink of carbon. This study also includes further details about the research site.

As part of the management regime, the Straits Inclosure is subject to thinning, whereby a proportion of trees are removed from the canopy in order to reduce competition and improve the quality of the final tree crop. At the Straits an intermediate thinning method is used with a portion of both subdominant and dominant trees being removed from the stand [Kerr and Haufe, 2011]. The whole of the stand was thinned in 1995. Subsequently the eastern side of the Straits was thinned in 2007 and then the western side in 2014. The flux tower at the site is situated on the boundary between these two sides. This allows for the use of a footprint model to split the flux record and thus analyse the effect of this disturbance on carbon fluxes at the site. In Wilkinson et al. [2015] a statistical analysis of the eddy covariance flux record found that there was no significant effect on the net carbon uptake of the eastern side after thinning in 2007. In this thesis we focus on the effect of disturbance on the western side after thinning in 2014 in chapter REF. We therefore refer to the western side as “thinned” forest and the eastern side as “unthinned” forest.




Figure 1: The Straits Inclosure research site in 2013.

3 Establishment of sampling points

For this fieldwork transects were designed to join up existing mensuration plots where measurements of woody biomass are made by Forest Research. This allowed for comparison with historic observations. In total 435 sampling points were marked at 10m intervals, these are shown in Figure 2. Python was used to calculate the exact latitude and longitude of each sampling point for the 3 transects, these locations were then entered into a GPS unit. When establishing the transects fluorescent spray paint was used to mark trees closest to each sampling point as shown on the GPS (see Figure 3). As parts of the forest site were extremely dense with vegetation a pair of loppers were used to clear a path in some areas to allow for the construction of relatively straight transects. Having all transect points numbered and corresponding to a latitude and longitude value allowed for comparison between methods and the splitting of observations between different distinct sections of the forest site.




Figure 2: Sampling transects. Black crosses: sampling points at 10m intervals, pink diamonds: Forest Research mensuration plots, black diamond: Forest Research flux tower.



Figure 3: Sampling point 291, showing fluorescent spray paint used to mark sampling points.

4 Leaf area index observations

Leaf Area Index (LAI) is an important variable in relation to the amount of CO₂ an ecosystem can remove from the atmosphere through photosynthesis. LAI is defined as the area of leaves per unit area of ground. Three different methods were used to find estimates for peak LAI (July - September) for the year 2015 along the three transects at different sampling intervals.

4.1 Ceptometer

A Decagon LP-80 ceptometer and an additional Photosynthetically Active Radiation (PAR) sensor were used to measure LAI. Here we measure below canopy PAR using the ceptometer while logging above canopy PAR using a data logger and PAR sensor positioned outside the canopy. We can then calculate LAI using the above and below canopy readings and a set of equations relying on some assumptions [Fassnacht et al., 1994]. The ceptometer represents the quickest method for estimating LAI, we therefore took readings with the ceptometer at every sampling point over two walks of the transects, giving us 870 observations in total.

In order to be sure that the PAR readings from the ceptometer and external PAR sensor were consistent we had to calibrate the PAR sensor against the ceptometer. This was done by leaving both the PAR sensor and ceptometer out logging next to each other every 10 seconds for a day in the Alice Holt Research Station met square. We can then calibrate the output of the PAR sensor with that of the ceptometer as shown in Figure 4.

Once the PAR sensor was calibrated measurements could be made along the transects. The PAR sensor positioned outside of the canopy was logged every 5 seconds using a Delta-T DL2e data logger, at the start of every set of measurements the clock on the data logger and ceptometer were synchronised to ensure comparison of measurements made at the same time. After sampling the transects we had a set of above canopy and below canopy PAR readings corresponding to each sampling point for both walks of the transects. We use the same calculation for LAI as given in the Decagon LP-80 manual. This is using a simple model of radiation transmission and scattering tested against the more complex model of Norman and Jarvis [1975]. The equation used to calculate




Figure 4: Calibration of above canopy Photosynthetically Active Radiation (PAR) sensor (measuring in mV) with LP-80 ceptometer measured PAR ($\mu\text{mol m}^{-2} \text{s}^{-1}$).

LAI from the above and below canopy PAR readings is,

$$LAI = \frac{((1 - \frac{1}{2K})f_b - 1)\ln\tau}{A(1 - 0.47f_b)}, \quad (1)$$

where K is the extinction coefficient, f_b is the beam fraction, $\tau = \frac{\text{below canopy PAR}}{\text{above canopy PAR}}$ and $A = 0.283 + 0.785a - 0.159a^2$ (where a is the leaf absorptivity, assumed to be 0.9 by Decagon). We assume a spherical leaf angle distribution parameter, $\chi = 1$, this means the extinction coefficient simplifies to $K = \frac{1}{2\cos\theta}$, where θ is the solar zenith angle. We took the mean of the two LAI observations at each point to give as an estimate to the peak LAI for the year 2015. We can see the LAI estimate for the Straits Inclosure in Figure 5.





Figure 5: Ceptometer derived LAI for Alice Holt.

4.2 Hemispherical photographs

The next method used to measure LAI was hemispherical photography. Hemispherical photographs show a complete view of the sky in all directions, from these images we use the HemiView software [Rich et al., 1999] which calculates the proportion of visible sky as a function of sky direction (gap fraction) which it then uses to calculate LAI [Jonckheere et al., 2004]. Hemispherical photographs were taken every 50m along the transects, giving a total of 89 images. It is important to note that hemispherical photographs are taken in overcast conditions so that the sun does not obscure areas of leaf area. It is important to note that we did not remove tree trunks and branches from our calculation of LAI with HemiView so that we are actually calculating plant area index. The impacts of this assumption are discussed in section 4.4. In Figure 6 we show an example of two hemispherical photographs taken in different areas of the Straits Inclosure.

4.3 Litter traps

Finally litter traps were used to find estimates to LAI and leaf mass per area. Here we placed litter traps under the canopy to catch leaf litter as it falls into a bag attached to the bottom of the trap.



(a) Unthinned forest

(b) Thinned forest

Figure 6: Hemispherical photographs from the Alice Holt flux site showing the difference between the thinned and unthinned sides of the forest.

The bags were changed every week during the litter fall period and the litter sorted into species. Every week the litter was dried in an oven at 70°C and weighed. This gave us the dry-weight of the leaf litter for the 2015 season. Towards the end of the season we scanned a subsample for each species of 100 leaves to find an area, we then dried and weighed each subsample, a relationship between dry-weight and leaf area was then built (leaf mass per area) and used to infer the total LAI for each trap once the whole seasons litter has been collected. This method of LAI calculation is the most time consuming.

A total of six litter traps were established at points along the transects (positions shown in Figure 7) allowing for comparison with the other methods. The 6 litter traps are not enough to describe the LAI for the research site [Kimmings, 1973]. We use these litter traps as a point of comparison and validation for the ceptometer and hemispherical photograph estimates of LAI made at the same locations and also for estimates to leaf mass per area. From our litter trap observations we find a leaf mass per area of 29 g C m^{-2} free soluble carbohydrates for both sides of the forest.

4.4 Comparison of methods

In Figure 8 and 9 we show a comparison of the different methods of estimating LAI for the unthinned and thinned forest respectively. We can see that in all cases LAI derived from the litter traps is always greater than LAI estimated from optical methods, this is expected from previous comparisons [Bréda, 2003].

Although the ceptometer is the fastest method for measuring LAI it is also the most variable,




Figure 7: Litter trap locations for Alice Holt.

being extremely sensitive to the solar zenith angle and clear sky conditions. If the sun is low in the sky the radiation will pass through much more photosynthetically active material than if the sun is directly above head, causing spikes in the LAI value. We can see that the LAI estimates from the hemispherical photographs are much less variable than the ceptometer. As discussed in section 4.2 the hemispherical estimate is actually of plant area index, as we have not removed trunks and branches from the gap fraction calculation. However, this does not appear to have a great impact on results as hemispherical photograph derived LAI is still the lowest estimate of all three.




Figure 8: LAI comparison for unthinned forest. Dots and solid line represent observations made at different points along transects, dotted lines represent the mean of the observations.




Figure 9: LAI comparison for thinned forest. Dots and solid line represent observations made at different points along transects, dotted lines represent the mean of the observations.

5 Point-centred quarter observations

We used the method of Point-Centred Quarters (PCQ) [Dahdouh-Guebas and Koedam, 2006] to determine an estimate of the woody biomass for both unthinned and thinned forest in the Straits Inclosure. The PCQ method is conducted at each sampling point as follows:

- Using a compass, map 4 regions from the central sampling point
- Measure the distance from the central sampling point to the nearest tree in each quarter
- Measure the Diameter at Breast Height (DBH) for each tree (shown in Figure 10) and record the species

There were 114 points samples along the three transects, from these measurements we derived estimates to tree density and mean DBH for both thinned and unthinned sides of the forest. We then used allometric relationships between DBH and total above ground biomass and coarse root biomass, found in work carried out by Forest Research and in McKay et al. [2003]. These relationships were as follows,

$$\text{above ground dry-mass} = 0.0678 \times \overline{\text{DBH}}^{2.619} \quad (2)$$

and

$$\text{below ground coarse root dry-mass} = 0.149 \times \overline{\text{DBH}}^{2.12}. \quad (3)$$

This gave us an estimate to the dry-mass in kilograms for the average tree in our sampling area. Assuming that half of all dry-mass is carbon we can find an estimate of total woody and coarse root carbon in g C m^{-2} using the equation,

total woody and coarse root carbon =

$$1000 \times 0.5 \times (\text{above ground dry-mass} + \text{below ground coarse root dry-mass}) \times \text{tree density}. \quad (4)$$

Forest Research have carried out their own mensuration studies at the site, these have been conducted at the mensuration points shown in Figure 2. As these plots are included in our transects this means that hopefully our measurements will be comparable with those from Forest Research.



Figure 10: Taking diameter at breast height measurements at Alice Holt.

6 Flux tower observations and data processing

Forest Research provided half-hourly raw flux tower data for the Straits Inclosure from January 1999 to December 2015. These consist of the NEE fluxes and meteorological driving data of temperature, irradiance and atmospheric CO₂ concentration for use in the DALEC model. The view from the top of the flux tower in the Straits Inclosure can be seen in Figure 11. Forest Research provided this data in the form of multiple excel spreadsheets corresponding to the flux tower measurement record for each year. To prepare this data for use with data assimilation we first had to convert these 16 excel files to one Python readable data file (here we chose NetCDF), this was then further processed. To process the NEE data we first performed u^* filtering, where any half-hourly flux observation corresponding to a friction velocity of 0.2 m s⁻¹ (this value represents the point at which flux measurements become unreliable and was found by Forest Research) or less were removed from the data set. We then subjected the observations of NEE to quality control procedures similar to those described by Papale et al. [2006]. For each year of the NEE dataset this procedure involved calculating the standard deviation of both the positive and negative half-hourly observations and

then removing any values that were ± 3 standard deviations away from the yearly positive/negative mean. This was also repeated on a month by month basis. Gap-filling procedures were not applied to the half-hourly NEE dataset so that only true observations were considered for assimilation. To match the time-step of the DALEC model we computed daily NEE observations by taking the mean over the 48 measurements made each day, selecting only days where there was no missing data.



Figure 11: At the top of the Alice Holt flux tower.

References

- N. J. Bréda. Ground-based measurements of leaf area index: a review of methods, instruments and current controversies. *Journal of experimental botany*, 54(392):2403–2417, 2003.
- F. Dahdouh-Guebas and N. Koedam. Empirical estimate of the reliability of the use of the point-centred quarter method (pcqm): Solutions to ambiguous field situations and description of the pcqm+ protocol. *Forest Ecology and management*, 228(1):1–18, 2006.
- K. S. Fassnacht, S. T. Gower, J. M. Norman, and R. E. McMurtrie. A comparison of optical

- and direct methods for estimating foliage surface area index in forests. *Agricultural and Forest Meteorology*, 71(1):183–207, 1994.
- I. Jonckheere, S. Fleck, K. Nackaerts, B. Muys, P. Coppin, M. Weiss, and F. Baret. Review of methods for in situ leaf area index determination Part I. Theories, sensors and hemispherical photography. *Agricultural and Forest Meteorology*, 121(1-2):19–35, 2004. ISSN 01681923. doi: 10.1016/j.agrformet.2003.08.027.
- G. Kerr and J. Haufe. Thinning practice: A silvicultural guide. *Forestry Commission*, page 54, 2011.
- J. Kimmins. Some statistical aspects of sampling throughfall precipitation in nutrient cycling studies in british columbian coastal forests. *Ecology*, pages 1008–1019, 1973.
- H. McKay, J. Hudson, and R. Hudson. Woodfuel resource in britain. *Forestry Commission Report*, 2003.
- J. Norman and P. Jarvis. Photosynthesis in sitka spruce (*picea sitchensis* (bong.) carr.): V. radiation penetration theory and a test case. *Journal of Applied Ecology*, pages 839–878, 1975.
- D. Papale, M. Reichstein, M. Aubinet, E. Canfora, C. Bernhofer, W. Kutsch, B. Longdoz, S. Rambal, R. Valentini, T. Vesala, et al. Towards a standardized processing of net ecosystem exchange measured with eddy covariance technique: algorithms and uncertainty estimation. *Biogeosciences*, 3(4):571–583, 2006.
- R. Pitman and M. Broadmeadow. Leaf area, biomass and physiological parameterisation of ground vegetation of lowland oak woodland. *Forestry Commission, Edinburgh*, 2001.
- P. M. Rich, J. Wood, D. Vieglais, K. Burek, and N. Webb. Hemiview user manual, version 2.1. *Delta-T Devices Ltd., Cambridge, UK*, 79, 1999.
- S. W. Schlich and W. Perrée. *Working plan for the Alice Holt Forest*. HM Stationery Office, 1905.
- M. Wilkinson, E. Eaton, M. Broadmeadow, and J. Morison. Inter-annual variation of carbon uptake by a plantation oak woodland in south-eastern england. *Biogeosciences*, 9(12):5373–5389, 2012.

M. Wilkinson, P. Crow, E. Eaton, and J. Morison. Effects of management thinning on co₂ exchange by a plantation oak woodland in south-eastern england. *Biogeosciences Discussions*, 12(19), 2015.