

Plasma multiphysics model (FM-WP2)

Michael Barnes¹, M. R. Hardman¹
J. Omotani², S. Newton², F. I.
Parra³

*¹Rudolf Peierls Centre for Theoretical
Physics, U. Oxford*

²Culham Centre for Fusion Energy

³Princeton Plasma Physics Laboratory

Limitations of current model

- Homogeneous, helical (or straight) magnetic field
- No intra-species collisions
- Assumed Boltzmann electron response
- Simple wall boundary condition
- Electrostatic fluctuations
- Simple models for collisions with neutrals

Limitations of current model

- **Homogeneous, helical (or straight) magnetic field**
- **No intra-species collisions**
- **Assumed Boltzmann electron response**
- Simple wall boundary condition
- Electrostatic fluctuations
- Simple models for collisions with neutrals

Unresolved numerical issues

- Current implementation of moment kinetic equations incompatible with wall BCs
- Inclusion of radial electric field leads to poor convergence near walls
- Parallelism of code could be improved?
- Anything else?

More realistic magnetic field geometry

- Keep helical geometry, but allow $B=B(r,z)$
- Requires treatment of non-periodic radial BC
- Introduces magnetic drifts and shear into the model
- New physics + sets up machinery for coupling to equilibrium solvers
- Allows for direct comparisons with established fluid (e.g., BOUT++) and kinetic (e.g., GKEYLL) codes

Dynamically evolved electrons

- Extant codes either use small term to solve for potential (via polarization density) or must solve iteratively:
;
- No explicit expression for electrostatic potential
- Rapid electron motion along field lines \Rightarrow small time steps
- Boltzmann model for electrons avoids possible numerical issues associated with evolution

Dynamically evolved electrons

- Extant codes either use small term to solve for potential (via polarization density) or must solve iteratively:
;
- We propose two approaches:
 - Treat electrons as a fluid (Braginskii closure and/or mass ratio expansion)
 - Moment-kinetic model with mass ratio expansion

Collisions between charged particles

- Wide range of collisionalities present in edge plasma: from collisional near material boundary to \sim collisionless near core
- Need to account for charged particle collisions, at least in intermediate region between fluid limit and collisionless limit
- Full Landau operator is nonlinear, integrodifferential
- Propose to explore hierarchy of models of increasing complexity: Krook, pitch-angle scattering, energy diffusion, conserving corrections

Miscellaneous related issues

- Transition between open and closed field line regions
- Transition between fluid and kinetic treatments
- Can moment kinetics be made to work with wall BC?
- More sophisticated kinetic wall BCs
- Parallelization/memory optimization
- Addition of further unit, integrated, and regression tests