

Quantum Field Theory Equation Sheet

quinten tupker

October 7 2020 - December 6, 2020

Here are some useful equations

Table 1: Equation Sheet

Name/Description	Equation	Remarks
Noether conserved current	$j^\mu = \partial_{\partial_\mu \phi} \mathcal{L} \delta \phi - F^\mu$	here $\mathcal{L}(x + \delta x) = \mathcal{L} + \delta x \partial_\mu F^\mu$, $\partial_\mu j^\mu = 0$
The conserved charge arising from a conserved current	$Q = \int d^3x j^0$	
The Energy-Momentum Tensor	$T^\mu_\nu = \partial_{\partial_\mu \phi} \mathcal{L} \partial_\nu \phi - \delta^\mu_\nu \mathcal{L}$	This is the Noether current under translation. This tensor can always be chosen to be symmetric. It is a Noether current, so conserved as $\partial_\mu T^{\mu\nu} = 0$
Ladder Operators	$[a_p, a_q^\dagger] = (2\pi)^3 \delta(p - q)$	
Field Operator	$\phi = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2\omega_p}} (a_p e^{ip \cdot x} + a_p^\dagger e^{-ip \cdot x})$	
Momentum Operator	$\pi = \int \frac{d^3p}{(2\pi)^3} (-i) \sqrt{\frac{\omega_p}{2}} (a_p e^{ip \cdot x} - a_p^\dagger e^{-ip \cdot x})$	
Dirac Equation	$(i\gamma^\mu \partial_\mu - m) \cdot \psi = (i\not{\partial} - m) \cdot \psi = 0$	
Chiral representation and Clifford Algebra	$\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu} I, \{\gamma^5, \gamma^\mu\} = 0, (\gamma^5)^2 = 0, \gamma^5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3, \gamma^0 = \begin{pmatrix} I & \\ & I \end{pmatrix}, \gamma^i = \begin{pmatrix} & \sigma^i \\ \sigma^i & \end{pmatrix}, \gamma^5 = \begin{pmatrix} I & \\ & -I \end{pmatrix}$	
General Solution to Dirac Equation	positive frequencies: $\psi(x) = u(p) e^{-ip \cdot x}, u^s(p) = \begin{pmatrix} \sqrt{p \cdot \bar{\sigma}} \xi^s \\ \sqrt{p \cdot \bar{\sigma}} \xi^s \end{pmatrix}$, for negative frequencies $\psi(x) = v(p) e^{ip \cdot x}, v^s(p) = \begin{pmatrix} \sqrt{p \cdot \bar{\sigma}} \eta^s \\ -\sqrt{p \cdot \bar{\sigma}} \eta^s \end{pmatrix}$	Here ξ^r, η^s form orthonormal bases for \mathbb{C}^2 and $\sqrt{p \cdot \bar{\sigma}} = \sqrt{m} e^{\chi \cdot \sigma / 2}, \sqrt{p \cdot \bar{\sigma}} = \sqrt{m} e^{-\chi \cdot \sigma}$