
a
BMI at diagnosis
AUCROC: 0.886
20
30
40
50
60
BMI (kg/m2) at diagnosis
b
Unintentional weight loss
AUCROC: 0.772
% unintentional weight loss
100
75
50
25
0
Type 1
Type 2
c
Waist−hip ratio
AUCROC: 0.771
0.50
0.75
1.00
1.25
1.50
Waist−hip ratio
d
Age at diagnosis
AUCROC: 0.768
20
30
40
50
Age at diagnosis (years)
e
HbA1c at diagnosis
AUCROC: 0.715
50
100
150
HbA1c (mmol/mol) at diagnosis
f
Parent with non−insulin−treated diabetes
AUCROC: 0.609
% Parent with non−insulin−treated diabetes
Type 1
Type 2
g
Osmotic symptoms
AUCROC: 0.605
% osmotic symptoms
Type 1
Type 2
h
DKA
AUCROC: 0.577
% DKA
Type 1
Type 2
i
Sex
AUCROC: 0.550
% female
Type 1
Type 2
j
Parent history of autoimmune disease
AUCROC: 0.541
% parent with autoimmune disease
Type 1
Type 2
k
Ethnicity
AUCROC: 0.536
%
Black
Mixed
Other
South Asian
White
l
Personal history of autoimmune disease
AUCROC: 0.520
% additional autoimmune disease
Type 1
Type 2