



SAPIENZA  
UNIVERSITÀ DI ROMA

“SAPIENZA” UNIVERSITY OF ROME  
FACULTY OF INFORMATION ENGINEERING,  
INFORMATICS AND STATISTICS  
DEPARTMENT OF COMPUTER SCIENCE

---

# Discrete Mathematics

---

Lecture notes integrated with the book "Discrete Mathematics",  
Norman L. Biggs

*Author*  
Simone Bianco

March 3, 2024

# Contents

<b>Information and Contacts</b>	<b>1</b>
<b>1 Introduction to Discrete Maths</b>	<b>2</b>
1.1 Solved exercises . . . . .	2

# Information and Contacts

Personal notes and summaries collected as part of the *Discrete Mathematics* course offered by the degree in Computer Science of the University of Rome "La Sapienza".

Further information and notes can be found at the following link:

<https://github.com/Exyss/university-notes>. Anyone can feel free to report inaccuracies, improvements or requests through the Issue system provided by GitHub itself or by contacting the author privately:

- Email: [bianco.simone@outlook.it](mailto:bianco.simone@outlook.it)
- LinkedIn: [Simone Bianco](#)

The notes are constantly being updated, so please check if the changes have already been made in the most recent version.

## Suggested prerequisites:

Preventive learning of material related to the *Algebra* course is recommended

## Licence:

These documents are distributed under the [GNU Free Documentation License](#), a form of copyleft intended for use on a manual, textbook or other documents. Material licensed under the current version of the license can be used for any purpose, as long as the use meets certain conditions:

- All previous authors of the work must be **attributed**.
- All changes to the work must be **logged**.
- All derivative works must be **licensed under the same license**.
- The full text of the license, unmodified invariant sections as defined by the author if any, and any other added warranty disclaimers (such as a general disclaimer alerting readers that the document may not be accurate for example) and copyright notices from previous versions must be maintained.
- Technical measures such as DRM may not be used to control or obstruct distribution or editing of the document.

# 1

## Introduction to Discrete Maths

### 1.1 Solved exercises

#### Problem 1

Let  $S = \{4n - 3 \mid n \in \mathbb{N}\}$  and let  $S_{\mathbb{P}} \subseteq S$  be the set of  $S$ -prime numbers, that being the numbers in  $S$  that have exactly two factors (1 and itself) in  $S$ .

1. Prove that  $S$  is closed under multiplication.
2. Are there infinitely many  $S$ -prime numbers?
3. Prove that  $1617 \in S$  and find two different factorizations of 1617 into  $S$ -primes.
4. Find a few more examples of  $S$ -integers with more than one factorization.

*Solution:*

First, we formally define  $S_{\mathbb{P}}$  as  $S_{\mathbb{P}} = \{x \in S \mid \nexists a, b \in S - \{1, x\} : x = ab\}$ . It's easy to notice that  $\mathbb{P} \cap S \subseteq S_{\mathbb{P}}$ , meaning that if a prime number is also in  $S$  then it's an  $S$ -prime number.

1. Given  $(4a - 3), (4b - 3) \in S$ , we show that:

$$\begin{aligned}(4a - 3)(4b - 3) &= 16ab - 12a - 12b + 9 = \\ 16ab - 12a - 12b + 12 - 3 &= 4(4ab - 3a - 3b + 4) - 3\end{aligned}$$

Since  $4ab - 3a - 3b + 4 \in \mathbb{N}$ , we conclude that  $(4a - 3)(4b - 3) \in S$ .

2. By way of contradiction, we suppose that  $S_{\mathbb{P}}$  is finite, meaning that  $S_{\mathbb{P}} = \{p_1, \dots, p_n\}$ .

Consider the number  $q := 4p_1 \dots p_n - 3$ . It's easy to see that  $q \in S - S_{\mathbb{P}}$ , meaning that  $q$  is  $S$ -composite and thus that  $\exists p_i, p_j \in S_{\mathbb{P}}$  such that  $p_i \mid q$  and  $p_j \mid q$ .

Without loss of generality, we proceed with  $p_i$ . By reflection, we have that  $p_i \mid p_i$ , which implies that  $p_i \mid 4p_1 \dots p_n$ . Then, since  $p_i \mid 4p_1 \dots p_n$  and  $p_i \mid q$ , it must also divide their difference, which equals 3, implying that  $p_i \mid 3$ .

Finally, since  $p_i \mid 3$ , it must hold that  $p_i \leq 3$ , implying that  $p_i \in \{1, 2, 3\}$ . However, if  $p_i = 2$  or  $p_i = 3$ , that would imply that  $2 \in S$  or  $3 \in S$ , which is a contradiction. By the same reasoning,  $p_i$  can't be equal to 1 since that would imply that  $p_j = 3$  and that  $3 \in S$ , which is a contradiction. Thus, the set  $S_{\mathbb{P}}$  must be infinite.

Another way to prove this result is by showing that  $\forall k \in \mathbb{N}$  it holds that  $4 \cdot 2^k - 3 \in S_{\mathbb{P}}$ . This can be easily done by way of contradiction. Moreover, this generator of infinite  $S$ -prime numbers can be extended to all primes, meaning that  $\forall p \in \mathbb{P}$  and  $\forall k \in \mathbb{N}$  it holds that  $4p^k - 3 \in S_{\mathbb{P}}$ .

3. It's easy to see that  $1617 = 4 \cdot 405 - 3$ , thus  $1617 \in S$ . We now consider the prime factorization  $1617 = 3 \cdot 11 \cdot 7^2$ , we notice that  $1617 = 33 \cdot 49$  and  $1617 = 21 \cdot 77$ .

Since  $33 = 4 \cdot 9 - 3$ , we get that  $33 \in S$ . However, since  $33 = 3 \cdot 11$  and  $3, 11 \notin S$ , the number 33 must be  $S$ -prime. By the same reasoning, we can show that  $49, 21, 77 \in S_{\mathbb{P}}$ , giving us two different  $S$ -prime factorizations of 1617.

4. Following the structure of the previous example, we can simply replace one of the numbers that form the prime factorization of 1617 with another prime number that isn't in  $S$ :

- The number  $441 = 3 \cdot 3 \cdot 7^2 \in S$  can be rewritten as  $441 = 9 \cdot 49 = 21 \cdot 21$ , where  $9, 21, 49 \in S_{\mathbb{P}}$ .
- The number  $2789 = 3 \cdot 19 \cdot 7^2 \in S$  can be rewritten as  $1029 = 57 \cdot 49 = 21 \cdot 133$ , where  $21, 57, 133 \in S_{\mathbb{P}}$ .