
a
Spearman ρ, all 8 tasks = −0.86 (exact p = 0.011)
7 QUBIQ tasks only = −0.79 (exact p = 0.048)
Evaluation-target (protocol) effect (Dice points, calibrated)
Rater agreement index |V ≥ N| / |V ≥ 1|
brain-tumour t2
LIDC-IDRI
brain-growth
prostate t2
brain-tumour t3
prostate t1
kidney
brain-tumour t1
b
Spearman ρ, all 8 tasks = −0.86 (exact p = 0.011)
7 QUBIQ tasks only = −1.00 (exact p = 0.0004)
Supervision-target effect (Dice points, calibrated)
Rater agreement index |V ≥ N| / |V ≥ 1|
QUBIQ, CT
QUBIQ, MRI
LIDC-IDRI (this study)
Marker size: rater count (3–7)
c
Union-trained (A)
Majority-trained (B)
Majority + soft (C)
brain-tumour t2 index 0.17
LIDC-IDRI index 0.33
brain-growth index 0.49
brain-tumour t3 index 0.64
prostate t2 index 0.77
brain-tumour t1 index 0.78
kidney index 0.87
prostate t1 index 0.96
Dice (points)
1/N
N/N
Evaluation reference V ≥ k as a fraction of raters, k/N (tasks ordered by agreement index)