



$$\neg \forall x \forall y [p(x,y)] \equiv \exists x \exists y [\neg p(x,y)] \quad \tanh(z) = -i \tan(iz)$$

How to solve the Arnold conjecture

By Ezra Guerrero
Mentor: Joye Chen



$$2ab + b^2$$



$$a_{1, n-1}$$


$$\sim(p \wedge q) \equiv \sim p \vee \sim q$$

$$\tanh(z) = -i \tan(iz)$$

$$S^2 = \sqrt{\sum_{i=1}^N (x_i - \bar{x})^2}$$

Table of contents



01 Hamiltonian Systems

We set up the problem and state the Arnold conjecture.

02 The Action Functional

We introduce the function at the center of our analysis.




03 Making a Chain Complex

We analyze the space of solutions and build a complex.

04 Floer Homology

We define Floer homology and present an example.


$$\text{scch}(z) = \text{ch}(z)$$




Symplectic Geometry

In classical mechanics, physicists solve problems by considering a *phase space* of possible states for a system.

In math, phase space becomes an even-dimensional manifold endowed with a *symplectic form*, ω . Such manifolds are called *symplectic manifolds*.




$$\omega_p(v, w) = \langle v \times w, n \rangle$$



01

Hamiltonian Systems

Counting periodic solutions of a differential system



$$\omega(\gamma, x_H) = dH(\gamma)$$




$$\sim \forall x \forall y [p(x,y)] \equiv \exists x \exists y [\sim p(x,y)]$$

$$\operatorname{sech}(z) = \sec(iz)$$

$$\frac{d}{dt} \left(\frac{1}{t} \right) = -\frac{1}{t^2}$$

Hamiltonians



$$H_t(x, y) = \frac{1}{2\pi} \left[\sin(2\pi y) + \cos(2\pi(x - t \cos(2\pi y))) \right]$$

A *time-dependent Hamiltonian* is a smooth function $H: W \times \mathbb{R} \rightarrow \mathbb{R}$. We write H_t for the Hamiltonian at time t , $H_t(p) = H(p, t)$.

Equipped with a symplectic form, we define the *symplectic gradient* of H_t by the formula:

$$\omega(Y, X_t) = dH_t(Y)$$


We are interested in solutions of period 1 to the *Hamiltonian system*

$$\dot{x}(t) = X_t(x(t))$$





The Arnold Conjecture



Conjecture 1: Let W be a compact symplectic manifold and let

$$H: W \times \mathbb{R} \rightarrow \mathbb{R}$$

be a time-dependent Hamiltonian. Then, there are at least

$$\sum_k \dim H_k(W; \mathbb{Z}/2)$$

periodic solutions to the associated Hamiltonian system.





$$\sim \forall x \forall y [p(x,y)] \equiv \exists x \exists y [\sim p(x,y)]$$

$$\operatorname{sech}(z) = \sec(iz)$$

$$\frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} f(x,y) \\ g(x,y) \end{pmatrix}$$

Autonomous Hamiltonians


Suppose H does not depend on time. We can prove the Arnold conjecture as follows:

Step 1: Find the critical points of H .

Step 2: Compute the *index of each critical point*.

Step 3: Count the number of trajectories connecting critical points.

Step 4: Build a chain complex whose homology coincides with the singular homology of W



$$H(x,y) = \cos(2\pi x) + \cos(2\pi y)$$

$$\begin{matrix} 0 & & 0 \\ \mathbb{Z}/2 & \xrightarrow{\quad} & \mathbb{Z}/2 \oplus \mathbb{Z}/2 & \xrightarrow{\quad} & \mathbb{Z}/2 \\ d & & b & c & a \end{matrix}$$

02

The Action Functional

Periodic solutions as critical points



$$\frac{\partial u}{\partial s} + J \frac{\partial u}{\partial t} + \text{grad } H_t = 0$$



The Loop Space



To follow our recipe, we need to work with the space of loops. We denote the space of contractible loops by $\mathcal{L}W$.

A tangent vector at $x \in \mathcal{L}W$ is given by a vector field along x .





The Action Functional

For each $x \in \mathcal{L}W$, pick an extension $u: D^2 \rightarrow W$.
We define


$$\mathcal{A}_H(x) = - \int_{D^2} u^* \omega + \int_0^1 H_t(x(t)) dt$$

This is well-defined under suitable assumptions. Thus, we have a function

$$\mathcal{A}_H: \mathcal{L}W \rightarrow \mathbb{R}$$

We can compute its derivative:

$$(d\mathcal{A}_H)_x(Y) = \int_0^1 \omega_{x(t)}(\dot{x}(t) - X_t(x), Y) dt$$





The Floer Equation

To define trajectories, we need a notion of gradient on our loop space.

We use an *almost complex structure* J , and compute

$$\operatorname{grad} \mathcal{A}_H = J\dot{x} + \operatorname{grad} H_t$$

Thus, the trajectories of the (negative) gradient are solutions $u: \mathbb{R}_s \times S^1_t \rightarrow W$ to the differential equation


$$\frac{\partial u}{\partial s} + J \frac{\partial u}{\partial t} + \operatorname{grad} H_t = 0$$



03

Making a Chain Complex

The space of trajectories and the differential





$$\neg \forall x \forall y [p(x,y)] \equiv \exists x \exists y [\neg p(x,y)]$$

$$\operatorname{sech}(z) = \sec(iz)$$


$$\frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} f(x,y) \\ g(x,y) \end{pmatrix}$$

The Space of Trajectories

Denote the space of trajectories from x to y by $\mathcal{L}(x, y)$. We can show that $\mathcal{L}(x, y)$ is a manifold of dimension $\mu(x) - \mu(y) - 1$

Thus, if $\mu(x) - \mu(y) = 1$, $\mathcal{L}(x, y)$ is a 0-dimensional manifold, and if $\mu(x) - \mu(z) = 2$, $\mathcal{L}(x, z)$ is a 1-dimensional manifold.

If we include *broken trajectories*, then $\overline{\mathcal{L}}(x, z)$ is compact, with boundary consisting of broken trajectories.





The Floer Complex

$$\dots \rightarrow CF_k(W) \xrightarrow{\partial} CF_{k-1}(W) \xrightarrow{\partial} \dots \xrightarrow{\partial} CF_0(W)$$

$$CF_k(W) = (\mathbb{Z}/2)^{\# \text{ crit. pts. of } \mu=k}$$

$$\partial(x) = \sum_{\mu(y)=\mu(x)-1} n(x,y) y$$



04

Floer Homology

A concrete computation

$$\# \text{ periodic solutions} \geq \sum_{k \geq 0} \dim HF_k(W; \mathbb{Z}/2)$$



$$\sim \forall x \forall y [p(x,y)] \equiv \exists x \exists y [\sim p(x,y)]$$

$$\operatorname{sech}(z) = \sec(iz)$$




Invariance of Floer Homology

We can prove that the homology of the Floer complex does not depend on the choice of H or J .

Picking a suitable pair (H, J) , we can show that this homology coincides with the singular homology of W .

$$\# \text{ periodic solutions} \geq \sum_{k \geq 0} \dim HF_k(W; \mathbb{Z}/2) = \sum_{k \geq 0} \dim H_k(W; \mathbb{Z}/2)$$






An Example

Consider the Hamiltonian

$$H_t(x, y) = \frac{1}{2\pi} [\sin(2\pi y) + \cos(2\pi(x - t \cos(2\pi y)))]$$

defined on the torus $T^2 = \mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z}$.

It has four contractible periodic solutions.





$$\neg \forall x \forall y [p(x,y)] \equiv \exists x \exists y [\neg p(x,y)]$$

$$\operatorname{sech}(z) = \sec(iz)$$

$$\frac{d}{dz} \left(\frac{1}{z} \right) = -\frac{1}{z^2}$$


Example (Continued)

According to the indices, the Floer complex must be

$$\mathbb{Z}/2 \longrightarrow \mathbb{Z}/2 \oplus \mathbb{Z}/2 \longrightarrow \mathbb{Z}/2$$



index = 1




index = 2

index = 0



index = 1



Thank you!



Questions?

CREDITS: This presentation template was created by [Slidesgo](#), and includes icons by [Flaticon](#), and infographics & images by [Freepik](#)

Followed Audin and Damian's *Morse Theory and Floer Homology*



$$S^2 = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}}$$



$$\neg \forall x \forall y [p(x, y)] \equiv \exists x \exists y [\neg p(x, y)] \quad \tanh(z) = -i \tanh(iz)$$

$$2ab + b^2$$

$$a_{1, n-1}$$