

## 2013 N8

Ezra Guerrero Alvarez

October 13, 2021

## 2003 N8

### 2003 N8

Let  $p$  be a prime number and let  $A$  be a set of positive integers that satisfies the following conditions:

- (i) the set of prime divisors of the elements in  $A$  consists of  $p - 1$  elements;
- (ii) for any nonempty subset of  $A$ , the product of its elements is not a perfect  $p$ -th power.

What is the largest possible number of elements in  $A$  ?

We claim the answer is  $(p - 1)^2$ . This is achievable by taking the set

$$A = \{q_i^{pj+1} | (i, j) \in \{1, \dots, p - 1\}^2\}$$

where  $\{q_i\}_{i=1}^\infty$  is the sequence of primes. Now we show  $|A| > (p - 1)^2$  is impossible. Let  $n = |A|$ . Note that by condition (i) we can represent the elements of  $A$  as vectors in  $\mathbb{F}_p^{p-1}$ . Let these be

$$v_i = (e_{i1}, \dots, e_{i(p-1)}) \text{ for } 1 \leq i \leq n.$$

Now, we define  $f_j: \mathbb{F}_p^n \rightarrow \mathbb{F}_p^n$  as

$$f_j(x_1, \dots, x_n) = \sum_{i=1}^n x_i^{p-1} e_{ji}.$$

Consider the system of equations  $f_j(x_1, \dots, x_n) = 0$  for  $1 \leq j \leq p - 1$ . Note that  $(0, \dots, 0)$  is a trivial solution. Suppose for the sake of contradiction that  $n > (p - 1)^2$ . Then, we have

$$n > (p - 1)^2 = (p - 1) \cdot (p - 1) = \sum_{j=1}^{p-1} \deg(f_j).$$

Thus, quoting *Chevalley's Theorem* there must exist a nontrivial solution to the system. However, since  $x_i^{p-1} \in 0, 1$ , this nontrivial solution specifies a nonempty subset of  $A$  for which the product is a perfect  $p$ -th power, yielding the desired contradiction. ■