

AI 1 Training

Contents

| | | |
|-----------|-------------------------------------|-----------|
| 1 | Linear Algebra | 5 |
| 2 | Introduction | 7 |
| 3 | Linear Systems | 9 |
| 4 | Matrix Representation | 11 |
| 4.1 | Determinant of a Matrix | 11 |
| 4.2 | Transpose of a Matrix | 11 |
| 4.3 | Inverse of a Matrix | 11 |
| 5 | Gaussian Elimination | 13 |
| 5.1 | Steps: | 13 |
| 5.2 | Example | 13 |
| 5.2.1 | Step 1: Augmented Matrix | 13 |
| 5.2.2 | Step 2: Row Reduction | 13 |
| 5.2.3 | Step 3: Back Substitution | 13 |
| 6 | Gauss–Jordan Elimination | 15 |
| 6.1 | Steps: | 15 |
| 6.2 | Resulting Form: | 15 |
| 7 | Row Echelon Form (REF) | 17 |
| 7.1 | Example | 17 |
| 7.2 | Advantages | 17 |
| 7.3 | Visual Representation | 17 |
| 7.4 | Solving with Row Echelon Form (REF) | 17 |
| 7.5 | Final Answer: | 18 |
| 8 | Vector | 19 |
| 9 | Vector Space | 21 |
| 10 | Linear Transformation | 23 |
| 11 | Introduction to SVMs | 25 |
| 12 | Basis | 27 |
| 13 | Eigenvalues and Eigenvectors | 29 |
| 14 | Calculus | 31 |

| | | |
|----|-----------------------------------------|----|
| 15 | Limit | 33 |
| 16 | Derivative | 35 |
| 17 | Partial Derivative | 37 |
| 18 | Gradient | 39 |
| 19 | Backpropagation | 41 |
| 20 | Newton, Hessian Method | 43 |
| 21 | Taylor Series, Taylor Theorem | 45 |
| 22 | Linearization | 47 |
| 23 | Optimization | 49 |
| 24 | Probability and Statistics | 51 |
| 25 | Probability | 53 |
| 26 | Probability Distributions | 55 |
| 27 | Sample | 57 |
| 28 | Estimating | 59 |
| 29 | Hypothesis Testing | 61 |

1. Linear Algebra

2. Introduction

Linear systems of equations form a core topic in linear algebra with widespread applications in mathematics, science, engineering, and technology. Solving linear systems efficiently is crucial, and among various techniques, *Gaussian elimination* is one of the most effective and widely used.

3. Linear Systems

A linear system of equations consists of multiple linear equations involving the same set of variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m \end{cases} \quad (3.1)$$

Where:

- a_{ij} are the coefficients,
- x_j are the unknowns,
- b_i are the constants.

4. Matrix Representation

A linear system can be written compactly in matrix form as:

$$A \cdot X = B \quad (4.1)$$

Where:

- A : $m \times n$ coefficient matrix
- X : $n \times 1$ unknown vector
- B : $m \times 1$ result vector

This representation is ideal for applying numerical methods.

4.1. Determinant of a Matrix

The determinant of a square matrix A , denoted $\det(A)$ or $|A|$, helps determine invertibility.

For a 2×2 matrix:

$$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc \quad (4.2)$$

4.2. Transpose of a Matrix

The transpose of a matrix A , denoted A^T , swaps its rows and columns.

Example:

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad (4.3)$$

$$A^T = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \quad (4.4)$$

4.3. Inverse of a Matrix

A square matrix A has an inverse A^{-1} if:

$$A \cdot A^{-1} = I \quad (4.5)$$

, where $\det(A) \neq 0$

Methods to compute A^{-1} :

- Augmented matrix (row operations)
- Adjoint method: $A^{-1} = \frac{1}{\det(A)} \cdot \text{adj}(A)$

5. Gaussian Elimination

A step-by-step method to reduce a matrix to upper triangular form:

5.1. Steps:

1. **Form the augmented matrix:** $[A \mid B]$
2. **Forward elimination:**
 - Identify pivot elements along the main diagonal
 - Use row operations to zero out entries below each pivot
3. **Back substitution:**
 - Solve for variables starting from the last row upward

5.2. Example

Given the system:

$$\begin{cases} 2x+3y-z=5 \\ 4x+y+2z=6 \\ -2x+5y+3z=7 \end{cases}$$

5.2.1. Step 1: Augmented Matrix

$$\left(\begin{array}{ccc|c} 2 & 3 & -1 & 5 \\ 4 & 1 & 2 & 6 \\ -2 & 5 & 3 & 7 \end{array} \right)$$

5.2.2. Step 2: Row Reduction

- Normalize the first row
- Eliminate entries below the pivot
- Repeat for next rows

5.2.3. Step 3: Back Substitution

Find values of z , then y , and finally x .

6. Gauss–Jordan Elimination

An extension of Gaussian Elimination that produces Reduced Row Echelon Form (RREF):

6.1. Steps:

- Form the augmented matrix $[A \mid B]$
- Use row operations to form leading 1s (pivots)
- Zero out **both** above and below the pivot in each column

6.2. Resulting Form:

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & x_1 \\ 0 & 1 & 0 & x_2 \\ 0 & 0 & 1 & x_3 \end{array} \right)$$

7. Row Echelon Form (REF)

A matrix is in *row echelon form* if:

- All nonzero rows are above zero rows
- The leading entry (pivot) in each nonzero row is 1
- Each pivot is to the right of the one in the row above
- All entries below a pivot are 0

7.1. Example

$$\left(\begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & 1 & 3 & -2 \\ 0 & 0 & 1 & 5 \end{array} \right)$$

7.2. Advantages

- Simplifies solving linear systems
- Easy to implement algorithmically
- Helps identify inconsistent or dependent systems

7.3. Visual Representation

$$\left(\begin{array}{cccc} 1 & & & \\ 0 & 1 & * & \\ 0 & 0 & 0 & 0 \end{array} \right)$$

Interpretation:

- Zero row at the bottom
- Leading 1s at $A_{\{11\}}$ and $A_{\{22\}}$
- All entries below pivots are 0

7.4. Solving with Row Echelon Form (REF)

We are given the system:

$$\begin{cases} x+y+z = 6 \\ 2x+3y+7z=20 \\ x+3y+4z = 13 \end{cases}$$

Step 1: Write the Augmented Matrix

$$\left(\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 2 & 3 & 7 & 20 \\ 1 & 3 & 4 & 13 \end{array} \right)$$

Step 2: Row Reduction to Row Echelon Form

- Keep Row 1 as is.
- Eliminate below the first pivot (Row 1, Col 1):

$$R_2 := R_2 - 2 \cdot R_1$$

$$R_3 := R_3 - R_1$$

New matrix:

$$\left(\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 1 & 5 & 8 \\ 0 & 2 & 3 & 7 \end{array} \right)$$

- Eliminate below the second pivot (Row 2, Col 2):

$$R_3 := R_3 - 2 \cdot R_2$$

Resulting Row Echelon Form:

$$\left(\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 1 & 5 & 8 \\ 0 & 0 & -7 & -9 \end{array}\right)$$

Step 3: Back Substitution

From the last row:

$$-7z = -9 \rightarrow z = \frac{9}{7} \quad (7.1)$$

Second row:

$$y + 5z = 8 \rightarrow y = 8 - 5 \cdot \left(\frac{9}{7}\right) = \frac{11}{7} \quad (7.2)$$

First row:

$$x + y + z = 6 \rightarrow x = 6 - \frac{11}{7} - \frac{9}{7} = \frac{22}{7} \quad (7.3)$$

7.5. Final Answer:

$$x = \frac{22}{7}, y = \frac{11}{7}, z = \frac{9}{7} \quad (7.4)$$

8. Vector

9. Vector Space

10. Linear Transformation

11. Introduction to SVMs

12. Basis

13. Eigenvalues and Eigenvectors

14. Calculus

15. Limit

16. Derivative

17. Partial Derivative

18. Gradient

19. Backpropagation

20. Newton, Hessian Method

21. Taylor Series, Taylor Theorem

22. Linearization

23. Optimization

24. Probability and Statistics

25. Probability

26. Probability Distributions

27. Sample

28. Estimating

29. Hypothesis Testing