

MX4557 Complex Analysis

F. Olukoya




Table of contents

Revision Guide	5
How to Prepare	5
Important statements and results	6
Use Of Calculators	11
 I Complex numbers and functions	 12
1 Geometry of the Complex plane	13
Questions for further discussion	21
1.1 Exercises	22
Further resources	28
 2 Complex-valued functions	 29
Image grid	33
Domain colouring	36
3-dimensional graphs	38
Questions for further discussion	38
2.1 Exercises	39
Further resources	41
 3 Topology of the complex plane	 42
Questions for further discussion	46
3.1 Exercises	47
Further resources	49
 II Differentiation and contour integrals	 50
 4 Holomorphic functions	 51

Questions for further discussion	59
4.1 Exercises	60
5 Power Series	63
Questions for further discussion	73
5.1 Exercises	74
6 Contour integrals	78
Questions for further discussion	88
6.1 Exercises	88
 III Cauchy's theorem and residues	 92
7 Cauchy's theorem	93
Questions for further discussion	101
7.1 Exercises	102
8 Applications of Cauchy's theorem	105
Questions for further discussion	114
8.1 Exercises	114
9 Laurent series and singularities	117
Questions for further discussion	126
9.1 Exercises	126
10 Residue theorem	129
10.1 Exercises	136
11 Applications to integrals	138
 Appendices	 142
A All Solutions	143
A.1 Chapter 1 solutions	143
A.2 Chapter 2 solutions	169
A.3 Chapter 3 solutions	178
A.4 Chapter 4 solutions	184
A.5 Chapter 5 solutions	192
A.6 Chapter 6 solutions	201
A.7 Chapter 7 solutions	207

A.8 Chapter 8 solutions	214
A.9 Chapter 9 solutions	218
References	225

Revision Guide

This guide offers some advice on preparing for the exam.

! Rule 1

You cannot begin revising too soon!

! Rule 2

Read the questions *carefully*.

! Rule 3

Learn from mistakes: revisit both the generic feedback and personalised annotations on your assessments to help avoid repeating similar errors.

How to Prepare

Go through your notes and ensure you understand:

- Key definitions
- Theorems and their statements

- Where possible, the proofs (or outlines) of important results

Work through:

- Quiz questions and their solutions
- Continuous assessments (feel free to arrange a meeting if there are parts you'd like to go over)
- Past exams and solutions
- Problem sheets and solutions

The exam may include:

- Recalling definitions and key results
- Stating or interpreting parts of proofs
- Solving problems similar to those in quizzes, problem sheets, and assessments

By engaging with all of the above, you'll be well prepared. You're also welcome to reach out by [email](#) or arrange a meeting if you'd like help with anything.

Important statements and results

Below is a *non-exhaustive* list of key results and concepts you should know, grouped by chapter.

Chapter 1

- Expressing complex numbers in polar Proposition 1.5 and exponential form (see the discussion following Definition 1.4)
- Computing the argument and principal argument Definition 1.3
- Finding all n th roots of a complex number Proposition 1.7
- Visualising complex numbers in the complex plane

Chapter 2

- Definition of a complex function Definition 2.1, especially polynomials and rational functions Example 2.2
- Identifying the domain of definition
- Real and imaginary parts of a complex function
- Definitions of $\exp$, $\sin$, $\cos$, $\sinh$, $\cosh$ via exponentials Example 2.4 and Example 2.3
- Principal branch of the logarithm Example 2.5
- Understanding how standard complex functions transform geometric sets (e.g. lines, circles)

Chapter 3

- Definitions of open discs Definition 3.1, open/closed/bounded sets Definition 3.2
- Convergence of sequences Definition 3.3 and series Definition 3.4
- Cauchy sequences and their equivalence with convergence in $\mathbb{C}$ Proposition 3.1
- Closure of sets Definition 3.5
- Limits and continuity for complex functions (Definition 3.5)

Chapter 4

- Complex differentiability Definition 4.1
- Linearity, product, quotient, and chain rules Proposition 4.1
- The Cauchy-Riemann equations and their use Proposition 4.3 and the examples that follow
- Derivative of the complex logarithm Example 4.6

Chapter 5

- Addition, multiplication, and inversion of power series Definition 5.2
- Radius of convergence using standard convergence tests

- Taylor series for $\frac{1}{1-z}$, $\exp$, $\cos$, $\sin$ Example [5.2](#)
- Absolute and uniform convergence within the radius Theorem [5.1](#)
- Power series define holomorphic functions Theorem [5.2](#)

Chapter 6

- Integral of a complex-valued function over a real interval Definition [6.1](#)
- Estimating integrals via modulus Proposition [6.1](#)
- Definitions and properties of curves, paths, and loops Definition [6.2](#)
- Parametrising standard curves
- Definition and properties of contour integrals Definition [6.3](#)
- Invariance under reparametrisation Lemma [6.1](#)
- Concatenation and reversal of curves Definition [6.4](#)
- Complex version of the Fundamental Theorem of Calculus Proposition [6.3](#)

Chapter 7

- Definitions of homotopy, path homotopy, and null-homotopy Definition [7.1](#)
- Cauchy's Theorem and its application Theorem [7.1](#)

Chapter 8

- Cauchy's integral formulae Theorem [8.1](#)
- Consequences such as Liouville's Theorem and the Fundamental Theorem of Algebra Theorem [8.3](#)
- Existence of primitives for entire functions and in simply connected regions Theorem [8.2](#)

Chapter 9

- Definition of bilateral series and their residues Definition [9.1](#)
- Laurent series and convergence within an annulus Proposition [9.1](#)
- Expansion of holomorphic functions as Laurent series Theorem [9.1](#) using the geometrix series
- Classification of isolated singularities Definition [9.4](#)

Chapter 10

- Residues of holomorphic functions Definition [10.1](#)
- Computing residues Proposition [10.1](#)
- Winding number of a loop Definition [10.2](#)
- Residue Theorem Theorem [10.1](#)

Chapter 11

- Applications of the Residue Theorem to compute real integrals

Use Of Calculators

Only silent, self-contained calculators with a single-line display are permitted in the examination. Calculator memories must be clear at the start of the examination. The type of calculator used must be specified on the cover sheet of your answer book.

Part I

Complex numbers and functions

Chapter 1

Geometry of the Complex plane

This section is a brief reminder of Sections 3 and 4 of MA1006 Algebra.

Definition 1.1. The *complex numbers* $\mathbb{C}$ are the set of all pairs $z = (x, y) \in \mathbb{R}^2$ of real numbers with the addition

$$z_1 + z_2 = (x_1 + x_2, y_1 + y_2) \quad (1.1)$$

and the multiplication

$$z_1 \cdot z_2 = (x_1x_2 - y_1y_2, x_1y_2 + x_2y_1), \quad (1.2)$$

where $z_1 = (x_1, y_1)$ and $z_2 = (x_2, y_2)$. We call $x = \Re(z)$ the *real part* and $y = \Im(z)$ the *imaginary part* of z .

From Section 3.8 in MA1006 Algebra we recall:

Proposition 1.1. *The complex numbers are a field.*

Remark 1.1.

We view the real numbers as a subset of $\mathbb{C}$ by identifying $x \in \mathbb{R}$ with $(x, 0) \in \mathbb{C}$. The **imaginary unit** is $i = (0, 1)$. With these conventions, a calculation using Equation 1.2 shows that

$$z = x + iy. \quad (1.3)$$

Using this notation, we can manipulate complex numbers in the same way as real numbers, keeping in mind the identity

$$i^2 = -1. \quad (1.4)$$

Of course, $\mathbb{C}$ is a one-dimensional vector space over itself. Restricting the scalar multiplication to $\mathbb{R}$ makes $\mathbb{C}$ into a vector space over $\mathbb{R}$, isomorphic to $\mathbb{R}^2$, of dimension two with standard basis $1, i \in \mathbb{C}$.

Proposition 1.2.

a. In the standard basis, every $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_{2 \times 2}(\mathbb{R})$ corresponds uniquely to an $\mathbb{R}$ -linear map $T_A: \mathbb{C} \rightarrow \mathbb{C}$, namely

$$T_A(x + iy) = (ax + by) + i(cx + dy). \quad (1.5)$$

b. T_A is $\mathbb{C}$ -linear $\iff a = d$ and $b = -c$. In this case,

$$T_A(z) = \alpha \cdot z, \quad \alpha = a + ic. \quad (1.6)$$

Proof.

(a) This is a recap from linear algebra. An $\mathbb{R}$ -linear map $T: \mathbb{C} \rightarrow \mathbb{C}$ is uniquely determined by the image of the basis vectors $T(1), T(i)$ which, conversely, may be prescribed arbitrarily. If we write $T(1) = a+ic, T(i) = b+id$, then the linear transform T is described entirely by $a, b, c, d \in \mathbb{R}$. The expression Equation 1.5 is obtained by expanding the left hand side by $\mathbb{R}$ -linearity.

(b) In the same way, $\mathbb{C}$ -linear maps correspond to $\alpha \in M_{1 \times 1}(\mathbb{C})$ as in Equation 1.6.

An $\mathbb{R}$ -linear map T_A is $\mathbb{C}$ -linear $\iff T_A(i) = T_A(i1) = iT_A(1) \iff$

$$\begin{pmatrix} b \\ d \end{pmatrix} = T_A(i) = iT_A(1) = i \begin{pmatrix} a \\ c \end{pmatrix} = \begin{pmatrix} -c \\ a \end{pmatrix}.$$

In this case, $T_A(x + iy) = (ax - cy) + i(cx + ay) = \alpha(x + iy)$.

□

Definition 1.2. The *conjugate* of $z \in \mathbb{C}$ is the complex number

$$\bar{z} = (x, -y),$$

and the *modulus* (also called *absolute value*) is

$$|z| = \sqrt{x^2 + y^2} \geq 0.$$

Proposition 1.3. The following formulas hold for $z, w \in \mathbb{C}$:

$$\overline{z \cdot w} = \bar{z} \cdot \bar{w} \qquad \overline{z + w} = \bar{z} + \bar{w} \qquad (1.7)$$

$$\overline{\bar{z}} = z \qquad \bar{i} = -i, \bar{1} = 1 \qquad (1.8)$$

$$z \cdot \bar{z} = |z|^2 \qquad |z \cdot w| = |z| \cdot |w| \qquad (1.9)$$

$$\Re(z) = \frac{z + \bar{z}}{2} \qquad \Im(z) = \frac{z - \bar{z}}{2i} \qquad (1.10)$$

$$z^{-1} = \frac{\bar{z}}{|z|^2} \quad \text{if } z \neq 0 \qquad (1.11)$$

Proposition 1.4. *The following inequalities hold for $z, w \in \mathbb{C}$:*

$$|z + w| \leq |z| + |w| \qquad |z - w| \geq \big||z| - |w|\big| \qquad (1.12)$$

$$|\Re(z)| \leq |z| \qquad |\Im(z)| \leq |z| \qquad (1.13)$$

Thinking of complex numbers as points in the plane, we can use **polar coordinates** to represent them (see Figure 1.1).




Figure 1.1: Polar coordinates

Proposition 1.5. *For every non-zero complex number $z = (x, y)$ there is an **argument** $\theta \in \mathbb{R}$ and a **radius** $r = |z| > 0$ such **that***

$$x = r \cos(\theta), y = r \sin(\theta). \quad (1.14)$$

This representation is unique up to replacing θ by $\theta + 2\pi k$ for any $k \in \mathbb{Z}$.

Proof. (omitted)

Since $x^2 + y^2 = r^2(\cos(\theta)^2 + \sin(\theta)^2) = r^2$, the radius must be $r = |z|$. Define the complex number $w = r^{-1}z$ and write $w = u + iv$ for its real and imaginary parts.

We will prove the existence of $\theta \in \mathbb{R}$ with $u = \cos(\theta)$, $v = \sin(\theta)$. This also proves the existence of a representation Equation 1.14, by multiplying by r . Since $u^2 + v^2 =$

$|w|^2 = r^{-2}|z|^2 = 1$, we know $|u| \leq 1$, $|v| \leq 1$. Recall that $\cos: [0, \pi] \rightarrow [-1, 1]$ and $\sin: [-\pi/2, \pi/2] \rightarrow [-1, 1]$ are bijections. Hence

$$\begin{aligned} u &= \cos(\alpha) && \text{for some } \alpha \in [0, \pi], \\ v &= \sin(\beta) && \text{for some } \beta \in [-\pi/2, \pi/2]. \end{aligned}$$

As $\sin(\beta)^2 = v^2 = 1 - u^2 = 1 - \cos(\alpha)^2 = \sin(\alpha)^2$, we have $\sin(\alpha) = \pm \sin(\beta) = \sin(\pm\beta)$. To produce the correct θ , we distinguish two cases.

Case 1 $\alpha \in [0, \pi/2]$. Then $\alpha = \pm\beta$ by the injectivity of the sine function on the interval $[-\pi/2, \pi/2]$. Setting $\theta = \pm\alpha = \beta$, we find that $u = \cos(\theta)$ and $v = \sin(\theta)$, as required.

Case 2 $\alpha \in [\pi/2, \pi]$. Then $\pi - \alpha, \beta \in [-\pi/2, \pi/2]$ and $\sin(\pi - \alpha) = \sin(\alpha) = \pm \sin(\beta)$, so $\pi - \alpha = \pm\beta$ by injectivity. Setting $\theta = \pm\alpha = \pm\pi - \beta$, we find $u = \cos(\theta), v = \sin(\theta)$, using trigonometric identities.

This completes the existence part of the proof. For uniqueness, we already know that $r = |z| > 0$ is unique, so it remains to consider

$$x = r \cos(\theta_1) = r \cos(\theta_2), \quad y = r \sin(\theta_1) = r \sin(\theta_2).$$

To translate the situation into an interval that we understand, pick $k_1, k_2 \in \mathbb{Z}$ so that $\theta_1 + 2\pi k_1, \theta_2 + 2\pi k_2 \in [-\pi, \pi)$. Then

$$\cos(\theta_1 + 2\pi k_1) = \cos(\theta_1) = \cos(\theta_2) = \cos(\theta_2 + 2\pi k_2).$$

Using the injectivity of the cosine function and considering cases as above, we find that $\theta_1 + 2\pi k_1 = \pm(\theta_2 + 2\pi k_2)$. If the sign is '+' we get $\theta_1 - \theta_2 = 2\pi(k_2 - k_1)$

and we are done, so suppose $\theta_1 + 2\pi k_1 = -(\theta_2 + 2\pi k_2)$. Then

$$\sin(\theta_1) = \sin(\theta_2) = \sin(\theta_2 + 2\pi k_1) = -\sin(\theta_1 + 2\pi k_1) = -\sin(\theta_1)$$

implies $\sin(\theta_1) = 0$. Therefore θ_1 is a multiple of 2π , which implies that $\theta_1 + 2\pi k_1 = -(\theta_2 + 2\pi k_2) = 0$ since these numbers were chosen in $[-\pi, \pi)$ and we have $2\pi\mathbb{Z} \cap [-\pi, \pi) = \{0\}$. Hence $\theta_1 - \theta_2 = 2\pi(k_2 - k_1)$. $\square$

To get around the non-uniqueness of the argument in polar coordinates, we restrict θ to lie in a half-open interval of length 2π . Here is the most common convention.

Definition 1.3. The **principal argument** of a non-zero $z \in \mathbb{C}$ is the unique $\theta \in (-\pi, \pi]$ such that Equation 1.14 holds, and we write $\arg(z) = \theta$.

Definition 1.4. The value of the **exponential function** at the complex number $z = x + i\theta$, where $x, \theta \in \mathbb{R}$, is defined as

$$e^{x+i\theta} = e^x(\cos(\theta) + i\sin(\theta)). \quad (1.15)$$

Proposition 1.5 implies that every complex number can be represented in **polar form**

$$z = re^{i\theta}. \quad (1.16)$$

The addition of complex numbers is the usual addition of vectors in $\mathbb{R}^2$. To visualize multiplication, the polar form is useful. Combining Equation 1.2 and Equation 1.15,

we find that

$$\begin{aligned} e^{i\theta_1} \cdot e^{i\theta_2} &= \cos(\theta_1) \cos(\theta_2) - \sin(\theta_1) \sin(\theta_2) \\ &\quad + i[\cos(\theta_1) \sin(\theta_2) + \cos(\theta_2) \sin(\theta_1)] \\ &= \cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2) = e^{i(\theta_1 + \theta_2)}. \end{aligned}$$

Here we have used the trigonometric addition formulas. Hence

$$z_1 = r_1 e^{i\theta_1}, z_2 = r_2 e^{i\theta_2} \implies z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}. \quad (1.17)$$

Complex multiplication adds the angles and multiplies the radii.

Proposition 1.6. $e^{z_1} \cdot e^{z_2} = e^{z_1 + z_2}$ for all $z_1, z_2 \in \mathbb{C}$. Moreover, we have $(e^z)^n = e^{nz}$ for all $z \in \mathbb{C}$, $n \in \mathbb{Z}$.

Proof.

The first assertion follows from Equation 1.17 combined with the identity $e^{x_1} e^{x_2} = e^{x_1 + x_2}$ for $x_1, x_2 \in \mathbb{R}$ from MA1005 Calculus. The second claim follows from this by induction.

□

The polar form can be applied to the construction of n^{th} roots.

For example, the **n^{th} root of unity** is $\zeta_n = e^{i\frac{2\pi}{n}}$ and satisfies

$$(\zeta_n)^n = (e^{i\frac{2\pi}{n}})^n = e^{2\pi i} = 1.$$

Proposition 1.7. Every complex number $z \neq 0$ has an n^{th} root w satisfying $w^n = z$.

If w is an n^{th} root of z , the set of all n^{th} roots of z is

$$\{w, \zeta_n \cdot w, \zeta_n^2 \cdot w, \dots, \zeta_n^{n-1} \cdot w\}.$$

Proof.

Write $z = re^{i\theta}$ and $w = se^{i\varphi}$ for $\theta, \varphi \in [0, 2\pi)$ and $r, s > 0$. By the uniqueness of the polar form, the equation $w^n = z$ is equivalent to $s^n = r$ and $n\varphi = \theta + 2\pi k$ for some $k \in \mathbb{Z}$. Of course, $s = \sqrt[n]{r}$ is the unique positive n^{th} root from MA1005 Calculus. From $0 \leq n\varphi < 2\pi n$ and $0 < -\theta \leq 2\pi$ we get

$$0 < k = \frac{n\varphi - \theta}{2\pi} < n.$$

Since k is an integer, this implies $k = 0, \dots, n-1$ and hence $\varphi = \frac{\theta + 2\pi k}{n}$ for such k are the only possible solutions for φ . In summary,

$$w_k = \sqrt[n]{r}e^{i\varphi} = \sqrt[n]{r}e^{i\frac{\theta + 2\pi k}{n}} = \zeta_n^k \cdot w_0, k = 0, \dots, n-1,$$

are all the possible n^{th} roots of z .

□

Questions for further discussion

- The complex numbers are obtained by ‘adjoining’ a symbol i with $i^2 = -1$. If instead we would have adjoined a different symbol ϵ with $\epsilon^2 = -1$, would the set of elements $x + \epsilon y$ still define a field?
- The real numbers have a total order ‘ $\leq$ ’. Why doesn’t it make sense to extend this definition to the complex numbers?

- Describe geometrically the set $R_n = \{1, \zeta_n, \dots, (\zeta_n)^{n-1}\}$ of n^{th} roots of unity. Find a connection between R_n and the cyclic group $C_n = \{\overline{0}, \dots, \overline{n-1}\}$ of integers modulo n from MX3020 Group Theory.

1.1 Exercises

Note

This problem sheet is intended as a recap and contains more problems than can be discussed during the tutorials.

Exercise 1.1

Verify

$$z = x + iy$$

and

$$i^2 = -1$$

straight from the definition Equation 1.2.

Show Solution 1.1 on P143

Exercise 1.2

How many real solutions x does $x^2 + 1 = 0$ have? Show that the polynomial equation $z^2 + 1 = 0$ has exactly two solutions $z \in \mathbb{C}$.

Show Solution 1.2 on P144

Exercise 1.3

Give examples of complex numbers $z, w \neq 0$ such that $z^2 + w^2 = 0$.

Show Exercise 1.3 on P145

Exercise 1.4

Sketch the position of the complex numbers $i, 1 + i, \frac{3+2i}{4}$ in the plane.

Show Solution 1.4 on P146

Exercise 1.5

Express the following complex numbers z in the form $x + iy$ with $x, y \in \mathbb{R}$.

$$(1 + i)^{20}, (5 + 3i)(1 + 2i), (1 - i)(2 + 3i), (1 - i)i(1 + i), \frac{2 + i}{1 - i}$$

Show Solution 1.5 on P146

Exercise 1.6

Express the following complex numbers z in the form $x + iy$ with $x, y \in \mathbb{R}$.

$$1/i, \frac{1}{1 + i}, \frac{3 + i}{3 - i}$$

Show Solution 1.6 on P147

Exercise 1.7

Find the modulus and the conjugate of the following complex numbers.

$$2 + i, i, 5 - 3i, \frac{1 + i}{2 + i}$$

Show Solution 1.7 on P147

Exercise 1.8

Describe the sets $A = \{z \in \mathbb{C} \mid \Im(z) > 0\}$, $B = \{z \in \mathbb{C} \mid \Re(z) \leq 1\}$, $C = \{z \in \mathbb{C} \mid \Re((1+i)z) = 0\}$, and $A \cap B$ geometrically.

Show Solution 1.8 on P148

Exercise 1.9

Describe the set $D = \{z \in \mathbb{C} \mid z \cdot \bar{z} = 1\}$ geometrically.

Hint: Write $z = re^{i\theta}$ in polar form.

Show Solution 1.9 on P151

Exercise 1.10

Draw all nine sets described by the following conditions on the complex number z .

$$\begin{array}{lll} |z| = 1, & |z| < 1, & 1 < |z| < 2, \\ |1+z| > 1, & |2-z| < 2, & 3 < |z+i| < 4, \\ |z-1| < |z+1|, & |z| = |z+1|, & |z-1| = |z+i|. \end{array}$$

Show Solution 1.10 on P152

Exercise 1.11

Let $S = \{x + iy \in \mathbb{C} \mid 0 \leq x, y \leq 1\}$. Draw S and the sets

$$\begin{array}{ll} A = \{2z \mid z \in S\}, & B = \{\bar{z} \mid z \in S\}, \\ C = \{-z \mid z \in S\}, & D = \{z^2 \mid z \in S\}. \end{array}$$

Show Solution 1.11 on P157

Exercise 1.12

Let $D = \{z \in \mathbb{C} \mid |z| < 1\}$ be the unit disk. Draw the sets

$$A = \{2z \mid z \in D\}, \quad B = \{z^2 \mid z \in D\}, \quad C = \{|z| \mid z \in D\}.$$

Show Solution 1.12 on P159

Exercise 1.13

Show that $i = e^{i\pi/2}$ and $-1 = e^{i\pi}$.

Show Solution 1.13 on P159

Exercise 1.14

Express the following complex numbers z in the form $x + iy$ with $x, y \in \mathbb{R}$.

$$e^{i\pi/4}, \quad e^{i\pi}, \quad e^{i\frac{2\pi}{3}}$$

Show Solution 1.14 on P159

Exercise 1.15

Write each of the following complex numbers in polar form $re^{i\theta}$ with $r > 0$ and $-\pi < \theta \leq \pi$.

$$i, \quad -1, \quad -i, \quad 1+i, \quad 1-i, \quad i-1, \quad \frac{1}{2} + i\frac{\sqrt{3}}{2}$$

Draw each of these numbers in the complex plane.

Show Solution 1.15 on P160

Exercise 1.16

Calculate i^{2021} and $(1 + i)^{20}$.

Show Solution 1.16 on P161

Exercise 1.17

Solve the equation $(1 - i)^n - 2075 = 2021$ and find $n \in \mathbb{N}$.

Show Solution 1.17 on P162

Exercise 1.18

Prove that for $z \in \mathbb{R}$ we have

$$\cos(z) = \frac{e^{iz} + e^{-iz}}{2}, \quad \sin(z) = \frac{e^{iz} - e^{-iz}}{2i}.$$

Use these equations to extend the definition of the functions $\cos(z)$, $\sin(z)$ to complex arguments $z \in \mathbb{C}$. Find $z \in \mathbb{C}$ with $\sin(z) = 2$.

Hint: Put $w = e^{iz}$ and reduce to a quadratic equation.

Show Solution 1.18 on P162

Exercise 1.19

Prove the following statements.

- a. $\overline{zw} = \bar{z} \cdot \bar{w}$ for all $z, w \in \mathbb{C}$
- b. $\overline{z_1 z_2 \cdots z_n} = \bar{z}_1 \bar{z}_2 \cdots \bar{z}_n$ for all $z_1, z_2, \dots, z_n \in \mathbb{C}$ (use induction)
- c. $\overline{(z^n)} = (\bar{z})^n$ for all $z \in \mathbb{C}$
- d. Let $p(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0$ be a polynomial with real coefficients $a_0, \dots, a_n \in \mathbb{R}$. Prove that $\overline{p(z)} = p(\bar{z})$. Deduce that all roots of $p(z)$ occur in complex conjugate pairs.

Show Solution 1.19 on P164

Exercise 1.20

Show that $|z| = |-z|$ and $|\bar{z}| = |z|$. Prove also that $|\lambda z| = \lambda |z|$ for all $\lambda \geq 0$.

Show Solution 1.20 on P165

Exercise 1.21

Prove that $\Re(z) = \frac{1}{2}(z + \bar{z})$, $\Im(z) = \frac{1}{2i}(z - \bar{z})$.

Show Solution 1.21 on P165

Exercise 1.22

Prove that $\overline{e^z} = e^{\bar{z}}$. Deduce that $|e^z| = e^{\Re(z)}$.

Show Solution 1.22 on P166

Exercise 1.23

Prove that $|z + w|^2 + |z - w|^2 = 2|z|^2 + 2|w|^2$.

Show Solution 1.23 on P166

Exercise 1.24

Show that $|z+w|^2 = |z|^2 + 2\Re(z\bar{w}) + |w|^2$. Use this to determine the conditions on z, w for $|z+w| = |z| + |w|$ to hold.

Show Solution 1.24 on P167

Exercise 1.25

Assuming we know the triangle inequality $|z+w| \leq |z| + |w|$ for all $z, w \in \mathbb{C}$, prove the reverse triangle inequality

$$|z-w| \geq ||z| - |w||.$$

Show Solution 1.25 on P168

Exercise 1.26

Let K be a field with $\mathbb{R} \subset K \subset \mathbb{C}$. Prove that $K = \mathbb{R}$ or $K = \mathbb{C}$.

Show Solution 1.26 on P169

Further resources

- Freitag–Busam ([Freitag and Busam 2009, chap. 1](#)) for additional exercises and historical background.
- https://en.wikipedia.org/wiki/Complex_number for overview and history
- <https://youtu.be/T647CGsuOVU> for some visualization

Chapter 2

Complex-valued functions

Definition 2.1. A complex function $f: D \rightarrow \mathbb{C}$ is a map with *domain of definition* $D \subset \mathbb{C}$ and codomain the complex plane. Thus, f assigns to each $z = x + iy \in D$ in the domain a complex number

$$f(z) = u(z) + iv(z).$$

We call $u: D \rightarrow \mathbb{R}$ the *real part* and $v: D \rightarrow \mathbb{R}$ the *imaginary part* of the complex function f .




Figure 2.1: Schematic picture of a complex function

We can visualize complex functions as in Figure 2.1. In practice, most complex functions are defined by a formula.

Example 2.1.

Let $a, b \in \mathbb{C}$ be fixed complex numbers. Then $f(z) = az + b$ is a complex function with domain $D = \mathbb{C}$, written $f: \mathbb{C} \rightarrow \mathbb{C}$. For instance,

$$f(z) = iz, f(z) = 2z, f(z) = z + 3.$$

Example 2.2.

A *polynomial* is a complex function $f: \mathbb{C} \rightarrow \mathbb{C}$ of the form

$$f(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0, \quad (2.1)$$

with complex *coefficients* $a_n, \dots, a_0 \in \mathbb{C}$ and with domain $D = \mathbb{C}$.

More generally, a *rational function* is a complex function for which there are polynomials f, g with $g \neq 0$ so that

$$h(z) = \frac{f(z)}{g(z)}. \quad (2.2)$$

The domain of h is $D = \{z \in \mathbb{C} \mid g(z) \neq 0\}$. However, if the polynomials f, g have a common factor, we can cancel that factor in the fraction Equation 2.2 and regard h as a complex function on a larger domain.

For example, $h(z) = \frac{z+1}{z^2-1}$ has domain $\mathbb{C} \setminus \{\pm 1\}$, but we can rewrite the fraction as $h(z) = \frac{1}{z-1}$, which makes sense on the extended domain $\mathbb{C} \setminus \{+1\}$.

We have already met several complex functions in the previous section.

Example 2.3.

The *exponential function* $\exp: \mathbb{C} \rightarrow \mathbb{C}$ is defined by $\exp(z) = e^z = e^x(\cos(y) + i \sin(y))$ and has domain $D = \mathbb{C}$.

Example 2.4.

The *sine function* $\sin: \mathbb{C} \rightarrow \mathbb{C}$ and the *cosine function* $\cos: \mathbb{C} \rightarrow \mathbb{C}$ are the complex functions defined by

$$\cos(z) = \frac{e^{iz} + e^{-iz}}{2}, \quad \sin(z) = \frac{e^{iz} - e^{-iz}}{2i}. \quad (2.3)$$

(See the corresponding [exercise](#) in Section 1.1)

Definition 2.2. We define the following subsets of the complex plane:

$$\mathbb{C}^\times = \{z \in \mathbb{C} \mid z \neq 0\}$$

punctured plane

$$\mathbb{C}^- = \mathbb{C} \setminus \{x \in \mathbb{R} \mid x \leq 0\}$$

slit plane

$$S = \{z = x + iy \mid y \in (-\pi, \pi)\}$$

principal strip

$$\mathbb{H} = \{z = x + iy \mid y > 0\}$$

upper half-plane




Figure 2.2: The principal strip S and the slit plane $\mathbb{C}^-$

Example 2.5.

The *principal branch* of the *complex logarithm* is the complex function

$$\log: \mathbb{C}^- \longrightarrow S \subset \mathbb{C}$$

defined by

$$\log(w) = \log(r) + i\theta \iff w = re^{i\theta}, r > 0, \theta \in (-\pi, \pi). \quad (2.4)$$

Here $\log(r)$ denotes the (real) logarithm from MA1005 Calculus. In other words, the real part of $\log(w)$ is the logarithm of the modulus $r = |w|$ and the imaginary part of $\log(w)$ is the argument function $\arg(w)$ from Definition 1.3.

To evaluate Equation 2.4, write w in polar coordinates, ensuring that $\theta \in (-\pi, \pi)$. For example, $i = e^{i\pi/2}$ and so $\log(i) = \log(1) + i\frac{\pi}{2} = i\frac{\pi}{2}$.

Proposition 2.1.

- a. The exponential function is surjective onto $\mathbb{C}^\times$.
- b. We have

$$\exp(\log(w)) = w \quad (\forall w \in \mathbb{C}^-) \quad (2.5)$$

$$\log(\exp(z)) = z \quad (\forall z \in S). \quad (2.6)$$

Hence the restriction $\exp|_S$ of the exponential to the principal strip is a bijection $\exp|_S: S \rightarrow \mathbb{C}^-$ onto the slit plane, with inverse $\log(w)$.

Proof.

- a. Recall from MA1005 Calculus that $e^x \neq 0$ for all $x \in \mathbb{R}$. As $|\exp(x + iy)| = e^x \neq 0$, the image of $\exp$ is contained in $\mathbb{C}^\times$. For proving that $\exp$ is onto $\mathbb{C}^\times$, recall that every non-zero number can be written in polar form $w = re^{i\theta}$, $\theta \in (-\pi, \pi]$. Then $\exp(z) = w$ for $z = \log(r) + i\theta$.

- b. Equations Equation 2.5, Equation 2.6 are straightforward to verify using Equation 2.4.

□

Since the *graph* $\Gamma(f) = \{(z, w) \in D \times \mathbb{C} \mid f(z) = w\}$ of a complex function is a subset of four-dimensional space, we cannot visualize complex functions as easily as real functions. We will now discuss some alternatives.

Image grid

A useful way to picture a complex function is to sketch its values on a grid G . The image grid $f(G)$ is a distorted version of the original grid which can be navigated easily using the grid lines. For example, to determine $f(1 + 2i)$, take one step in x -direction and two steps in y -direction on the distorted grid. Formally, let $G = \{z = x + iy \in \mathbb{C} \mid x \in \mathbb{Z} \text{ or } y \in \mathbb{Z}\}$ be the *unit grid* and define the *image grid* as (see Figure 2.3)

$$f(G) = \{w = u + iv \in \mathbb{C} \mid \exists z \in G : f(z) = w\}.$$




Figure 2.3: The unit grid G and the image grid $f(G)$ for $f(z) = \frac{1+i}{\sqrt{2}}z$

In practice, the image grid can often be described by finding a familiar equation that all of its members $u + iv$ satisfy. When this is not possible, a computer will help sketching an approximate image grid.

Example 2.6.

Consider $f(z) = z^2$. Then

$$u = x^2 - y^2, \quad v = 2xy. \quad (2.7)$$

To determine $f(G)$, first fix $x = \pm 1, \pm 2, \pm 3, \dots$ to be a non-zero integer. Using Equation 2.7 we find $u = \frac{-1}{4x^2}v^2 + x^2$. This is a downward parabola in the (u, v) -plane rotated by 90 degrees with vertex at $(u, v) = (x^2, 0) = (1, 0), (4, 0), (9, 0), \dots$. Similarly, if we fix y to be a non-zero integer, then $u = \frac{1}{4y^2}v^2 - y^2$ is an upward parabola in the (u, v) -plane rotated by 90 degrees. For $x = 0$, we get $(u, v) = (-y^2, 0)$ which parameterizes the negative u -axis. Similarly, for $y = 0$ we get the positive u -axis. All this is summarized in Figure 2.4.




Figure 2.4: Unit grid and image grid of $f(z) = z^2$

Example 2.7.

Consider $f(z) = 1/z$. Then

$$u = \frac{x}{x^2 + y^2}, \quad v = \frac{-y}{x^2 + y^2}. \quad (2.8)$$

Fixing $x = \pm 1, \pm 2, \dots$, we have $(u - \frac{1}{2x})^2 + v^2 = \frac{1}{4x^2}$ (verify by substituting Equation 2.8 into this equation). This is a circle of radius $\frac{1}{2x}$ and with center $(u, v) = (\frac{1}{2x}, 0)$. For $x = 0$, equations Equation 2.8 become $(u, v) = (0, -1/y)$ which parameterizes the v -axis. Similarly, for $0 \neq y \in \mathbb{Z}$ we find $u^2 + (v + \frac{1}{2y})^2 = \frac{1}{4y^2}$ and for $y = 0$ we obtain a parametrization of the u -axis. This is summarized in Figure 2.5 (and on the title page).




Figure 2.5: Unit grid and image grid of $f(z) = 1/z$. Notice that $z = 0$ gets sent to a 'point at infinity' that is imagined to surround the complex plane.

The previous example can be generalized.

Example 2.8.

Rational functions Equation 2.2 with $f(z) = az + b$ and $g(z) = cz + d$ affine linear, where $a, b, c, d \in \mathbb{C}$, are *Möbius transformations*. Thus

$$f(z) = \frac{az + b}{cz + d}$$

with domain $D = \mathbb{C} \setminus \{-d/c\}$ if $c \neq 0$ and domain $D = \mathbb{C}$ if $c = 0$. To exclude constant functions, we also assume that $ad - bc \neq 0$.

Domain colouring

We represent each unit complex number $e^{i\theta}$ by a color on the color wheel. The modulus r of an arbitrary complex number $re^{i\theta}$ will be represented by the lightness of the color. This assigns a unique color to each complex number, see Figure 2.6. Pure white is never used and would correspond to infinity. Pure black corresponds to the origin.

This is less useful for calculations by hand but generates artistic images using a computer.




Figure 2.6: Representing complex numbers z by color

This can be used for visualizing complex functions. Draw each point z in the domain of $w = f(z)$ using the color for w .




Figure 2.7: Domain colouring of $f(z) = \frac{(z-2)(z-1)^2}{z^2+1}$. Notice that near $z = \pm i$, where f is undefined, the function tends to infinity (pure white). The zeros of f at $z = 1, 2$ can also be seen.

3-dimensional graphs

Another approach is to plot the 3-dimensional graph of any of the following real-valued functions $D \rightarrow \mathbb{R}$

$$u, v, |f| = \sqrt{u^2 + v^2}.$$

Again, the missing information can be color-coded (see Figure 2.8).




Figure 2.8: The graph of $|f(z)|$ colored by $\arg(z)$

Questions for further discussion

- What are the functions in Example 2.1 geometrically?
- Does Equation 2.6 remain valid for all $z \in \mathbb{C}$? What is the correct modification?
- The two zeros in Figure 2.7 have a slightly different character. What is the difference between the zeros that might account for this?
- In Example 2.6 and Example 2.7} almost all the image gridlines meet at a right angle. The only exception is at $f(0)$ in Example 2.6. Use polar coordinates to explain this behavior of $f(z) = z^2$ at $z = 0$.
- Is it always possible to extend the domain of definition of a complex function?

Is this always sensible?

- Can you think of other branches of the logarithm?

2.1 Exercises

Exercise 2.1

Describe the image set of the complex function $f(z) = \frac{1+z}{1-z}$ with domain $D = \mathbb{C} \setminus \{1\}$. In other words, determine the set of all $w \in \mathbb{C}$ for which $w = \frac{1+z}{1-z}$ has a solution $z \in D$.

Show Solution 2.1 on P169

Exercise 2.2

Sketch the following curves $z(t)$ in the complex plane, where t is a real parameter.

- $z(t) = t(1 + i)$ for $0 \leq t \leq 1$
- $z(t) = \cos(t) + i \sin(t)$ for $0 \leq t \leq \pi$
- $z(t) = \cos(t) - i \sin(t)$ for $0 \leq t \leq \pi/2$
- $z(t) = \frac{1}{1+it}$ for $t \in \mathbb{R}$

Show Solution 2.2 on P170

Exercise 2.3

Let $\log(z)$ be the principal branch of the logarithm. Compute

$$\log(2i), \log(1 + i), \log(-3i), \log(5).$$

Show Solution 2.3 on P173

Exercise 2.4

Describe the following complex functions geometrically.

$$f(z) = 3z, f(z) = iz, f(z) = \frac{(1+i)}{\sqrt{2}}z$$

Show Solution 2.4 on P174

Exercise 2.5

Determine a domain of definition for the following complex functions.

$$f(z) = \frac{1}{z}, f(z) = \frac{1+z}{z-1}, f(z) = \frac{z^2-4}{z^2+2z}, f(z) = \frac{1}{\exp(z)}.$$

Show Solution 2.5 on P174

Exercise 2.6

Determine the domain of definition for $f(z) = \frac{1}{\sin(z)}$.

Show Solution 2.6 on P175

Exercise 2.7

Find all solutions to the following equations:

- a. $e^z = -1$,
- b. $\sin(z) = -i$

Show Solution 2.7 on P176

Exercise 2.8

Prove that the composition $f \circ f'$ of two Möbius transformations is again a Möbius transformation. If we associate to f, f' the matrices

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, A' = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in M_{2 \times 2}(\mathbb{C}),$$

show that $f \circ f'$ is associated to the matrix product AA' .

Show Solution 2.8 on P176

Exercise 2.9

Draw the image grid for $\exp: \mathbb{C} \rightarrow \mathbb{C}^\times$.

Show Solution 2.9 on P177

Further resources

- <https://youtu.be/NtolXhUgqSk> 5 ways to visualize a complex function
- Animation of Möbius transformations: <https://youtu.be/0z1flsUNhO4> and <https://www-users.cse.umn.edu/~arnold/moebius/>

Chapter 3

Topology of the complex plane

In this section we explain that most of the facts about limits, series, and continuity carry over from real analysis essentially without change.

The modulus of complex numbers defines a *distance* $d(z, w) = |z - w|$ on the plane (this is the usual Euclidean distance), which determines the following standard terminology for metric spaces.

Definition 3.1. The *open disc* of radius $0 \leq r \leq +\infty$ centered at $z_0 \in \mathbb{C}$ is

$$D_r(z_0) = \{z \in \mathbb{C} \mid |z - z_0| < r\}.$$

The *closed disc* $\overline{D}_r(z_0)$ is the set of all $z \in \mathbb{C}$ with $|z - z_0| \leq r$.

Definition 3.2. A subset $O \subset \mathbb{C}$ is *open* if for every point $z_0 \in O$ there exists $r > 0$ such that $D_r(z_0) \subset O$ (see Figure 3.1). A subset $C \subset \mathbb{C}$ is called *closed* if the complement $O = \mathbb{C} \setminus C$ is an open subset.

A subset $B \subset \mathbb{C}$ is *bounded* if $B \subset D_r(0)$ for some $0 < r < \infty$.




Figure 3.1: The gray open disc is an open subset of $\mathbb{C}$.

Remark 3.1. The above notion of open set determines a topology on $\mathbb{C}$.

Definition 3.3. A sequence of complex numbers $(z_n)_{n \in \mathbb{N}}$ has the *limit* $\zeta \in \mathbb{C}$ (or is *convergent* to ζ), written $\lim_{n \rightarrow \infty} z_n = \zeta$, if

$$\forall \epsilon > 0 \exists n_0 \in \mathbb{N} \forall n \geq n_0 : |z_n - \zeta| < \epsilon.$$

Equivalently, $\lim_{n \rightarrow \infty} |z_n - \zeta| = 0$. We call $(z_n)_{n \in \mathbb{N}}$ a *Cauchy sequence* if

$$\forall \epsilon > 0 \exists n_0 \in \mathbb{N} \forall n, m \geq n_0 : |z_n - z_m| < \epsilon.$$

The same argument as in real analysis shows that the limit w is unique and that every convergent sequence is a Cauchy sequence. The converse is also true by the completeness of real numbers.

Proposition 3.1. *For a sequence $(z_n)_{n \in \mathbb{N}}$ in $\mathbb{C}$, the following are equivalent:*

- a. *There exists $\zeta \in \mathbb{C}$ such that $\zeta = \lim_{n \rightarrow \infty} z_n$.*
- b. *$(z_n)_{n \in \mathbb{N}}$ is a Cauchy sequence.*

Proof.

To prove (b) $\implies$ (a) write $z_n = x_n + iy_n$ and notice that

$$|z_n - z_m| = \sqrt{(x_n - x_m)^2 + (y_n - y_m)^2} \geq |x_n - x_m|, |y_n - y_m|$$

implies that both $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ are Cauchy sequences in $\mathbb{R}$. By the completeness of $\mathbb{R}$, these sequences have limits $\chi, v \in \mathbb{R}$. Set $\zeta = \chi + iv$ and pick $n_0 \in \mathbb{N}$ such that $|x_n - \chi| < \frac{\epsilon}{\sqrt{2}}$ and $|y_n - v| < \frac{\epsilon}{\sqrt{2}}$ for all $n \geq n_0$. Then

$$|z_n - \zeta| = \sqrt{(x_n - \chi)^2 + (y_n - v)^2} < \sqrt{\epsilon^2/2 + \epsilon^2/2} = \epsilon$$

for all $n \geq n_0$. The converse, (a) $\implies$ (b), is left as an exercise.

□

The advantage of Cauchy sequences is that one does not need to know the value of the limit in advance.

Definition 3.4. A series of complex numbers $(z_k)_{k \in \mathbb{N}}$ converges to the limit ζ , written $\zeta = \sum_{k=0}^{\infty} z_k$, if the sequence of partial sums $(w_n = \sum_{k=0}^n z_k)_{n \in \mathbb{N}}$ converges to ζ . We call a series *absolutely convergent* if the series $\sum_{k=0}^{\infty} |z_k|$ is convergent.

As in real analysis, the Cauchy criterion implies that every absolutely convergent series is convergent. Absolutely convergent series may be rearranged and orders of summation may be exchanged.

Although $\infty \notin \mathbb{C}$, it will be convenient to define $\lim_{n \rightarrow \infty} z_n = \infty$ to mean that the sequence $(z_n)_{n \in \mathbb{N}}$ eventually leaves every disk. Symbolically,

$$\forall r > 0 \exists n_0 \in \mathbb{N} \forall n \geq n_0 : |z_n| > r.$$

We call $\mathbb{C} \cup \{\infty\}$ the *extended complex plane*.

Definition 3.5. A point $\zeta \in \mathbb{C} \cup \{\infty\}$ is in the *closure* of $D \subset \mathbb{C}$ if there exists a sequence $(z_n)_{n \in \mathbb{N}}$ with $z_n \in D$ and $\lim_{n \rightarrow \infty} z_n = \zeta$.

Let $f: D \rightarrow \mathbb{C}$ be a complex function and let $\zeta \in \mathbb{C} \cup \{\infty\}$ be in the closure of D . The function $f(z)$ has the *limit* $w \in \mathbb{C} \cup \{\infty\}$ as $z \rightarrow \zeta$, written $\lim_{z \rightarrow \zeta} f(z) = w$, if for every sequence $(z_n)_{n \in \mathbb{N}}$ with $z_n \in D$ and $\lim_{n \rightarrow \infty} z_n = \zeta$ we have $\lim_{n \rightarrow \infty} f(z_n) = w$. An equivalent ε - δ -definition is

$$\forall \varepsilon > 0 \exists \delta > 0 : 0 < |z - \zeta| < \delta, z \in D \implies |f(z) - w| < \varepsilon.$$

A complex function $f: D \rightarrow \mathbb{C}$ is *continuous* at $\zeta \in D$ if $\lim_{z \rightarrow \zeta} f(z) = f(\zeta)$. We call f *continuous on* D if f is continuous at every $\zeta \in D$.




Figure 3.2: The ε - δ -definition of $\lim_{z \rightarrow \zeta} f(z) = w$.

Convention

There is a some ambiguity in writing an expression of the form $\lim_{n \rightarrow \infty} a_n = f(n)$ as this could be interpreted either as a limit as a complex number z tends to infinity or as a limit over a sequence a_n .

In order to resolve this, unless explicitly stated otherwise, in this context, n shall refer to a natural number and a limit $\lim_{n \rightarrow \infty}$ should be interpreted as a limit over a sequence indexed by the natural numbers.

Questions for further discussion

- Explain the difference between the notions 'limit of a function', 'limit of a sequence', and 'limit of a series'.

3.1 Exercises

Exercise 3.1

Recall the ratio and root test for series of real numbers.

Show Solution 3.1 on P178

Exercise 3.2

For which $z \in \mathbb{C}$ do the following limits exist?

$$\lim_{n \rightarrow \infty} n^{1/n} z, \lim_{n \rightarrow \infty} z^n, \lim_{n \rightarrow \infty} \frac{z^n}{n}, \lim_{n \rightarrow \infty} \frac{z^n}{n!}, \lim_{n \rightarrow \infty} \frac{z^n}{n^n}, \lim_{n \rightarrow \infty} n! z.$$

Show Solution 3.2 on P179

Exercise 3.3

Show that every convergent sequence $(z_n)_{n \in \mathbb{N}}$ of complex numbers is bounded.

Show Solution 3.3 on P180

Exercise 3.4

Let $\sum_{k=0}^{\infty} z_k$ be a convergent series of complex numbers. Show that $\lim_{k \rightarrow \infty} z_k = 0$.

Show Solution 3.4 on P180

Exercise 3.5

For which $z \in \mathbb{C}$ do the following series converge?

$$\sum_{k=0}^{\infty} k z^k, \quad \sum_{k=0}^{\infty} (k z)^k$$

Show Solution 3.5 on P181

Exercise 3.6

Let $f: \mathbb{C} \rightarrow \mathbb{C}$ be a complex function. Show that

$$\lim_{z \rightarrow \infty} f(z) = w \iff \lim_{z \rightarrow 0} f\left(\frac{1}{z}\right) = w.$$

Show Solution 3.6 on P182

Exercise 3.7

The Riemann sphere is $\mathbb{S} = \{(a, b, c) \in \mathbb{R}^3 \mid a^2 + b^2 + c^2 = 1\}$. Show that the stereographic projection

$$F: \mathbb{S} \rightarrow \mathbb{C} \cup \{\infty\}, F(a, b, c) = \begin{cases} \frac{a+ib}{1-c} & \text{if } c \neq 1, \\ \infty & \text{if } c = 1. \end{cases}$$

is a bijection between the Riemann sphere and the extended complex plane. Find a formula for the inverse function.




Figure 3.3: Stereographic projection from the north pole

Show Solution 3.7 on P182

Further resources

Part II

Differentiation and contour integrals

Chapter 4

Holomorphic functions

The following definition is a key concept for this course.

Definition 4.1. A complex function $f: U \rightarrow \mathbb{C}$ with domain an open set $U \subset \mathbb{C}$ is *complex differentiable* at a point $z_0 \in U$ if the limit

$$f'(z_0) = \frac{\partial f}{\partial z}(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} \quad (4.1)$$

exists. We call f *holomorphic* on U if f is complex differentiable at every $z_0 \in U$. The set of all holomorphic functions on U is denoted $\mathcal{O}(U)$. A function that is holomorphic¹ on $U = \mathbb{C}$ is *entire*.

This definition is structurally the same as for real functions. For example, f is continuous at z_0 since multiplying Equation 4.1 by $0 = \lim_{z \rightarrow z_0} (z - z_0)$ gives

$$0 = \lim_{z \rightarrow z_0} f(z) - f(z_0) \implies f(z_0) = \lim_{z \rightarrow z_0} f(z).$$

¹From Greek *holos* 'whole, complete' and *morphē* 'form', shape

Other formal proofs carry over as well and show the following.

Proposition 4.1. *Let $f, g: U \rightarrow \mathbb{C}$ be complex differentiable at z_0 . Then:*

a. $f + g$ is complex differentiable at z_0 with

$$(f + g)'(z_0) = f'(z_0) + g'(z_0).$$

b. fg is complex differentiable at z_0 with

$$(fg)'(z_0) = f'(z_0)g(z_0) + f(z_0)g'(z_0).$$

c. If $g(z_0) \neq 0$, then f/g is complex differentiable at z_0 with

$$(f/g)'(z_0) = \frac{f'(z_0)g(z_0) - f(z_0)g'(z_0)}{g(z_0)^2}.$$

Proof.

Exercise, using the algebra of limits.

□

Example 4.1.

Directly from Equation 4.1 we find that the constant function $f(z) = c$ is entire with $f'(z) = 0$. The identity function $f(z) = z$ is entire with $f'(z) = 1$. Proposition 4.1 and induction then imply that all polynomials

$$f(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0$$

are entire with

$$f'(z) = na_n z^{n-1} + (n-1)a_{n-1} z^{n-2} + \dots + a_1.$$

It then follows from Proposition 4.1 part c. that rational functions are holomorphic on their domain of definition.

Proposition 4.2. *Let $f: U \rightarrow \mathbb{C}$, $g: V \rightarrow \mathbb{C}$ be complex functions with $f(U) \subset V$, and U, V open. Suppose that f is complex differentiable at $z_0 \in U$ and that g is complex differentiable at $w_0 = f(z_0) \in V$.*

Then $g \circ f$ is complex differentiable at z_0 and

$$(g \circ f)'(z_0) = g'(w_0)f'(z_0). \quad (4.2)$$

Proof.

We can equivalently rewrite Equation 4.1 as

$$f(z_0 + h) = f(z_0) + hf'(z_0) + h\rho(h), \quad \lim_{h \rightarrow 0} \rho(h) = 0, \quad (4.3)$$

where the function $\rho(h)$, $h \neq 0$, is defined by this equation. Similarly,

$$g(w_0 + k) = g(w_0) + kg'(w_0) + k\sigma(k), \quad \lim_{k \rightarrow 0} \sigma(k) = 0. \quad (4.4)$$

Then

$$\begin{aligned} \frac{(g \circ f)(z_0 + h) - (g \circ f)(z_0)}{h} &= \frac{g(w_0 + \overbrace{hf'(z_0) + h\rho(h)}^{k=}) - g(w_0)}{h} \\ &= f'(z_0)g'(w_0) + [\rho(h)g'(w_0) + (f'(z_0) + \rho(h))\sigma(hf'(z_0) + h\rho(h))], \end{aligned}$$

where the first equality follows from Equation 4.3, the second from Equation 4.4 and the square bracket tends to zero as $h \rightarrow 0$, using the chain rule for limits to see that $\lim_{h \rightarrow 0} \sigma(hf'(z_0) + h\rho(h)) = \lim_{k \rightarrow 0} \sigma(k) = 0$.

□

One should view Equation 4.3 as a short Taylor series expansion: near the point z_0 the function f is given to zeroth order by a constant $f(z_0)$, with first order correction given by a *complex linear* function $h \mapsto hf'(z_0)$, plus a higher order term $h\rho(h)$ which vanishes infinitesimally (after dividing by h).

We next discuss how complex differentiability compares to real differentiability. Regard a complex function as a real multivariable map

$$f = (u, v): U \longrightarrow \mathbb{R}^2, \quad U \subset \mathbb{R}^2 \text{ open set.}$$

Recall that f is *real differentiable* at $z_0 = (x_0, y_0)$ if there exists a matrix $J_f(z_0) \in M_{2 \times 2}(\mathbb{R})$, called the *Jacobian*, so that Equation 4.3 holds with the complex multiplication $hf'(z_0)$ replaced by the matrix vector multiplication $J_f(z_0)h$. Real differentiability at z_0 implies that the partial derivatives at z_0 exist and

$$J_f(z_0) = \begin{pmatrix} \frac{\partial u}{\partial x}(z_0) & \frac{\partial u}{\partial y}(z_0) \\ \frac{\partial v}{\partial x}(z_0) & \frac{\partial v}{\partial y}(z_0) \end{pmatrix}. \quad (4.5)$$

Recall that the converse requires the additional assumptions that the partial derivatives exist *in a neighborhood* of z_0 and are *continuous* at z_0 .

Proposition 4.3. *For a complex function $f = u + iv$ the following are equivalent:*

- a. *f is complex differentiable at z_0*

b. f is real differentiable at z_0 and the Cauchy–Riemann equations

$$\frac{\partial u}{\partial x}(z_0) = \frac{\partial v}{\partial y}(z_0), \quad \frac{\partial u}{\partial y}(z_0) = -\frac{\partial v}{\partial x}(z_0), \quad (4.6)$$

hold.

In this case, $f'(z_0) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$.

Proof.

The two definitions are identical except that the complex multiplication $f'(z_0)h$ by $h \in \mathbb{C}$ is replaced by the matrix vector multiplication $J_f(z_0)h$ by $h \in \mathbb{R}^2$. Our task is to compare these two notions. For this we simply observe that by Proposition 1.2 $J_f(z_0)$ is $\mathbb{C}$ -linear if and only if Equation 4.6 holds.

□




Figure 4.1: Bernhard Riemann. Familienarchiv Thomas Schilling, ca. 1862. [WikiMedia Commons](#). Public Domain

Example 4.2.

For the exponential function, $u = e^x \cos(y)$, $v = e^x \sin(y)$. Then the partial derivatives exist

$$\frac{\partial u}{\partial x} = e^x \cos(y), \quad \frac{\partial u}{\partial y} = -e^x \sin(y), \quad \frac{\partial v}{\partial x} = e^x \sin(y), \quad \frac{\partial v}{\partial y} = e^x \cos(y),$$

are continuous functions on $\mathbb{R}^2$, and satisfy the Cauchy–Riemann equations. It follows that $\exp$ is entire with

$$\exp'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \exp(z). \quad (4.7)$$

Example 4.3.

The *hyperbolic cosine* $\cosh(z) = \frac{e^z + e^{-z}}{2}$ and the *hyperbolic sine* function $\sinh(z) = \frac{e^z - e^{-z}}{2}$ are entire with

$$\sinh'(z) = \cosh(z), \quad \cosh'(z) = \sinh(z).$$

Since $\cosh(z) = 0 \iff e^{2z} = -1 \iff z \in \frac{\pi i}{2} + \pi i \mathbb{Z}$ the *hyperbolic tangent* $\tanh(z) = \frac{\sinh(z)}{\cosh(z)}$ is a holomorphic function

$$\tanh: \mathbb{C} \setminus \left(\frac{\pi i}{2} + \pi i \mathbb{Z} \right) \longrightarrow \mathbb{C},$$

with

$$\tanh'(z) = \frac{\cosh(z)^2 - \sinh(z)^2}{\cosh(z)^2} = 1 - \tanh(z)^2.$$

Similarly, the *hyperbolic cotangent* $\coth(z) = \frac{\cosh(z)}{\sinh(z)}$ is a holomorphic function $\coth: \mathbb{C} \setminus \pi i \mathbb{Z} \rightarrow \mathbb{C}$ with

$$\coth'(z) = 1 - \coth(z)^2.$$

Example 4.4.

The trigonometric sine $\sin(z) = -i \sinh(iz)$ and the cosine $\cos(z) = \cosh(iz)$ are entire with $\sin'(z) = \cos(z)$, $\cos'(z) = -\sin(z)$. The tangent $\tan(z) = \frac{\sin(z)}{\cos(z)}$ is a holomorphic function on $\mathbb{C} \setminus (\frac{\pi}{2} + \pi \mathbb{Z})$ with $\tan'(z) = 1 + \tan(z)^2$. The cotangent

$\cot(z) = \frac{\cos(z)}{\sin(z)}$ is a holomorphic function on $\mathbb{C} \setminus \pi\mathbb{Z}$ with $\cot'(z) = 1 - \cot(z)^2$.

Example 4.5.

$f(z) = \bar{z} = x - iy$ has all partial derivatives of all orders, but is *not* complex differentiable at any $z_0 \in \mathbb{C}$ since $\frac{\partial u}{\partial x} = 1 \neq \frac{\partial v}{\partial y} = -1$.

Theorem 4.1. *Let $f: U \rightarrow \mathbb{C}$ be a holomorphic function and $z_0 \in U$ with $f'(z_0) \neq 0$. Then there exist open sets $V, W \subset \mathbb{C}$ with $z_0 \in V \subset U$ and $w_0 = f(z_0) \in W$ with the property that the restriction $f|_V$ becomes a bijection $V \rightarrow W$ with holomorphic inverse function $(f|_V)^{-1}: W \rightarrow V$ and*

$$\frac{\partial(f|_V)^{-1}}{\partial w}(w) = \frac{1}{f'((f|_V)^{-1}(w))}. \quad (4.8)$$

Proof.

As in the proof of Proposition 4.3, we view f as a differentiable real multivariable map $U \rightarrow \mathbb{R}^2$. Since $f'(z_0) = \frac{\partial u}{\partial x}(z_0) + i\frac{\partial u}{\partial y}(z_0) \neq 0$ we have

$$\begin{aligned} \det J_f(z_0) &= \frac{\partial u}{\partial x}(z_0) \frac{\partial v}{\partial y}(z_0) - \frac{\partial v}{\partial x}(z_0) \frac{\partial u}{\partial y}(z_0) \\ &= \left(\frac{\partial u}{\partial x}(z_0) \right)^2 + \left(\frac{\partial u}{\partial y}(z_0) \right)^2 \neq 0. \end{aligned}$$

where the first equality follows from Equation 4.5 and the second from Equation 4.6. We may thus apply the inverse function theorem from multivariable calculus. This gives open sets with $z_0 \in V \subset U, w_0 \in W$ such that $f|_V: V \rightarrow W$ is a bijection and $(f|_V)^{-1}: W \rightarrow V$ is real differentiable. In particular, $\det J_f(z) \neq 0$ for all $z \in V$ (chain rule). Let $w \in W$ and $z = f^{-1}(w)$. Then

$$J_{(f|_V)^{-1}}(w) = J_f(z)^{-1} = \frac{1}{\det J_f(z)} \begin{pmatrix} \frac{\partial v}{\partial y}(z) & -\frac{\partial u}{\partial y}(z) \\ -\frac{\partial v}{\partial x}(z) & \frac{\partial u}{\partial x}(z) \end{pmatrix},$$

so the Cauchy–Riemann equations are satisfied at every $w \in W$ for the inverse function, so $(f|_V)^{-1}$ is holomorphic. Finally, Equation 4.8 follows by applying the chain rule Equation 4.2 to $f|_V \circ (f|_V)^{-1} = \text{id}_W$ which gives

$$(f|_V)'((f|_V)^{-1}(w)) \cdot \frac{\partial (f|_V)^{-1}}{\partial w}(w) = 1.$$

□

Remark 4.1. Our proof is a simple application of the ordinary inverse function theorem. We will see two different proofs later in the course.

Example 4.6.

Recall from Proposition 2.1 that $\exp|_S: S \rightarrow \mathbb{C}^-$ is a bijection with the logarithm as inverse function. As $\exp'(z) = \exp(z) \neq 0$, we conclude from Theorem 4.1 that $\log: \mathbb{C}^- \rightarrow S$ is holomorphic with

$$\log'(z) = 1/z. \tag{4.9}$$

Questions for further discussion

- $x \mapsto x^3$ is a real differentiable bijection whose inverse function is not differentiable at $x = 0$. What about the complex analogue $z \mapsto z^{1/3}$?

4.1 Exercises

Exercise 4.1

Write the following complex functions in the form $f = u + iv$:

$$f_1(z) = \sin(z), \quad f_2(z) = e^{z^2}, \quad f_3(z) = \cosh(z)$$

Show Solution 4.1 on P184

Exercise 4.2

At which points $z \in \mathbb{C}$ are the following functions complex differentiable? At which points are they real differentiable?

$$\begin{aligned} f_1(z) &= z, & f_2(z) &= \bar{z}, & f_3(z) &= z^3 + z, \\ f_4(z) &= \frac{1}{2iz}, & f_5(z) &= |z|^2, & f_6(z) &= \frac{|z|^2}{\bar{z}} \end{aligned}$$

Show Solution 4.2 on P185

Exercise 4.3

Let $f: V \rightarrow W$ be a bijection of open sets $V, W \subset \mathbb{C}$. Assume that f is complex differentiable at z_0 and that f^{-1} is complex differentiable at $w_0 = f(z_0)$. Show that $f'(z_0) \neq 0$. (The same result holds for real differentiable maps to show $\det J_f(z_0) \neq 0$.)

Show Solution 4.3 on P186

Exercise 4.4

Let $f: U \rightarrow \mathbb{C}$ be a holomorphic function with domain an open set $U \subset \mathbb{C}$. Suppose also that $f'(z)$ is holomorphic. Write $f(z) = u(x, y) + iv(x, y)$, where $z = x + iy$. As we will see, a consequence of f being holomorphic, is that u and v have continuous second order derivatives. Show that:

- $|f'(z)|^2 = \left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial v}{\partial x}\right)^2 = \left(\frac{\partial u}{\partial y}\right)^2 + \left(\frac{\partial v}{\partial y}\right)^2$
- Both u, v satisfy the Laplace equation $\Delta(u) = \Delta(v) = 0$, where the Laplace operator is defined as $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$.
- Fix $n \in \mathbb{N}$. Use b. to find a solution to $\Delta(u) = 0$ that satisfies $u(z) = \cos(n\theta)$ for all $|z| = 1$, $z = e^{i\theta}$ on the unit circle.
- Does the converse to b. hold?

Show Solution 4.4 on P186

Exercise 4.5

Find subdomains where $\sin, \cos, \sinh, \cosh, \tan, \tanh$ are injective. Compute the derivative of an inverse using the chain rule. Deduce an explicit formula for the derivative of an inverse.

Show Solution 4.5 on P188

Exercise 4.6

Show that the set $\mathcal{O}(U)$ of holomorphic functions on U becomes a ring under the operations of point-wise addition and multiplication:

$$(f + g)(z) = f(z) + g(z), \quad (fg)(z) = f(z)g(z).$$

Show Solution 4.6 on P189

Exercise 4.7

Let $f: \mathbb{C} \rightarrow \mathbb{C}$ be an entire function with image $f(\mathbb{C}) \subset \mathbb{R}$. Prove that f is constant.

Show Solution 4.7 on P190

Exercise 4.8

Let U be an open set and assume that U is *path-connected*, meaning that for all $z_0, z_1 \in U$ there exists a continuously differentiable map $\gamma: [0, 1] \rightarrow U$ such that $\gamma(0) = z_0$, $\gamma(1) = z_1$.

- a. Let $f: U \rightarrow \mathbb{C}$ be a holomorphic function such that $f'(z) = 0$ for all $z \in U$. Prove that $f(z)$ is a constant function.

Hint: Consider the real and imaginary parts of $f \circ \gamma$.

- b. Show that the only entire functions $f(z)$ satisfying $f'(z) = Cf(z)$ for a constant $C \in \mathbb{C}$ are the functions $f(z) = De^{Cz}$, $D \in \mathbb{C}$.

- c. Find all entire solutions $f(z)$ of $f''(z) = f(z)$.

Hint: Consider $g = f + f'$, $h = f - f'$

Show Solution 4.8 on P190

Chapter 5

Power Series

Definition 5.1. A *formal power series* is an expression of the form

$$P = \sum_{n=0}^{\infty} a_n T^n \quad (5.1)$$

with *coefficients* $a_n \in \mathbb{C}$. As no convergence is required, this is really just a sequence $(a_n)_{n \in \mathbb{N}}$ of complex numbers.

We call a_0 the *constant term* of P . The *order* of P is the smallest $n \in \mathbb{N}$ such that $a_n \neq 0$.

Example 5.1.

Every polynomial is a formal power series.

Definition 5.2. Let $\mathbb{C}[[T]]$ be the set of all formal power series. Define the addition and multiplication of $P = \sum_{n=0}^{\infty} a_n T^n$, $Q = \sum_{n=0}^{\infty} b_n T^n \in \mathbb{C}[[T]]$ by

$$P + Q = \sum_{n=0}^{\infty} (a_n + b_n) T^n, \quad (5.2)$$

$$PQ = \sum_{n=0}^{\infty} c_n T^n, c_n = \sum_{i+j=n} a_i b_j. \quad (5.3)$$

Proposition 5.1.

- a. These operations make $\mathbb{C}[[T]]$ a commutative ring with unit 1, the series with constant term 1 and all higher coefficients zero.
- b. $P \in \mathbb{C}[[T]]$ has a multiplicative inverse $\iff$ the constant term of P is non-zero.

Proof.

- a. is straightforward and left as an exercise.
- b. Having an inverse amounts to the existence of $Q = \sum_{n=0}^{\infty} b_n T^n$ such that $PQ = 1$. By Equation 5.3 this means

$$a_0 b_0 = 1 \quad \sum_{i+j=n} a_i b_j = 0 (\forall n > 0). \quad (5.4)$$

The first equation shows that a_0 is invertible, proving ' $\implies$ '. For ' $\impliedby$ ', the idea is to use Equation 5.4 to construct the coefficients b_n of the inverse inductively. Of course, $b_0 = a_0^{-1}$. For $n = 1$, Equation 5.4 gives $a_0 b_1 + a_1 b_0 = 0$ so

$$b_1 = -a_0^{-2} a_1.$$

For $n = 2$, $a_0 b_2 + a_1 b_1 + a_2 b_0 = 0$. Inserting the known b_0, b_1 implies

$$b_2 = a_0^{-3} (a_1^2 - a_0 a_2).$$

For general n , given $b_0, \dots, b_{n-1}$ we rearrange Equation 5.4 as

$$b_n = -a_0^{-1} \sum_{j=0}^{n-1} a_{n-j} b_j,$$

which defines b_n by recursion.

□

It is unnecessary to remember Equation 5.4. It is enough to recall the ansatz $PQ = 1$ and the strategy of computing the coefficients of Q inductively.

Example 5.2.

Let $P = 1 - T$. Then $P^{-1} = \sum_{n=0}^{\infty} T^n$ is the geometric series. We verify $(1 - T) \sum_{n=0}^{\infty} T^n = \sum_{n=0}^{\infty} T^n - \sum_{n=1}^{\infty} T^n = 1$ (telescope sum).

Definition 5.3. The *domain* $D(P)$ of a formal power series Equation 5.1 is the set of all $z \in \mathbb{C}$ such that the series of complex numbers

$$P(z) = \sum_{n=0}^{\infty} a_n z^n,$$

obtained by substituting T by z , converges. We obtain a complex function

$$D(P) \longrightarrow \mathbb{C}, z \longmapsto P(z).$$

More generally, fix a *center* $z_0 \in \mathbb{C}$. We then have a complex function

$$D(P, z_0) := z_0 + D(P) \longrightarrow \mathbb{C}, z \longmapsto P(z - z_0) = \sum_{n=0}^{\infty} a_n (z - z_0)^n \quad (5.5)$$

which differs from $P(z)$ only by a translation.

Notational clash

There is a small clash in notation. Recall that for a complex number z and a positive real number r , we write $D_r(z)$ for the open disc of radius r about z , while for a formal power series P , $D(P)$ represents its domain of convergence. You can tell these apart by whether or not there is a subscript and also by referring to the context.

So for instance in interpreting a chain of the form:

$$D_\rho(z_0) \subset D(P, z_0) \subset \overline{D}_\rho(z_0)$$

- the first term refers to the open disc of radius ρ about the z_0 ,
- the second is the domain of the power series P with centre z_0 , and,
- the last is the closed disc of radius ρ about z_0 .

Example 5.3.

The geometric series $P = \sum_{n=0}^{\infty} T^n$ converges for all $|z| < 1$ since the partial sums $P_n(z) = \frac{z^{n+1}-1}{z-1}$ tend to $\frac{1}{1-z}$ as $n \rightarrow \infty$. The geometric series diverges for all $|z| \geq 1$, since in this case $(z^n)_{n \in \mathbb{N}}$ is not a null sequence. Hence $D(P) = D_1(0)$.

Recall the following concept from analysis.

Definition 5.4. For a complex function $f: D \rightarrow \mathbb{C}$, the *uniform norm* is

$$\|f\|_{\infty, D} = \sup_{z \in D} |f(z)|. \quad (5.6)$$

A sequence of functions $(f_n)_{n \in \mathbb{N}}$ is *uniformly convergent* to f on D if

$$\|f - f_n\|_{\infty, D} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

A series $\sum_{n=0}^{\infty} f_n(z)$ is a *uniformly convergent series* on D if the sequence of *partial sums* $P_n(z) = \sum_{k=0}^n f_k(z)$ converges uniformly.

Uniform convergence on a subset $C \subset D$ refers to the uniform convergence of the functions restricted to C .

Theorem 5.1. *Every formal power series P has a unique radius of convergence $0 \leq \rho \leq +\infty$ such that for every center z_0*

$$D_\rho(z_0) \subset D(P, z_0) \subset \overline{D}_\rho(z_0). \quad (5.7)$$

In fact,

$$\rho = \sup \{r \geq 0 \mid (|a_k| r^k)_{k \in \mathbb{N}} \text{ is a bounded sequence}\}. \quad (5.8)$$

Moreover, $P(z - z_0)$ converges absolutely and uniformly on every smaller disk $D_r(z_0)$ with $0 < r < \rho$ and diverges at every point $z \in \mathbb{C}$ with $|z - z_0| > \rho$.

Proof.

For uniqueness, suppose $D_{\rho_\ell}(z_0) \subset D(P, z_0) \subset \overline{D}_{\rho_\ell}(z_0)$ for $\ell = 1, 2$ and $\rho_1 \neq \rho_2$, say $0 \leq \rho_1 < \rho_2$. Pick $z \in \mathbb{C}$ with $\rho_1 < |z - z_0| < \rho_2$. Then $z \in D_{\rho_2}(z_0) \subset D(P, z_0) \subset \overline{D}_{\rho_1}(z_0)$, so $|z - z_0| \leq \rho_1$, a contradiction.

It suffices to prove the last statement in the theorem for ρ defined by Equation 5.8, because this implies Equation 5.7. For $z \in \mathbb{C}$ with $|z - z_0| > \rho$ the terms $a_k(z - z_0)^k$

do not tend to zero (they are unbounded), so the series $P(z - z_0)$ diverges.

Let $0 < r < \rho$. Using that ρ is the *least* upper bound, there exists $r < s < \rho$ with $|a_k|s^k < B$ bounded. Pick $w \in \mathbb{C}$ with $s = |w - z_0|$. For all $z \in D_r(z_0)$ we then have

$$\sum_{k=0}^{\infty} |a_k(z - z_0)^k| = \sum_{k=0}^{\infty} \left| a_k(w - z_0)^k \underbrace{\left(\frac{z - z_0}{w - z_0} \right)^k}_{q^k} \right| \leq B \sum_{k=0}^{\infty} q^k. \quad (5.9)$$

Since $|q| < 1$, the geometric series on the right converges. Hence the comparison test implies that $P(z - z_0)$ converges absolutely. As for uniform convergence, let P_n be the n -th partial sum. As in Equation 5.9, for all $z \in D_r(z_0)$

$$|P(z - z_0) - P_n(z - z_0)| = \left| \sum_{k=n}^{\infty} a_k(z - z_0)^k \right| \leq B \sum_{k=n}^{\infty} q^k.$$

The right hand side is independent of z , so $\|P - P_n\|_{\infty, D_r(z_0)} \xrightarrow{n \rightarrow \infty} 0$.

□

Remark 5.1. The domain $D(P)$ of a formal power series is roughly a disk of radius ρ , with uncertain behavior on the boundary circle $|z| = \rho$. Determining the behavior on the boundary can be very difficult. On the other hand, the radius of convergence is easy to compute. If you find $z \in \mathbb{C}$ for which $P(z)$ converges (for example, using the ratio or the root test), you can deduce $|z| < \rho$. If $P(z)$ diverges at $z \in \mathbb{C}$, you know $|z| \geq \rho$.

Example 5.4 (optional).

For $P = \sum_{n=1}^{\infty} \frac{z^n}{n}$ we have $\rho = 1$ by Equation 5.8. We investigate the behavior on the boundary circle $|z| = 1$. For $z = 1$, this is the divergent harmonic series. We claim

that P converges for all other boundary points, so $D(P) = \overline{D_1}(0) \setminus \{1\}$. Consider

$$\begin{aligned} (z-1) \sum_{k=1}^n \frac{z^k}{k} &= \sum_{k=2}^{n+1} \frac{z^k}{k-1} - \sum_{k=1}^n \frac{z^k}{k} \\ &= \frac{z^{n+1}}{n} - z + \sum_{k=2}^n \left(\frac{z^k}{k-1} - \frac{z^k}{k} \right) \\ &= \frac{z^{n+1}}{n} - z + \sum_{k=2}^n \frac{z^k}{k(k-1)}. \end{aligned}$$

The series $\sum_{k=2}^{\infty} \frac{z^k}{k(k-1)}$ is (absolutely) convergent for $|z| \leq 1$. It follows that the left hand sequence of partial sums converges as $n \rightarrow \infty$. Dividing by $z-1 \neq 0$ implies that P converges.

Corollary 5.1. *The restriction of the complex function $f(z) = P(z - z_0)$ defined in Equation 5.5 to $D_\rho(z_0) \subset D(P, z_0)$ is continuous.*

Proof.

Since the translation $z \mapsto z - z_0$ is a homeomorphism, it suffices to consider the case $z_0 = 0$. We prove continuity at each $w \in D_\rho(0)$. Let $\epsilon > 0$. By uniform convergence, we can pick $n \in \mathbb{N}$ with $\|P_n - P\|_{\infty, D_\rho(0)} < \epsilon/3$. Since the polynomial $P_n(z)$ is continuous, there is $\delta > 0$ such that $|P_n(z) - P_n(w)| < \epsilon/3$ for all $z \in D_\delta(w)$. By shrinking δ , we may suppose $D_\delta(w) \subset D_\rho(0)$. Then have

$$|P(z) - P(w)| \leq |P(z) - P_n(z)| + |P_n(z) - P_n(w)| + |P_n(w) - P(w)| < \epsilon.$$

for all $z \in D_\delta(w)$.

□

Remark 5.2. Continuity need not hold on $D(P, z_0)$ (Sierpinski, 1916).

Example 5.5.

The series $P = \sum_{n=1}^{\infty} \frac{z^n}{n^2}$ has $\rho = 1$. On the boundary circle, $|z^n/n^2| \leq 1/n^2$ and the series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges and is independent of z . This implies uniform convergence and the proof of Corollary 5.1 shows that P is continuous on its domain $D(P) = \overline{D}_1(0)$.

When the radius of convergence is positive, we drop the word ‘formal’ and simply speak of a (convergent) *power series*.

Example 5.6.

The *exponential series*

$$\exp(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$$

has radius of convergence $+\infty$, since $\frac{z^n}{n!}$ is bounded (even tends to zero), as factorials grow faster than powers. Alternatively, for each $z \in \mathbb{C}$ we have absolute convergence by the ratio test

$$\left| \frac{z^{n+1}/(n+1)!}{z^n/n!} \right| = \frac{|z|}{n+1} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Therefore the power series defines a complex function $\exp: \mathbb{C} \rightarrow \mathbb{C}$. Mathematically, this is a good definition for the exponential function. Using Equation 2.3 we obtain from this the series expansions of the sine and the cosine:

$$\cos(z) = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!}, \quad \sin(z) = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!}.$$

Example 5.7 (optional).

We first generalize the definition of *binomial coefficients* to complex numbers $\alpha \neq 0$

and $k \in \mathbb{N}$. Set

$$\binom{\alpha}{k} = \begin{cases} \frac{\alpha(\alpha-1)\cdots(\alpha-k+1)}{k!} & \text{if } k > 0, \\ 1 & \text{if } k = 0. \end{cases}$$

The *binomial series* is

$$B_\alpha = \sum_{k=0}^{\infty} \binom{\alpha}{k} z^k.$$

For $\alpha \in \mathbb{C} \setminus \mathbb{N}$ we have

$$\left| \frac{\binom{\alpha}{k+1} z^{k+1}}{\binom{\alpha}{k} z^k} \right| = \left| z \frac{\alpha - k}{k+1} \right| \rightarrow |z| \text{ as } k \rightarrow \infty.$$

Therefore, the ratio test implies $\rho = 1$ when $\alpha \notin \mathbb{N}$. When $\alpha \in \mathbb{N}$ only finitely many binomial coefficients $\binom{\alpha}{k}$ are non-zero, so B_α is a polynomial and $\rho = +\infty$ and indeed by the binomial theorem we have

$$B_\alpha(z) = (z+1)^\alpha, \quad \forall \alpha \in \mathbb{N}.$$

Theorem 5.2. *Let P be a formal power series. Assume the radius of convergence $\rho > 0$ is positive. Then for each center $z_0 \in \mathbb{C}$ the function*

$$P: D_\rho(z_0) \rightarrow \mathbb{C}, z \mapsto P(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n \quad (5.10)$$

is holomorphic with derivative given by termwise differentiation,

$$P'(z) = \sum_{n=1}^{\infty} n a_n (z - z_0)^{n-1}. \quad (5.11)$$

Hence P is infinitely complex differentiable, by induction.

Proof.

We show that P is complex differentiable at every $w \in D_\rho(z_0)$. As $\sum_{k=0}^{\infty} kq^k$ converges for $|q| < 1$, the same argument given in Equation 5.9 shows that Equation 5.11 is a convergent series of complex numbers. For simplicity of notation, suppose $z_0 = 0$. We have absolutely convergent series

$$\begin{aligned} \frac{P(w+h) - P(w)}{h} - P'(w) &= \sum_{n=0}^{\infty} a_n \frac{(w+h)^n - w^n}{h} - P'(w) \\ &= \sum_{n=0}^{\infty} a_n \left(\sum_{k=1}^n \binom{n}{k} w^{n-k} h^{k-1} \right) - P'(w) \\ &= \sum_{n=0}^{\infty} \sum_{k=2}^n a_n \binom{n}{k} w^{n-k} h^{k-1} \\ &= \sum_{k=2}^{\infty} \sum_{n=0}^{\infty} a_n \binom{n}{k} w^{n-k} h^{k-1}, \end{aligned}$$

where we use $\binom{n}{k} = 0$ for $k > n$ and where we have exchanged the order of summation, by absolute convergence. The above is a power series $R(z) = \sum_{k=0}^{\infty} c_k z^k$ with $c_{k-1} = \sum_{n=0}^{\infty} a_n \binom{n}{k} w^{n-k}$ for $k \geq 2$ and $c_0 = 0$. Our calculation shows that $R(h)$ converges, so the radius of convergence of R is at least h . Therefore $R(h)$ is continuous on $D_h(z_0)$, so $\lim_{h \rightarrow 0} R(h) = R(0) = c_0 = 0$. In other words, $\frac{P(w+h) - P(w)}{h} - P'(w) = R(h) \rightarrow 0$ as $h \rightarrow 0$.

□

The *identity theorem* for power series is the following result.

Corollary 5.2. *Let $P = \sum_{n=0}^{\infty} a_n z^n$, $Q = \sum_{n=0}^{\infty} b_n z^n$ be formal power series with positive radius of convergence. Let $0 \neq z_\ell \in D(P) \cap D(Q)$ be a null sequence, $\lim_{\ell \rightarrow \infty} z_\ell = 0$, of non-zero complex numbers in the common domain. If $P(z_\ell) = Q(z_\ell)$ for all ℓ , then all coefficients $a_n = b_n$ agree.*

Proof.

We prove this by induction.

Base case $n = 0$. By continuity of P and Q at 0,

$$\begin{aligned} a_0 &= P(0) = P\left(\lim_{\ell \rightarrow \infty} z_\ell\right) = \lim_{\ell \rightarrow \infty} P(z_\ell) \\ &= \lim_{\ell \rightarrow \infty} Q(z_\ell) = Q\left(\lim_{\ell \rightarrow \infty} z_\ell\right) = Q(0) = b_0. \end{aligned}$$

Inductive step $n+1$. Assume that $a_0 = b_0, \dots, a_n = b_n$. By subtracting $\sum_{k=0}^n a_k z_\ell^k = \sum_{k=0}^n b_k z_\ell^k$ from $P(z_\ell)$ and $Q(z_\ell)$ we find

$$\sum_{k=n+1}^{\infty} a_k z_\ell^k = \sum_{k=n+1}^{\infty} b_k z_\ell^k.$$

As $z_\ell \neq 0$, we can divide this equation by z_ℓ^{n+1} and get $\tilde{P}(z_\ell) = \tilde{Q}(z_\ell)$ for the formal power series $\tilde{P} = \sum_{k=0}^{\infty} a_{k+n+1} z_\ell^k$ and $\tilde{Q} = \sum_{k=0}^{\infty} b_{k+n+1} z_\ell^k$, which have the same (positive) radius of convergence. An application of the base case to $\tilde{P}$ and $\tilde{Q}$ then shows that $a_{n+1} = b_{n+1}$ for the constant terms, as required.

□

Questions for further discussion

- Give precise statements of the comparison test, the ratio test and root test. Recall how the ratio test is proven by comparison with the geometric series.
- Find a power series P with $D(P) = \overline{D_1}(0) \setminus \{\pm 1, \pm i\}$.
- Explain why ‘is a bounded sequence’ in Equation 5.8 can be replaced by ‘is a null sequence’.

5.1 Exercises

Exercise 5.1

Find the radius of convergence for the following power series centered at the origin.

- i. $\sum_{n=0}^{\infty} \frac{z^n}{n^3},$
- ii. $\sum_{n=0}^{\infty} z^{3n},$
- iii. $\sum_{n=0}^{\infty} \frac{z^n}{n^n}.$

Hint: Recall the ratio and root tests for series of complex numbers

Show Solution 5.1 on P192

Exercise 5.2

Treating e^z , $\sin(z)$, $\cos(z)$ as formal power series with their usual Taylor expansion, find the terms of order ≤ 3 of the following power series:

- i. $e^z \sin(z),$
- ii. $\sin(z) \cos(z),$
- iii. $1/\cos(z)$

Show Solution 5.2 on P193

Exercise 5.3

Let $P = \sum_{n=0}^{\infty} a_n z^n$, $Q = \sum_{n=0}^{\infty} b_n z^n$ be power series with positive radii of convergence $\rho_P, \rho_Q > 0$. Show that:

- a. $P + Q = \sum_{n=0}^{\infty} (a_n + b_n) z^n$ has radius of convergence $\rho \geq \min(\rho_P, \rho_Q)$.
- b. $PQ = \sum_{n=0}^{\infty} \left(\sum_{i+j=n} a_i b_j \right) z^n$ has radius of convergence $\rho \geq \min(\rho_P, \rho_Q)$.

Show Solution 5.3 on P194

Exercise 5.4

Find a solution to the non-linear differential equation

$$f'(z) + f(z)^2 = 0, \quad f(0) = 1$$

on a disk centered at $z_0 = 0$ by making the ansatz $f(z) = \sum_{n=0}^{\infty} a_n z^n$, inductively determining the coefficients a_n , and finding the radius of convergence.

Show Solution 5.4 on P194

Exercise 5.5

Let $P(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series centered at $z_0 = 0$ and assume that the radius of convergence $\rho > 0$ is positive. Suppose that $P(z) \in \mathbb{R}$ for all $z \in D_\rho(0) \cap \mathbb{R}$. Prove that all coefficients a_n , $n \in \mathbb{N}$, must be real numbers. Deduce that $\overline{P(z)} = P(\bar{z})$ for all $z \in D_\rho(0)$.

Show Solution 5.5 on P196

Exercise 5.6

Prove that the ring of formal power series $\mathbb{C}[[T]]$ is an *integral domain*. In other words, show that $PQ = 0 \implies P = 0$ or $Q = 0$.

Show Solution 5.6 on P197

Exercise 5.7

The *binomial coefficient* of a complex number $\alpha \neq 0$ and $k \in \mathbb{N}$ is defined as

$$\binom{\alpha}{k} = \begin{cases} \frac{\alpha(\alpha-1)\cdots(\alpha-k+1)}{k!} & \text{if } k > 0, \\ 1 & \text{if } k = 0. \end{cases}$$

The *binomial series* is the formal power series

$$B_\alpha = \sum_{k=0}^{\infty} \binom{\alpha}{k} T^k.$$

- Determine the radius of convergence of B_α .
- Prove that if $\alpha \in \mathbb{N}$, then $B_\alpha(z) = (z+1)^\alpha$.
- Show the *generalized Vandermonde* identity for $\alpha, \beta, \alpha + \beta \in \mathbb{C}^\times$,

$$\sum_{k=0}^n \binom{\alpha}{k} \binom{\beta}{n-k} = \binom{\alpha + \beta}{n}. \quad (\star)$$

- Prove that for $\alpha = 1/k$ the complex function $B_\alpha(z)$ satisfies $B_\alpha(z)^k = z+1$ on its domain. Hence $B_\alpha(z)$ is a k -th root of the function $z+1$.

Show Solution 5.7 on P198

Exercise 5.8

For $a > 0$ and $z \in \mathbb{C}$ define $a^z = \exp(\log(a)z)$. Show that:

- a. $a^z b^z = (ab)^z$ for all $a, b > 0, z \in \mathbb{C}$
- b. $a^z a^w = a^{z+w}$ for all $a > 0, z, w \in \mathbb{C}$
- c. $|a^z| = a^{\Re(z)}$ for all $a > 0, z \in \mathbb{C}$

The *Riemann ζ -function* is defined as

$$\zeta(z) = \sum_{n=1}^{\infty} \frac{1}{n^z} \quad \text{for } \Re(z) > 1. \quad (\star)$$

- d. Prove that the series $(\star)$ converges absolutely for all $z \in \mathbb{C}$ with $\Re(z) > 1$ and uniformly on every subset $S_\delta = \{z \in \mathbb{C} \mid \Re(z) > 1 + \delta\}$ with $\delta > 0$.

Show Solution 5.8 on P200

Chapter 6

Contour integrals

In this section, we will generalize the integral $\int_a^b f(x)dx$ from calculus in two ways. Firstly, we allow $f = u + iv$ to be a complex function. Secondly, we will define the integral over more general curves γ than intervals $[a, b] \subset \mathbb{R}$.




Figure 6.1: Generalizing the integral (the area, up to sign) of a real-valued function over an interval γ_1 to a general differentiable curve γ_2 by $\int_a^b f(\gamma(t))|\gamma'(t)|dt$. We will use a *different* generalization where we omit the modulus.

Definition 6.1. Let $f: [a, b] \rightarrow \mathbb{C}$ be a continuous complex-valued function on an interval with real and imaginary parts $f = u + iv$, $u, v: [a, b] \rightarrow \mathbb{R}$. Define

$$\int_a^b f(x)dx = \int_a^b u(x)dx + i \int_a^b v(x)dx. \quad (6.1)$$

Familiar rules for integrals (linearity, substitution) carry over to complex-valued functions, but the following estimate is more tricky to prove.

Proposition 6.1. $\left| \int_a^b f(x)dx \right| \leq \int_a^b |f(x)|dx$

Proof.

Pick a complex number $0 \neq c \in \mathbb{C}$ such that $c \int_a^b f(x)dx \in \mathbb{R}$. Then

$$\begin{aligned} |c| \left| \int_a^b f(x)dx \right| &= \left| c \int_a^b f(x)dx \right| \\ &= \left| \Re \int_a^b cf(x)dx \right| && \text{as } c \int_a^b f(x)dx \in \mathbb{R} \\ &= \left| \int_a^b \Re(cf(x))dx \right| \\ &\leq \int_a^b |\Re(cf(x))|dx && \text{from calculus} \\ &\leq \int_a^b |cf(x)|dx \\ &= |c| \int_a^b |f(x)|dx, \end{aligned}$$

where the third line follows from Equation 6.1 and the fifth from Equation 1.13. The result follows by dividing by $|c| \neq 0$.

□

Definition 6.2. A **curve** (or **path**) in the plane is a continuous map

$$[a, b] \xrightarrow{\gamma} \mathbb{C}$$

on a closed interval. Decompose $\gamma(t) = u(t) + iv(t)$ into real and imaginary parts. The curve γ is **differentiable** if u, v are differentiable on $[a, b]$ (including one-sided derivatives at the endpoints), and the curve γ is **continuously differentiable** (or C^1) if the derivatives $u'(t), v'(t)$ are continuous on $[a, b]$. The curve γ is **piecewise C^1** if there exists a subdivision of the interval

$$a = t_0 < t_1 < \dots < t_n = b \quad (6.2)$$

such that each of the restrictions $\gamma|_{[t_{k-1}, t_k]}$, $k = 1, \dots, n$, is a continuously differentiable curve. In this case we call the subdivision **admissible**. For any admissible subdivision, the **length** of the curve γ is

$$L(\gamma) = \sum_{k=1}^n \int_{t_{k-1}}^{t_k} |\gamma'(t)| dt. \quad (6.3)$$

We call $\gamma([a, b]) \subset \mathbb{C}$ the **image** of the curve. When the image is contained in a subset $D \subset \mathbb{C}$, we say that γ is a **curve in D** .

A curve is **closed** if $\gamma(a) = \gamma(b)$, and then we call $\gamma(a)$ the **base** of the **loop** (or **contour**) γ .




Figure 6.2: A piecewise C^1 curve

Definition 6.3. Let $f: D \rightarrow \mathbb{C}$ be a continuous complex function. Let γ be a piecewise C^1 curve in D . Pick an admissible subdivision Equation 6.2. The **path integral** (or **contour integral** if γ is closed) of f over the curve γ is

$$\int_{\gamma} f(z) dz = \sum_{k=1}^n \int_{t_{k-1}}^{t_k} f(\gamma(t)) \gamma'(t) dt. \quad (6.4)$$

Example 6.1.

The curve $\gamma(t) = t$ parameterizes the interval $[a, b]$. In this case, Equation 6.4 reduces to the integral $\int_a^b f(x) dx$ from Equation 6.1.

Example 6.2.

Let $\gamma(t) = p$ be constant. Then $\int_{\gamma} f(z) dz = 0$ for all f .

Example 6.3 (important).

Let $f(z) = z^n$, $n \in \mathbb{Z}$, and $\gamma(t) = e^{it}$, $t \in [a, b]$.

$$\int_{\gamma} f(z) dz = \int_a^b e^{int} i e^{it} dt = \begin{cases} i(b-a) & \text{if } n = -1, \\ \frac{e^{i(n+1)b} - e^{i(n+1)a}}{n+1} & \text{if } n \neq -1. \end{cases}$$

In particular, for the **boundary curve** of the disk $\overline{D}_r(z_0)$ defined by

$$[0, 2\pi] \xrightarrow{\gamma_{\partial D_r(z_0)}} \overline{D}_r(z_0), \gamma_{\partial D_r(z_0)}(t) = z_0 + r e^{it}, \quad (6.5)$$

which we denote by $\partial D_r(z_0)$, we have

$$\int_{\partial D_r(z_0)} (z - z_0)^n = \begin{cases} 2\pi i & \text{if } n = -1, \\ 0 & \text{if } n \neq -1. \end{cases} \quad (6.6)$$

Lemma 6.1.

- a. The integral Equation 6.4 and the length of a curve Equation 6.3 are independent of the choice of admissible subdivision.
- b. Let $\varphi: [c, d] \rightarrow [a, b]$ be a strictly monotone increasing, continuously differentiable bijection. Let γ be a piecewise C^1 curve. Then $\gamma \circ \varphi$ is also a piecewise C^1 curve and

$$\int_{\gamma \circ \varphi} f(z) dz = \int_{\gamma} f(z) dz.$$

Hence the curve integral is independent of the parameterization of γ . Similarly, $L(\gamma \circ \varphi) = L(\gamma)$ for the lengths.

Proof.

- a. Since every two admissible subdivisions have a common admissible subdivision,

it suffices to prove that Equation 6.4 remains unchanged upon passing to a subdivision. Every subdivision is obtained by inductively inserting points, so it suffices to consider the case of a single insertion, say $t_{k-1} < t_* < t_k$. Then only one summand in Equation 6.4 changes, and our claim follows from

$$\int_{t_{k-1}}^{t_k} f(\gamma(t))\gamma'(t)dt = \int_{t_{k-1}}^{t_*} f(\gamma(t))\gamma'(t)dt + \int_{t_*}^{t_k} f(\gamma(t))\gamma'(t)dt.$$

The same argument applies to the length of a curve.

- b. As φ is bijective, there are unique $s_k \in [c, d]$ with $\varphi(s_k) = t_k$ for $k = 0, \dots, n$. Of course, $s_0 = c$ and $s_n = d$. The monotonicity implies that $s_{k-1} < s_k$ for all k and that φ maps $[s_{k-1}, s_k]$ bijectively onto $[t_{k-1}, t_k]$. The subdivision $c = s_0 < s_1 < \dots < s_n = d$ is again admissible, since $\gamma \circ \varphi|_{[s_{k-1}, s_k]}$ is continuously differentiable for all k by the chain rule.

Using the change of variables $s = \varphi(t)$, we compute

$$\begin{aligned} \int_{\gamma \circ \varphi} f(z)dz &= \sum_{k=1}^n \int_{s_{k-1}}^{s_k} f(\gamma(\varphi(t)))\gamma'(\varphi(t))\varphi'(t)dt \\ &= \sum_{k=1}^n \int_{t_{k-1}}^{t_k} f(\gamma(s))\gamma'(s)ds = \int_{\gamma} f(z)dz. \end{aligned}$$

□

Due to Lemma 6.1(b) we view curves differing only by a parametrization as **equivalent**. In particular, we may translate and scale the domain $[a, b]$.

Definition 6.4.

- a. Let $[a, b] \xrightarrow{\gamma_1} \mathbb{C}$, $[b, c] \xrightarrow{\gamma_2} \mathbb{C}$. Assume $\gamma_1(b) = \gamma_2(b)$. Then the **concatenation**

of γ_1 with γ_2 is the curve

$$[a, c] \xrightarrow{\gamma_1 * \gamma_2} \mathbb{C}, (\gamma_1 * \gamma_2)(t) = \begin{cases} \gamma_1(t) & \text{if } t \in [a, b], \\ \gamma_2(t) & \text{if } t \in [b, c]. \end{cases}$$

b. The **opposite** of a curve $[a, b] \xrightarrow{\gamma} \mathbb{C}$ is the curve

$$[-b, -a] \xrightarrow{-\gamma} \mathbb{C}, (-\gamma)(t) = \gamma(-t).$$

Proposition 6.2. Let $f, g: D \rightarrow \mathbb{C}$ be continuous complex functions and let $[a, b] \xrightarrow{\gamma} D$ be a piecewise C^1 curve.

a. The integral over a curve is linear: for all $\lambda \in \mathbb{C}$ we have

$$\int_{\gamma} \lambda f(z) + g(z) dz = \lambda \int_{\gamma} f(z) dz + \int_{\gamma} g(z) dz.$$

b. For the opposite curve,

$$\int_{-\gamma} f(z) dz = - \int_{\gamma} f(z) dz.$$

c. For the concatenation of curves,

$$\int_{\gamma_1 * \gamma_2} f(z) dz = \int_{\gamma_1} f(z) dz + \int_{\gamma_2} f(z) dz.$$

d. Suppose that $|f(z)| \leq M$ for all $z \in \gamma([a, b])$. Then

$$\left| \int_{\gamma} f(z) dz \right| \leq M \cdot L(\gamma). \quad (6.7)$$

Proof.

a. follows from the linearity of the ordinary integral and b. is simply the change of variables $s = -t$ in the integral. The argument for c. is the same as for Lemma 6.1(a). We prove d. for a trivial subdivision ($n = 1$):

$$\begin{aligned} \left| \int_{\gamma} f(z) dz \right| &= \left| \int_a^b f(\gamma(t)) \gamma'(t) dt \right| \\ &\leq \int_a^b |f(\gamma(t))| |\gamma'(t)| dt \\ &\leq M \int_a^b |\gamma'(t)| dt. \end{aligned}$$

where the second line follows from Proposition 6.1.

The general case is obtained by applying this to each subinterval $[t_{k-1}, t_k]$ and then summing the resulting estimates over k .

□

Remark 6.1. There is a generalization of the path integral to curves γ of bounded variation called the Riemann–Stieltjes integral. Proposition 6.2 continues to hold with the length $L(\gamma)$ replaced by the total variation. For example, Lipschitz continuous maps are of bounded variation.

From Equation 6.7 we obtain:

Corollary 6.1. *Let $(f_n)_{n \in \mathbb{N}}$ be a uniformly convergent sequence of complex functions on D . For every curve $\gamma: [a, b] \rightarrow D$ we have*

$$\lim_{n \rightarrow \infty} \int_{\gamma} f_n(z) dz = \int_{\gamma} \lim_{n \rightarrow \infty} f_n(z) dz.$$

Proposition 6.3 (Complex FTC). *Let f be a continuous complex function on an open set U and let $[a, b] \xrightarrow{\gamma} U$ be a piecewise C^1 curve. Suppose F is a holomorphic function on U with $F' = f$ (we call F a **primitive** of f). Then*

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)). \quad (6.8)$$

In particular, for every closed curve

$$\int_{\gamma} f(z) dz = 0. \quad (6.9)$$

Proof.

By the chain rule,

$$\frac{d}{dt} F(\gamma(t)) = f(\gamma(t)) \gamma'(t).$$

Using the (ordinary) fundamental theorem of calculus to evaluate Equation 6.4 gives

$$\int_{\gamma} f(z) dz = \sum_{k=1}^n F(\gamma(t_k)) - F(\gamma(t_{k-1})) = F(\gamma(t_n)) - F(\gamma(t_0)).$$

□

If F is holomorphic with $F' = 0$ and U is path-connected, see Definition 8.1(a) below, we recover from Equation 6.8 the familiar fact that F is a constant. Of course, this holds more generally for differentiable functions F .

Questions for further discussion

- Is there an analogue of Proposition 6.3 where F is only assumed to be real differentiable?
- Find the integral of a power series $P = \sum_{n=0}^{\infty} a_n z^n$ with positive radius of convergence $\rho > 0$ along an arbitrary curve γ in $D_\rho(0)$?

6.1 Exercises


Exercise 6.1

Recall from Example 4.6 that $\log: \mathbb{C}^- \rightarrow \mathbb{C}$ is a holomorphic primitive of $1/z$ on the slit plane $\mathbb{C}^-$. Combine Equation 6.9 and Example 6.3 to show that there is *no* holomorphic primitive of $1/z$ on the punctured plane $\mathbb{C}^\times$.

Show Solution 6.1 on P201

Exercise 6.2

For the curves γ_k , $k = 1, 2, 3, 4$, as in the following sketch



- find piecewise C^1 parameterisations $\gamma_k: [a, b] \rightarrow \mathbb{C}$,
- compute the length $L(\gamma_k)$ (γ_3 may be omitted)
- evaluate $\int_{\gamma_k} z^2 dz$.

Show Solution 6.2 on P202

Exercise 6.3

Define $\gamma_\epsilon: [-\pi + \epsilon, \pi - \epsilon] \rightarrow \mathbb{C}$, $\gamma_\epsilon(t) = e^{it}$ for $0 < \epsilon < \pi$. Compute

$$\lim_{\epsilon \rightarrow 0} \int_{\gamma_\epsilon} z^{1/2} dz,$$

where the square root $z^{1/2} = \exp(\log(z)/2)$ is defined as on Sheet 5 using the principal branch of the logarithm.

(*Aside:* $z^{1/2}$ is integrable in the sense of measure theory and using the more general integral defined there we can rewrite the above integral as $\int_{\partial D_1(0)} z^{1/2} dz$.)

Show Solution 6.3 on P204

Exercise 6.4

Let γ be the straight line from $z = 1$ to $z = i$. Let $f(z) = 1/z^4$. Determine the maximum of $|f(z)|$ over all $z \in \mathfrak{I}(\gamma)$. Use this to estimate

$$\left| \int_{\gamma} f(z) dz \right| \leq 4\sqrt{2}.$$

Then compute $\int_{\gamma} f(z) dz$ directly and compare this to the estimate.

Show Solution 6.4 on P205

Exercise 6.5

Let $f(z): \partial D_r(z_0) \rightarrow \mathbb{C}$ be a continuous function. Show that

$$\overline{\int_{\partial D_r(z_0)} f(z) dz} = -r^2 \int_{\partial D_r(z_0)} \overline{f(z)} (z - z_0)^{-2} dz.$$

Show Solution 6.5 on P206

Part III

Cauchy's theorem and residues

Chapter 7

Cauchy's theorem

Definition 7.1. Let $D \subset \mathbb{C}$ and $\gamma_0, \gamma_1 : [a, b] \rightarrow D$ curves in D .

- a. A **homotopy**¹ in D between paths γ_0, γ_1 is a continuous map

$$\Gamma : [0, 1] \times [a, b] \longrightarrow D, (s, t) \longmapsto \Gamma_s(t),$$

such that $\Gamma_0(t) = \gamma_0(t)$ and $\Gamma_1(t) = \gamma_1(t)$ for all $t \in [a, b]$. Then γ_0, γ_1 are called (freely) **homotopic paths** in D .

If, additionally, $\Gamma_s(a) = p$ and $\Gamma_s(b) = q$ are constant in $s \in [0, 1]$, we call Γ a **path homotopy** in D and γ_0, γ_1 **path-homotopic** in D .

- b. If γ_0, γ_1 are loops, a **homotopy of loops** in D is a homotopy Γ in D with the additional property that Γ_s is a loop for each $s \in [0, 1]$. Then γ_0, γ_1 are called (freely) **homotopic loops** in D .

A loop is **null-homotopic** in D if there is a homotopy of loops in D to the constant loop.

¹From Greek *homos* 'one and the same' and *topos* 'place, region, space'




Figure 7.1: A path homotopy is a video of paths with fixed endpoints.

Remark 7.1. In the following we will also suppose that Γ is piecewise C^1 , meaning there exist subdivisions

$$0 = s_0 < s_1 < \cdots < s_m = 1, \quad a = t_0 < t_1 < \cdots < t_n = b$$

such that each restriction $\Gamma|_{[s_{j-1}, s_j] \times [t_{k-1}, t_k]}$ is continuously differentiable.

Example 7.1.

A set D is **star-shaped** if there exists a **focal point** $z_0 \in D$ such that for each $z \in D$ the straight line segment $tz + (1 - t)z_0$, $t \in [0, 1]$, is contained in D . Disks, the complex plane, and rectangles are star-shaped. Every loop γ in a star-shaped domain is null-homotopic. If the focal point agrees with the base of the loop, the path homotopy is

$$\Gamma_s(t) = (1 - s)\gamma(t) + sz_0. \quad (7.1)$$

In general, the path homotopy is more complicated to write down. We omit it, since we will not need this fact below.




Figure 7.2: Augustin Louis, Baron Cauchy. Lithograph by J. Boilly, 1821. [Wellcome Collection](#). Public Domain Mark

Theorem 7.1 (Cauchy's theorem). *Let $f: U \rightarrow \mathbb{C}$ be a holomorphic function on an open set. Let γ_0, γ_1 be piecewise C^1 curves in U that are path-homotopic in U . Then*

$$\int_{\gamma_0} f(z)dz = \int_{\gamma_1} f(z)dz.$$

Proof.

Pick a path homotopy $\Gamma: [0, 1] \times [a, b] \rightarrow U$. By reparameterizing we may assume $[a, b] = [0, 1]$. Write $R^{(0)} = [0, 1] \times [0, 1]$ for the domain of Γ and let $\partial R^{(0)}$ be its piecewise linear boundary path, with the obvious counterclockwise parameterization (see Figure 7.3).




Figure 7.3: Path homotopy when γ_0, γ_1 are loops.

The boundary path has four segments and the path integral along each constant path vanishes. Using Proposition 6.2(b),(c), we then find that

$$\int_{\gamma_0} f(z)dz - \int_{\gamma_1} f(z)dz = \int_{\Gamma(\partial R^{(0)})} f(z)dz.$$

Subdivide $R^{(0)}$ into four congruent rectangles $R_i^{(0)}$ as in Figure 7.4. Using the fact that the path integrals in opposite directions cancel, we find

$$\left| \int_{\Gamma(\partial R^{(0)})} f(z)dz \right| = \left| \sum_{i=1}^4 \int_{\Gamma(\partial R_i^{(0)})} f(z)dz \right| \leq \sum_{i=1}^4 \left| \int_{\Gamma(\partial R_i^{(0)})} f(z)dz \right|.$$




Figure 7.4: Subdivision and cancellation of opposite paths

Let $R^{(1)}$ be the rectangle $R_i^{(0)}$ for which $\left| \int_{\Gamma(\partial R_i^{(0)})} f(z) dz \right|$ is maximal. Then

$$\left| \int_{\Gamma(\partial R^{(0)})} f(z) dz \right| \leq 4 \left| \int_{\Gamma(\partial R^{(1)})} f(z) dz \right|.$$

Now repeat this process with $R^{(1)}$ to obtain $R^{(2)}$ and so forth. This yields a sequence of rectangles $R^{(n)}$ as in Figure 7.5 with sides of length 2^{-n} and

$$\left| \int_{\Gamma(\partial R^{(0)})} f(z) dz \right| \leq 4^n \left| \int_{\Gamma(\partial R^{(n)})} f(z) dz \right|. \quad (7.2)$$




Figure 7.5: Convergent sequence of rectangles $\bigcap R^{(n)} = \{(s_0, t_0)\}$

As the side lengths tend to zero, the midpoints $(s^{(n)}, t^{(n)})$ of the rectangles $R^{(n)}$ are a Cauchy sequence, so converge to some limit (s_0, t_0) . Let $z_0 = \Gamma_{s_0}(t_0)$. Since f is complex differentiable at z_0 , we may write

$$f(z) = f(z_0) + (z - z_0)f'(z_0) + (z - z_0)\rho(z), \quad \lim_{z \rightarrow z_0} \rho(z) = 0. \quad (7.3)$$

By Example 6.4,

$$\int_{\Gamma(\partial R^{(n)})} (f(z_0) + (z - z_0)f'(z_0)) dz = 0. \quad (7.4)$$

As Γ is piecewise C^1 , its derivative is bounded in norm by some $C > 0$.

We may also estimate Γ in a neighbourhood of (s_0, t_0) as

$$\begin{aligned}\Gamma(s, t) &= \Gamma(s_0, t_0) + ((s, t) - (s_0, t_0)) \Gamma'(s_0, t_0) + ((s, t) - (s_0, t_0)) \phi(s, t), \\ \lim_{(s, t) \rightarrow (s_0, t_0)} \phi(s, t) &= 0.\end{aligned}$$

Let $\epsilon > 0$. Pick $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, $z \in \Gamma(R^{(n)})$, $(s, t) \in R^{(n)}$, we have $|\rho(z)|, |\phi(s, t)| < \epsilon$. As the side lengths of $R^{(n)}$ are 2^{-n} and $z_0 \in \Gamma(R^{(n)})$, for all $z \in \Gamma(\partial R^{(n)})$, we have

$$\begin{aligned}|z - z_0| &= |\Gamma(s, t) - \Gamma(s_0, t_0)| \\ &\leq |(s, t) - (s_0, t_0)| |\Gamma'(s_0, t_0)| + |(s, t) - (s_0, t_0)| |\phi(s, t)| \\ &\leq 2^{-n} \sqrt{2} C + 2^{-n} \sqrt{2} \epsilon \\ &\leq \sqrt{2} 2^{-n} (C + \epsilon)\end{aligned}$$

where $(s, t) \in \partial(R^{(n)})$ is any point in $\Gamma^{-1}(z)$

Combining this with Equation 7.3 and Equation 7.4, we can estimate

$$\begin{aligned}\left| \int_{\Gamma(\partial R^{(n)})} f(z) dz \right| &= \left| \int_{\Gamma(\partial R^{(n)})} (z - z_0) \rho(z) dz \right| \\ &\leq L(\Gamma(\partial R^{(n)})) \sqrt{2} 2^{-n} (C + \epsilon) \epsilon = 4C \cdot 2^{-n} \sqrt{2} 2^{-n} (C + \epsilon) \epsilon\end{aligned}$$

The last inequality follows by Equation 6.7.

Hence

$$\left| \int_{\gamma} f(z) dz \right| = \left| \int_{\Gamma(\partial R^{(0)})} f(z) dz \right| \leq 4C(C + \epsilon)\sqrt{2}\epsilon.$$

The final inequality follows from Equation 7.2.

As $\epsilon > 0$ is arbitrary, the left hand side must be zero.

□

Theorem 7.2. *Let γ_0, γ_1 be piecewise C^1 loops in U that are (freely) homotopic in U . Suppose $f: U \rightarrow \mathbb{C}$ is a holomorphic function. Then*

$$\int_{\gamma_0} f(z) dz = \int_{\gamma_1} f(z) dz.$$

In particular, if γ is a loop that is homotopic in U to the constant loop, then

$$\int_{\gamma} f(z) dz = 0. \tag{7.5}$$

Proof.

Let Γ be the homotopy between γ_0 and γ_1 . Let $\eta(s) = \Gamma_s(0)$. Figure 7.6 describes the construction of a path homotopy between $\eta * \gamma_0 * (-\eta)$ and γ_1 .




Figure 7.6: Construction of path homotopy

Hence Theorem 7.1 implies our claim, using the fact that the path integrals over η and $-\eta$ cancel each other.

□

Questions for further discussion

- How should Figure 7.6 be interpreted? Can you give a formula for the path homotopy (consider cases)?
- Give a counterexample to Cauchy's theorem when (i) γ is not null-homotopic
(ii) $f(z)$ is not holomorphic
- Does Cauchy's theorem hold for $f(z) = \bar{z}$?

7.1 Exercises

Exercise 7.1

a. Sketch the curves

$$\begin{aligned}\gamma_0: [-1, 1] &\longrightarrow \mathbb{C}, & \gamma_0(t) &= t, \\ \gamma_1: [-1, 1] &\longrightarrow \mathbb{C}, & \gamma_1(t) &= e^{i\pi \frac{1-t}{2}}\end{aligned}$$

and show that they are path-homotopic.

b. Let $0 < r_0 < r_1$ and $z_0 \in \mathbb{C}$. Prove that the loops $\partial D_{r_0}(z_0)$ and $\partial D_{r_1}(z_0)$ are freely homotopic in the closed annulus $\overline{A}_{r_0, r_1}(z_0)$.

Show Solution 7.1 on P207

Exercise 7.2

Let $\alpha + i\beta \in \mathbb{C}$. Determine

$$\int_a^b e^{(\alpha+i\beta)t} dt$$

to compute

$$\int_a^b e^{\alpha t} \cos(\beta t) dt.$$

Show Solution 7.2 on P208

Exercise 7.3

Let $\mathbb{C}^- = \mathbb{C} \setminus (-\infty, 0]$ be the slit plane.

- Show that any two points in $\mathbb{C}^-$ may be connected by a path in $\mathbb{C}^-$. Hence $\mathbb{C}^-$ is *path-connected*.
- Show that every closed curve $\gamma: [a, b] \rightarrow \mathbb{C}^-$ is null-homotopic in $\mathbb{C}^-$ by finding a homotopy

$$\Gamma: [0, 1] \times [a, b] \longrightarrow \mathbb{C}^-, (s, t) \longmapsto \Gamma_s(t)$$

satisfying $\Gamma_0(t) = \gamma(t)$ and $\Gamma_1(t) = 1$ for all $t \in [a, b]$. Hence $\mathbb{C}^-$ is *simply-connected*.

- Prove the analogues of a. and b. for a disk $D_r(z_0)$.
- Use Cauchy's Theorem to prove that the punctured plane $\mathbb{C}^\times = \mathbb{C} \setminus \{0\}$ is not simply-connected. That is, there exists a closed curve in $\mathbb{C}^\times$ that is not null-homotopic in $\mathbb{C}^\times$.

Show Solution 7.3 on P209

Exercise 7.4

Let $0 < b < 1$.

- a. Using the geometric series, find the power series expansion

$$\frac{1}{z - 1/b} = \sum_{n=0}^{\infty} a_n (z - b)^n$$

with center $z_0 = b$ and determine the radius of convergence ρ .

- b. Use a. to show that for all $0 < r < \rho$

$$\int_{\partial D_r(b)} \frac{dz}{(z - b)(z - 1/b)} = \frac{2\pi i}{b - 1/b}.$$

- c. Use b. to compute

$$\int_0^{2\pi} \frac{dt}{1 - 2b \cos(t) + b^2}.$$

Show Solution 7.4 on P211

Exercise 7.5

Let $\gamma: [0, 1] \rightarrow D$ be a curve in $D \subset \mathbb{C}$ and let $-\gamma: [0, 1] \rightarrow D$, $(-\gamma)(t) = \gamma(1 - t)$ be the opposite curve. Prove that $\gamma * (-\gamma)$ is path-homotopic to a constant loop.

Show Solution 7.5 on P213

Chapter 8

Applications of Cauchy's theorem

Theorem 8.1 (Cauchy's integral formula). *Let $f: U \rightarrow \mathbb{C}$ be a holomorphic function and assume that $\overline{D}_r(z_0) \subset U$ with boundary curve*

$$[0, 2\pi] \xrightarrow{\gamma_{\partial D_r(z_0)}} \overline{D}_r(z_0), \gamma_{\partial D_r(z_0)}(t) = z_0 + re^{it}.$$

Then

$$f(z) = \frac{1}{2\pi i} \int_{\partial D_r(z_0)} \frac{f(\zeta)}{\zeta - z} d\zeta, \quad \forall z \in D_r(z_0). \quad (8.1)$$

Proof.

Fix $z \in D_r(z_0)$. Since f is complex differentiable at z , we have

$$f(\zeta) = f(z) + (\zeta - z)f'(z) + (\zeta - z)\rho(\zeta - z), \lim_{\zeta \rightarrow z} \rho(\zeta - z) = 0. \quad (8.2)$$

There is a homotopy in $U \setminus \{z\}$ between the loops $\partial D_r(z_0)$ and $\partial D_s(z)$, for all $s > 0$ with $D_s(z) \subset D_r(z_0)$, see Figure 8.1.




Figure 8.1: Homotopy between $\partial D_r(z_0)$ and $\partial D_s(z)$

Moreover, $\zeta \mapsto \frac{f(\zeta)}{\zeta - z}$ is a holomorphic function on $U \setminus \{z\}$.

By Theorem 7.1 we have

$$\int_{\partial D_r(z_0)} \frac{f(\zeta)}{\zeta - z} d\zeta = \int_{\partial D_s(z)} \frac{f(\zeta)}{\zeta - z} d\zeta,$$

applying Equation 8.2,

$$\int_{\partial D_s(z)} \frac{f(\zeta)}{\zeta - z} d\zeta = \int_{\partial D_s(z)} \frac{f(z)}{\zeta - z} d\zeta + \int_{\partial D_s(z)} f'(z) d\zeta + \int_{\partial D_s(z)} \rho(\zeta - z) d\zeta$$

applying Equation 6.6 first with $n = -1$ and then $n = 0$, we have

$$\int_{\partial D_s(z)} \frac{f(z)}{\zeta - z} d\zeta + \int_{\partial D_s(z)} f'(z) d\zeta + \int_{\partial D_s(z)} \rho(\zeta - z) d\zeta = 2\pi i f(z) \quad (8.3)$$

$$+ 0 \quad (8.4)$$

$$+ \int_{\partial D_s(z)} \rho(\zeta - z) d\zeta. \quad (8.5)$$

Consequently,

$$\int_{\partial D_r(z_0)} \frac{f(\zeta)}{\zeta - z} d\zeta = 2\pi i f(z) + 0 + \int_{\partial D_s(z)} \rho(\zeta - z) d\zeta.$$

Since $\lim_{\zeta \rightarrow z} \rho(\zeta - z) = 0$ there exists a bound $|\rho(\zeta - z)| \leq M$. Hence the final term can be estimated using Equation 6.7 as

$$\left| \int_{\partial D_s(z)} \rho(\zeta - z) d\zeta \right| \leq 2\pi s M.$$

This tends to zero as $s \rightarrow 0$, and the left hand side is independent of s .

□




Figure 8.2: Boundary curve of largest disk fitting inside U

Theorem 8.2 (Higher Cauchy integral formula). *Let $f: U \rightarrow \mathbb{C}$ be a holomorphic function and assume that $\overline{D}_r(z_0) \subset U$. Then f is equal on $D_r(z_0)$ to its Taylor power series*

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n, \quad (8.6)$$

which has radius of convergence $\rho \geq r$. In particular, f is infinitely complex differentiable on the open set U and has a primitive on $D_r(z_0)$. Moreover,

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \int_{\partial D_r(z_0)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta. \quad (8.7)$$

Proof.

Let $z \in D_r(z_0)$. The integrand in Equation 8.1 can be rewritten as

$$\frac{f(\zeta)}{\zeta - z} = \frac{\frac{f(\zeta)}{\zeta - z_0}}{1 - \frac{z - z_0}{\zeta - z_0}} = \frac{f(\zeta)}{\zeta - z_0} \sum_{n=0}^{\infty} \left(\frac{z - z_0}{\zeta - z_0} \right)^n, \quad (8.8)$$

where the geometric series converges uniformly for all ζ with $|z - z_0| < |\zeta - z_0|$.




Figure 8.3: The region $|z - z_0| < |\zeta - z_0|$ and the contour of integration

In particular, this holds for all $\zeta \in \partial D_r(z_0)$ and using Corollary 6.1 we may exchange the limit and the integral. Hence for all $|z - z_0| < r$ we have a convergent series

$$f(z) = \sum_{n=0}^{\infty} \left(\frac{1}{2\pi i} \int_{\partial D_r(z_0)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta \right) (z - z_0)^n. \quad (8.9)$$

In particular, the series Equation 8.9 has radius of convergence $\rho \geq r$.

Finally, Equation 8.7 (and therefore Equation 8.6) follow from the fact that the power series Equation 8.9 is differentiated termwise, see Theorem 5.2.

□

Theorem 8.3 (Liouville). *Every bounded entire function is constant.*

Proof.

Suppose $|f(z)| < C$ for all $z \in \mathbb{C}$. Then by Equation 8.7 with $z_0 = 0$ we have

$$|f^{(n)}(0)| = \left| \frac{n!}{2\pi i} \int_{\partial D_r(0)} \frac{f(\zeta)}{\zeta^{n+1}} d\zeta \right|,$$

by Equation 6.7,

$$\left| \frac{n!}{2\pi i} \int_{\partial D_r(0)} \frac{f(\zeta)}{\zeta^{n+1}} d\zeta \right| \leq \frac{n!}{2\pi} L(\partial D_r(0)) \frac{C}{r^{n+1}} = \frac{n!C}{r^n}$$

and so

$$|f^{(n)}(0)| \leq \frac{n!C}{r^n}$$

for all $r > 0$. Letting $r \rightarrow \infty$, this implies that $f^{(n)}(0) = 0$ for all $n > 0$. Now Equation 8.6 proves that $f(z)$ is constant.

□



Figure 8.4: Carl Friedrich Gauß, 1777-1855, [Österreichische Nationalbibliothek](#). Public Domain

Theorem 8.4 (Fundamental theorem of algebra). *Every non-constant polynomial $P(z) = a_n z^n + \dots + a_1 z + a_0$ with $a_i \in \mathbb{C}$ has a complex root.*

Proof.

Assume by contradiction that $P(z) \neq 0$ for all $z \in \mathbb{C}$. Then $1/P(z)$ is holomorphic and bounded. Indeed, assuming $a_n \neq 0$, we find

$$|P(z)| \geq |a_n||z|^n - |a_{n-1}||z|^{n-1} - \dots - |a_0| \rightarrow +\infty \text{ as } z \rightarrow \infty.$$

Hence $1/|P(z)| \rightarrow 0$ as $z \rightarrow \infty$. In particular, $|1/P(z)|$ is bounded.

□

Definition 8.1.

- a. A subset $D \subset \mathbb{C}$ is **path-connected** if for all $z_0, z_1 \in D$ there exists a (piecewise C^1) curve γ in D with $\gamma(0) = z_0, \gamma(1) = z_1$.
- b. A path-connected subset D is **simply connected** if every loop in D is (freely) homotopic in D to a constant loop.

Example 8.1.

Every star-shaped set D is simply connected. Firstly, D is path-connected since any $z \in D$ can be connected by a straight line $tz + (1 - t)z_0$ to the focal point z_0 . Secondly, if $\gamma: [a, b] \rightarrow D$ is a loop, then $\Gamma_s(t) = sz_0 + (1 - s)\gamma(t)$ is a homotopy in D from γ to the constant loop.

Theorem 8.5. *Let $f: U \rightarrow \mathbb{C}$ be a holomorphic function. Suppose that U is simply connected and let $z_0 \in U$. Define $F(z)$ for each $z \in U$ by choosing a piecewise C^1 curve γ with $\gamma(0) = z_0, \gamma(1) = z$ and defining*

$$F(z) = \int_{\gamma} f(\zeta) d\zeta. \quad (8.10)$$

Then F is well-defined and is the unique holomorphic function on U such that $F' = f$ and $F(z_0) = 0$.

Proof.

For any two curves γ_0, γ_1 as in the statement of the theorem, the concatenation $\gamma_0 * (-\gamma_1)$ is a loop which is null-homotopic by assumption. By Theorem 7.2 we have

$$\int_{\gamma_0} f(\zeta) d\zeta - \int_{\gamma_1} f(\zeta) d\zeta = 0,$$

hence Equation 8.10 is well-defined. It remains to check that F is complex differentiable

at every point $z_* \in U$ for which it suffices to restrict attention to the disk $\overline{D}_r(z_*) \subset U$.

Fix a path γ_* in U from z_0 to z_* . For each $z \in D_r(z_*)$, let γ be the concatenation of γ_* with a path η from z_* to z in the disk $D_r(z_*)$. Then

$$F(z) = \int_{\gamma} f(\zeta) d\zeta = \int_{\gamma_*} f(\zeta) d\zeta + \int_{\eta} f(\zeta) d\zeta.$$

The first integral is a constant independent of z . By Theorem 8.2, f has a holomorphic primitive g on $D_r(z_*)$ and η is a path in that disk, so we can apply the complex FTC Equation 6.8 to the second integral and write

$$F(z) = \int_{\gamma_*} f(\zeta) d\zeta + g(z) - g(z_*).$$

This shows that F is a holomorphic function of z with $F' = g' = f$.

To prove uniqueness, suppose that $\tilde{F}$ is another holomorphic function with $\tilde{F}' = f$ and $\tilde{F}(z_0) = 0$. Then $F - \tilde{F} = c$ is constant on U and by evaluating at z_0 we find $c = F(z_0) - \tilde{F}(z_0) = 0 - 0 = 0$.

□

Example 8.2.

Let $U \subset \mathbb{C}$ be a simply connected subset with $0 \notin U$ and $f(z) = 1/z$. Fix $w_0 \in \mathbb{C}$ such that $z_0 = e^{w_0} \in U$. Define the holomorphic function $F(z)$ on U by Equation 8.10. Then $\ell(z) = F(z) + w_0$ is the unique **branch of the logarithm** on U such that $\ell(z_0) = w_0$.

For $z_0 = 1$, $w_0 = 0$, and $U = \mathbb{C}^-$, we recover the principal branch of the logarithm, since $\log'(z) = 1/z$ by Equation 4.9 and $\log(1) = 0$.

Questions for further discussion

- How should $\int_{-i}^{1+i} z^2 dz$ be interpreted? What is the result?
- Why is $\mathbb{C}^\times$ path-connected but not simply connected?

Hint: consider $\frac{1}{2\pi i} \int_{\gamma} \frac{1}{z} dz$ for a loop γ in $\mathbb{C}^\times$. +

It is a mysterious calculus fact that $\arctan(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}$ diverges for $|x| > 1$ although $\arctan: \mathbb{R} \rightarrow \mathbb{R}$ is smooth. In terms of the principal logarithm, $\arctan(z) = \frac{i}{2} \log\left(\frac{i+z}{i-z}\right)$. Apply this to give a geometric explanation of the divergence using Theorem 8.2.

8.1 Exercises

Exercise 8.1

Compute the following integrals:

- $\int_{\partial D_1(0)} \frac{\sin(z)}{z^2} dz,$
- $\int_{\partial D_{1/2}(0)} \frac{\cos(z)}{z-1} dz,$
- $\int_{\partial D_2(1)} \frac{\sin(\cos(z))}{z-1} dz.$

Show Solution 8.1 on P214

Exercise 8.2

Let $f: \mathbb{C} \rightarrow \mathbb{C}$ be entire. Suppose there is a constant $C > 0$ such that $|f(z)| \leq C|z|^d$ for all $z \in \mathbb{C}$. Prove that $f(z)$ is a polynomial of degree $\leq d$.

Hint: Generalize the proof of Liouville's theorem.

Show Solution 8.2 on P214

Exercise 8.3

Let $f: U \rightarrow \mathbb{C}$ be holomorphic and $\overline{D}_r(z_0) \subset U$. Show that

$$f(z_0) = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{i\varphi}) d\varphi$$

and interpret this equation geometrically.

Show Solution 8.3 on P215

Exercise 8.4

Let $f: U \rightarrow \mathbb{C}$ be holomorphic and $\overline{D}_r(z_0) \subset U$.

- Prove that $g(x) = f(z_0 + re^{ix})$ is a 2π -periodic function $\mathbb{R} \rightarrow \mathbb{C}$.
- Show that the Cauchy integral formula implies an absolutely and uniformly convergent *Fourier expansion*

$$g(x) = \sum_{n=0}^{\infty} \gamma_n e^{inx}, \quad \forall x \in \mathbb{R},$$

with only non-negative Fourier modes. Moreover, show that

$$\gamma_n = \frac{1}{2\pi} \int_0^{2\pi} g(x) e^{-inx} dx.$$

Note. More generally, a bi-infinite Fourier series $\sum_{n=-\infty}^{\infty} \gamma_n e^{inx}$ can be obtained as a superposition $g_1(x) + g_2(-x)$.

Show Solution 8.4 on P216

Exercise 8.5

Let $f: U \rightarrow \mathbb{C}$ be holomorphic and $\overline{D}_r(z_0) \subset U$. Using polar coordinates show that for the (surface) integral over the unit disk $\overline{D}_r(x_0) = \{(x, y) \in \mathbb{R}^2 \mid (x - x_0)^2 + (y - y_0)^2 \leq r^2\}$ we have

$$f(z_0) = \frac{1}{\pi r^2} \int_{\overline{D}_r(z_0)} f(x + iy) dx dy$$

and interpret this equation geometrically.

Show Solution 8.5 on P217

Chapter 9

Laurent series and singularities

Definition 9.1. A **formal bilateral series** is an expression of the form

$$P = \sum_{n=-\infty}^{\infty} a_n T^n \quad (9.1)$$

with **coefficients** $a_n \in \mathbb{C}$. This is just a bi-infinite sequence $(a_n)_{n \in \mathbb{Z}}$ of complex numbers. Equivalently, we have a pair of formal power series

$$P_+ = \sum_{n=0}^{\infty} a_n T^n, \quad P_- = \sum_{n=1}^{\infty} a_{-n} T^n, \quad (9.2)$$

such that

$$P = P_+(T) + P_-(T^{-1}).$$

P_- is called the **principal part**. The **residue** of P is the coefficient a_{-1} .

Definition 9.2. Let P be a bilateral series Equation 9.1. The **domain** $D(P)$ is the

set of all $z \in \mathbb{C} \setminus \{0\}$ such that both series of complex numbers

$$P_+(z) = \sum_{n=0}^{\infty} a_n z^n, \quad P_-(z^{-1}) = \sum_{n=1}^{\infty} a_{-n} z^{-n}$$

converge. We obtain a complex function

$$D(P) \longrightarrow \mathbb{C}, P(z) := \sum_{n=-\infty}^{\infty} a_n z^n := P_+(z) + P_-(z^{-1}). \quad (9.3)$$

More generally, fix a **center** $z_0 \in \mathbb{C}$. We then have a complex function

$$P(z - z_0) := \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n := P_+(z - z_0) + P_-(z - z_0)^{-1}.$$

with domain $D(P, z_0) = z_0 + D(P)$ differing from $P(z)$ only by a translation.

Remark 9.1. Bi-infinite series of complex numbers $\sum_{n \in \mathbb{Z}} a_n$ must be treated with care. We have avoided these issues by viewing them as pairs of ordinary series. Another approach would be to pick a bijection $\nu: \mathbb{N} \rightarrow \mathbb{Z}$ and consider the ordinary series $\sum_{k=0}^{\infty} a_{\nu(k)}$. However, the limit depends on the choice of ν unless the series is absolutely convergent.

Definition 9.3. The **open annulus** $A_{r,R}(z_0)$ centered at $z_0 \in \mathbb{C}$ with radii $0 \leq r, R \leq +\infty$ is the (possibly empty) subset

$$A_{r,R}(z_0) = \{z \in \mathbb{C} \mid r < |z - z_0| < R\} = D_R(z_0) \cap (\mathbb{C} \setminus \overline{D}_r(z_0)).$$

The **punctured open disk** centered at $z_0 \in \mathbb{C}$ with radius $0 < R \leq +\infty$ is

$$D_R^\times(z_0) = D_R(z_0) \setminus \{z_0\} = A_{0,R}(z_0).$$

The **closed annulus** is $\overline{A}_{r,R}(z_0) = \overline{D}_R(z_0) \cap (\mathbb{C} \setminus D_r(z_0))$.




Figure 9.1: A closed annulus and a punctured open disc centered at z_0

Using the inversion

$$\iota: \mathbb{C}^\times \longrightarrow \mathbb{C}^\times, z \longmapsto w = z^{-1}, \quad (9.4)$$

we can rewrite Equation 9.3 as $P(z) = P_+(z) + P_-(w)$. Therefore we can treat the principal part as a power series as well, but on a second complex w -plane corresponding to the original complex z -plane via Equation 9.4.

Lemma 9.1. *Let $0 \leq r \leq +\infty$. Then*

$$\iota^{-1}(D_r(0)) = \mathbb{C}^\times \setminus \overline{D}_{1/r}(0), \quad \iota^{-1}(\overline{D}_r(0)) = \mathbb{C}^\times \setminus D_{1/r}(0), \quad (9.5)$$

using the conventions $1/0 = +\infty$ and $1/+\infty = 0$.

Proposition 9.1. *Let P be a bilateral series. If $D(P, z_0) \neq \emptyset$, there are unique **radii of convergence** $0 \leq r, R \leq +\infty$ such that*

$$A_{r,R}(z_0) \subset D(P) \subset \overline{A}_{r,R}(z_0). \quad (9.6)$$

Here $r = 1/\rho_-$, $R = \rho_+$ for the radii of convergence $\rho_\pm$ of the power series $P_\pm$ as defined in Theorem 5.1.

Moreover, the series $P_+(z)$, $P_-(1/z)$ converge absolutely and uniformly on every annulus $A_{r',R'}(z_0)$ with $r < r' < R' < R$.

Proof.

For simplicity, put $z_0 = 0$. According to Theorem 5.1 the power series P_+, P_- have radii of convergence $\rho_\pm \geq 0$ and disklike domains

$$D_{\rho_\pm}(0) \subset D(P_\pm) \subset \overline{D}_{\rho_\pm}(0). \quad (9.7)$$

As $P(z) = P_+(z) + P_-(w)$ with $\iota(z) = w$, we can rewrite Definition 9.2 in terms of the preimage operation $\iota^{-1}(\sqcup)$ as

$$D(P) = D(P_+) \cap \iota^{-1}(D(P_-)).$$

Applying ι^{-1} to Equation 9.7 for P_- and using Equation 9.5 gives

$$\mathbb{C}^\times \setminus \overline{D}_{1/\rho_-}(0) \subset \iota^{-1}(D(P_-)) \subset \mathbb{C}^\times \setminus D_{1/\rho_-}(0).$$

Combining this with Equation 9.7 for P_+ implies

$$\underbrace{(\mathbb{C}^\times \setminus \overline{D}_{1/\rho_-}(0)) \cap D_{\rho_+}(0)}_{A_{r,R}(0)=} \subset D(P) \subset \underbrace{(\mathbb{C}^\times \setminus D_{1/\rho_-}(0)) \cap \overline{D}_{\rho_+}(0)}_{\subset \overline{A}_{r,R}(0)}$$

where we have set $r = 1/\rho_-$, $R = \rho_+$ (see Figure 9.2).




Figure 9.2: Annulus of convergence for a bilateral series. The series $P_+(z)$ converges on the red disk in the z -plane and the series $P_-(w)$ on the blue disk in the w -plane, corresponding to the complement of a disk on the z -plane. $P(z) = P_+(z) + P_-(w)$ converges on the overlap, an annulus in the z -plane.

The absolute and uniform convergence on smaller annuli follows from the corresponding properties for power series stated in Theorem 5.1.

□

If the domain of a bilateral series P with center z_0 contains a non-empty open annulus $A_{r,R}(z_0) \subset D(P, z_0)$, then P is called a (convergent) **Laurent series** at z_0 .



Figure 9.3: Pierre Laurent, 1813-1854, [Wikipedia](#). Public Domain

Theorem 9.1 (Laurent expansion). *Let $f: U \rightarrow \mathbb{C}$ be a holomorphic function on an open set containing a closed annulus $\overline{A}_{r,R}(z_0)$. Then f is equal on $A_{r,R}(z_0)$ to the convergent Laurent series*

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n.$$

Moreover,

$$a_n = \frac{1}{2\pi i} \int_{\partial D_r(z_0)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta \quad (9.8)$$

Proof.

Let $z \in A_{r,R}(z_0)$. Let γ be the contour shown in Figure 9.4 that runs around and connects the two boundary components of the annulus. Pick $s > 0$ such that $D_s(z) \subset A_{r,R}(z_0)$. Then γ is freely homotopic to $\partial D_s(z)$.




Figure 9.4: Contour of integration for Laurent expansion

In the path integral over γ , the pieces connecting the two boundary components cancel and the inner circle is negatively oriented. Hence, by Equation 8.1

$$f(z) = \frac{1}{2\pi i} \int_{\partial D_s(z)} \frac{f(\zeta)}{\zeta - z} d\zeta,$$

applying Theorem 7.2

$$\frac{1}{2\pi i} \int_{\partial D_s(z)} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\partial A_{r,R}(z_0)} \frac{f(\zeta)}{\zeta - z} d\zeta,$$

thus,

$$f(z) = \frac{1}{2\pi i} \int_{\partial D_R(z_0)} \frac{f(\zeta)}{\zeta - z} d\zeta - \frac{1}{2\pi i} \int_{\partial D_r(z_0)} \frac{f(\zeta)}{\zeta - z} d\zeta$$

Rewrite the first integral as in Equation 8.8,

$$\frac{f(\zeta)}{\zeta - z} = \frac{\frac{f(\zeta)}{\zeta - z_0}}{1 - \frac{z - z_0}{\zeta - z_0}} = \frac{f(\zeta)}{\zeta - z_0} \sum_{n=0}^{\infty} \left(\frac{z - z_0}{\zeta - z_0} \right)^n.$$

The geometric series converges uniformly because $|z - z_0| < |\zeta - z_0| = R$. We then apply Corollary 6.1 to exchange the limit and the integral. We similarly rewrite the second integral using

$$\frac{f(\zeta)}{z - \zeta} = \frac{\frac{f(\zeta)}{z - z_0}}{1 - \frac{\zeta - z_0}{z - z_0}} = \frac{f(\zeta)}{z - z_0} \sum_{n=0}^{\infty} \left(\frac{\zeta - z_0}{z - z_0} \right)^n$$

and again apply Corollary 6.1, now using $r = |\zeta - z_0| < |z - z_0|$.

□

Definition 9.4. Let $f: U \rightarrow \mathbb{C}$ be a holomorphic function. A point $z_0 \in \mathbb{C} \setminus U$ is an **isolated singularity** of f if there exists $R > 0$ such that $D_R^\times(z_0) \subset U$.

By applying Theorem 9.1 to $A_{r,R}(z_0)$ for all $0 < r < R$ we obtain a Laurent expansion

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n \tag{9.9}$$

that converges on $D_R^\times(z_0)$.

Definition 9.5. An isolated singularity of f is called

- a. **removable**,
- b. a **pole** of order $m \geq 1$,
- c. **essential**,

if the principal part $\sum_{n=-\infty}^{-1} a_n (z - z_0)^n$ of the Laurent expansion Equation 9.9

- a. is zero,
- b. has $a_{-m} \neq 0$ and $a_n = 0$ for all $n < -m$,
- c. has infinitely many non-zero terms.

We can restate Definition 9.5 without reference to Laurent expansions. If z_0 is a removable singularity, then Equation 9.9 is a power series which by Theorem 5.1 defines a holomorphic $D_r(z_0)$. In other words,

$$z_0 \text{ removable singularity} \iff f \text{ extends holomorphically to } U \cup \{z_0\}.$$

If z_0 is a pole of order m , then $(z - z_0)^m f(z)$ has a removable singularity. So $g(z) = (z - z_0)^m f(z)$ is a holomorphic function on $U \cup \{z_0\}$ and

$$z_0 \text{ pole of order } m \iff f(z) = \frac{g(z)}{(z - z_0)^m}, \quad g \text{ holomorphic on } U \cup \{z_0\} \text{ with } g(z_0) \neq 0.$$

Theorem 9.2 (Riemann removable singularities). *Let z_0 be an isolated singularity of a holomorphic function $f: U \rightarrow \mathbb{C}$. Suppose there exist $R, C > 0$ such that $|f(z)| < C$ for all $z \in D_R^\times(z_0)$. Then z_0 is a removable singularity.*

Proof.

For all $n < 0$ we can estimate Equation 9.8 using Equation 6.7 and get

$$|a_n| \leq \frac{1}{2\pi} L(\partial D_R(z_0)) \frac{C}{R^{n+1}} = CR^{-n} \xrightarrow{R \rightarrow 0} 0.$$

□

Questions for further discussion

- Define the sum of formal Laurent series in the same way as Equation 5.2. Why can't we use Equation 5.3 to define the product of formal Laurent series?
- In Definition 9.2, why did we exclude $z = z_0$?
- For an isolated singularity, why is $U \cup \{z_0\}$ always an open set?
- Does the converse of the Riemann removable singularities theorem hold?

9.1 Exercises

Exercise 9.1

Determine whether the singularities of the following holomorphic functions $f(z)$ are isolated. If so, compute the residue. When applicable, find also the order of the pole or a holomorphic extension.

- i. $\frac{z+i}{z^2+1}$,
- ii. $\frac{1-\cos(z)}{z^2}$,
- iii. $\frac{1}{\cos(z)-1}$,
- iv. $e^{1/z}$

Show Solution 9.1 on P218

Exercise 9.2

Determine whether the domain $D(P)$ of the following formal bilateral series P is non-empty and, if so, find the radii of convergence. i. $\sum_{n=-\infty}^{\infty} \frac{z^n}{|n|!}$,

- ii. $\sum_{n=-\infty}^{\infty} z^n$,
- iii. $\sum_{n=-\infty}^{\infty} \frac{|n|^{|n|}}{|n|!} z^n$.

Show Solution 9.2 on P220

Exercise 9.3

Use the geometric series to find the Laurent expansion of

$$f(z) = \frac{1}{z(1-z)}$$

on the domains

- i. $0 < |z| < 1$,
- ii. $0 < |z - 1| < 1$,
- iii. $1 < |z| < +\infty$.

Show Solution 9.3 on P221

Exercise 9.4

Let $P = \sum_{n=-\infty}^{\infty} a_n T^n$, $Q = \sum_{n=-\infty}^{\infty} b_n T^n$ be formal bilateral series. Suppose that both domains $D(P, z_0)$, $D(Q, z_0)$ contain a non-empty annulus $A_{r,R}(z_0)$. Assume that for some radius s with $r < s < R$ we have

$$P(z) = Q(z), \quad \text{for all } |z - z_0| = s.$$

Prove the *identity theorem*, namely that $a_n = b_n$ for all $n \in \mathbb{Z}$.

Show Solution 9.4 on P222

Exercise 9.5

Let U be open and $z_0 \in U$. Let $f: U \rightarrow \mathbb{C}$ and $g: U \setminus \{z_0\} \rightarrow \mathbb{C}$ be holomorphic functions.

- i. If $f(z_0) \neq 0$ and g has a pole of order one at z_0 , prove that fg has a pole of order one at z_0 with

$$\operatorname{Res}_{z_0}(fg) = f(z_0) \operatorname{Res}_{z_0}(g).$$

- ii. If $f(z_0) = 0$ and $f'(z_0) \neq 0$, prove that $1/f(z)$ has a pole of order one at z_0 with $\operatorname{Res}_{z_0}(1/f) = 1/f'(z_0)$.

Show Solution 9.5 on P223

Chapter 10

Residue theorem

Definition 10.1. Let z_0 be an isolated singularity of a holomorphic function $f: U \rightarrow \mathbb{C}$. The **residue** $\text{Res}_{z_0}(f)$ is the residue of the Laurent series Equation 9.9. In other words,

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n \implies \text{Res}_{z_0}(f) = a_{-1}.$$

Proposition 10.1. Let z_0 be an isolated singularity of both the holomorphic functions $f, g: U \rightarrow \mathbb{C}$.

- a. $\text{Res}_{z_0}(f + g) = \text{Res}_{z_0}(f) + \text{Res}_{z_0}(g)$
- b. $\text{Res}_{z_0}(\lambda f) = \lambda \text{Res}_{z_0}(f)$ for all $\lambda \in \mathbb{C}$
- c. If z_0 is a pole of order m , then $\text{Res}_{z_0}(f) = \frac{1}{(m-1)!} \lim_{z \rightarrow z_0} h^{(m-1)}(z)$ for $h(z) = (z - z_0)^m f(z)$.

Proof.

a. and b. are left as an Exercise.

b. If $f = \sum_{n=-m}^{\infty} a_n(z - z_0)^n$, then $h(z) = \sum_{n=-m}^{\infty} a_n(z - z_0)^{n+m}$ is a power

series which we can differentiate termwise by Theorem 5.2 to get

$$h^{(m-1)}(z) = \sum_{n=-m}^{\infty} (n+m)(n+m-1)\cdots(n+2)a_n(z-z_0)^{n+1}.$$

The limit $z \rightarrow z_0$ of this power series is its constant term, $(m-1)!a_{-1}$.

□

Example 10.1.

$$\operatorname{Res}_0(z^n) = \begin{cases} 0 & \text{if } n \neq -1, \\ 1 & \text{if } n = -1. \end{cases}$$

Example 10.2.

The best way to compute a residue is often to work out the Laurent expansion using the calculus of formal power series from Chapter 5.

(a) Consider $f(z) = \frac{\sin(z^2)}{z^7}$. As $\sin(z) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} z^{2n+1}$, we have

$$f(z) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} z^{4n+2-7},$$

hence $\operatorname{Res}_0 f(z) = -1/6$ is the coefficient of the term with $n = 1$.

(b) Let $g(z) = \frac{\cos(z)-1}{(\exp(z)-1)^3}$. It suffices to work with finite portions of the involved power series. For this, introduce the notation $O(z^m)$ to mean 'terms involving z^n of order $m \leq n$ '. We will use this in an informal way, which can be made precise in algebra by saying that $O(z^n)$ is the ideal in the ring of formal power

series generated by z^n . Notice the properties

$$O(z^m)O(z^n) = O(z^{m+n}), \quad (10.1)$$

$$O(z^m)^n = O(z^{mn}), \quad (10.2)$$

$$O(z^m) + O(z^n) = O(z^{\min(m,n)}), \quad (10.3)$$

$$1/O(z^n) = O(z^{-n}). \quad (10.4)$$

Now compute

$$\begin{aligned} \exp(z) - 1 &= z + z^2/2 + O(z^3), \\ \Rightarrow (\exp(z) - 1)^3 &= \frac{z^6}{8} + \frac{3z^5}{4} + \frac{3z^4}{2} + z^3 + O(z^5) \\ &= z^3 + \frac{3z^4}{2} + O(z^5). \end{aligned}$$

To find the expansion of the inverse $1/(\exp(z)-1)^3$, we apply the proof strategy of Proposition 5.1(b) to $z^{-3}(\exp(z)-1)^3 = 1 + \frac{3z}{2} + O(z^2)$. In other words, we compare coefficients to solve

$$\left(1 + \frac{3z}{2} + \dots\right) (a_0 + a_1z + a_2z^2 + \dots) = 1$$

recursively for a_0, a_1, a_2 . This gives

$$1/(\exp(z)-1)^3 = z^{-3} \left(1 - \frac{3z}{2} + \frac{9z^2}{4} + O(z^3)\right) = z^{-3} - \frac{3z^{-2}}{2} + \frac{9z^{-1}}{4} + O(z^0).$$

Finally, multiply this by $\cos(z) - 1 = -\frac{z^2}{2} + \frac{z^4}{24} + O(z^5)$ to get

$$g(z) = -\frac{1}{2z} + \frac{3}{4} - \frac{13z}{12} - \frac{z^2}{16} + O(z^2)$$

from which we read off $\text{Res}_0 g(z) = -1/2$.

Definition 10.2. Let $\gamma: [a, b] \rightarrow \mathbb{C}$ be a closed piecewise C^1 curve and let $z_0 \in \mathbb{C} \setminus \gamma([a, b])$. The **winding number** of γ around z_0 is

$$W_{z_0}(\gamma) = \frac{1}{2\pi i} \int_{\gamma} \frac{dz}{z - z_0}.$$

Example 10.3.

Let $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$, $\gamma(t) = z_0 + re^{it}$, be the boundary curve of the disk $\overline{D}_r(z_0)$. Then Equation 6.6 implies $W_{z_0}(\gamma) = 1$. More generally, for every $w_0 \in D_r(z_0)$ the Cauchy integral formula Equation 8.1 gives

$$W_{w_0}(\gamma) = 1.$$

Pick a subdivision $a = t_0 < t_1 < \dots < t_n = b$ such that $\gamma|_{[t_{k-1}, t_k]}$ is continuously differentiable. Using branches of the logarithm, we can write $\gamma(t) = z_0 + r(t)e^{i\theta(t)}$ for $r(t) = |\gamma(t) - z_0|$ and a continuous function $\theta: [a, b] \rightarrow \mathbb{R}$ with $\theta|_{[t_{k-1}, t_k]}$ continuously differentiable. With this notation, the following holds.

Proposition 10.2.

- (a) We have $W_{z_0}(\gamma) = \frac{\theta(b) - \theta(a)}{2\pi}$, which is an integer.
- (b) For closed curves γ_0, γ_1 in $\mathbb{C} \setminus \{z_0\}$ we have

$$W_{z_0}(\gamma_0) = W_{z_0}(\gamma_1) \iff \gamma_0, \gamma_1 \text{ are homotopic in } \mathbb{C} \setminus \{z_0\}.$$

Proof.

(a) Over $[t_{k-1}, t_k]$ we have $\gamma'(t) = r'(t)e^{i\theta(t)} + r(t)\theta'(t)e^{i\theta(t)}$. Using

$$\frac{\gamma'(t)}{\gamma(t) - z_0} = \frac{r'(t)}{r(t)} + i\theta'(t) \quad (10.5)$$

we compute

$$2\pi i W_{z_0}(\gamma) = \sum_{k=1}^n \int_{t_{k-1}}^{t_k} \frac{\gamma'(t)}{\gamma(t) - z_0} dt$$

by Equation 10.5 we have

$$\sum_{k=1}^n \int_{t_{k-1}}^{t_k} \frac{\gamma'(t)}{\gamma(t) - z_0} dt = \sum_{k=1}^n \left(\int_{t_{k-1}}^{t_k} \frac{d}{dt} \log r(t) dt + i \int_{t_{k-1}}^{t_k} \theta'(t) dt \right)$$

using FTC, the right hand side becomes

$$\sum_{k=1}^n \left(\log r(t_k) - \log r(t_{k-1}) + i\theta(t_k) - i\theta(t_{k-1}) \right)$$

observing that these are telescoping sums, we conclude that

$$2\pi i W_{z_0}(\gamma) = i(\theta(t_n) - \theta(t_0)) = i(\theta(b) - \theta(a))$$

and the claimed formula follows. From $\gamma(a) = \gamma(b)$ we have $e^{i\theta(a)} = e^{i\theta(b)}$, so Proposition 1.5 implies $\theta(b) - \theta(a) = 2\pi k$ for some $k \in \mathbb{Z}$.

(b) By Theorem 7.2 applied to $U = \mathbb{C} \setminus \{z_0\}$ and $f(z) = 1/(z - z_0)$, the implication ' $\Leftarrow$ ' holds.

Conversely, assume $W_{z_0}(\gamma_0) = W_{z_0}(\gamma_1)$. Write $\gamma_k(t) = z_0 + r_k(t)e^{i\theta_k(t)}$ for $k = 0, 1$. As $\gamma_k(a) = \gamma_k(b)$, we have $r_k(a) = r_k(b)$ and $\theta_k(a) = \theta_k(b) - 2\pi n_k$ for some $n_k \in \mathbb{Z}$. By assumption, $\frac{\theta_0(b) - \theta_0(a)}{2\pi} = \frac{\theta_1(b) - \theta_1(a)}{2\pi}$, hence $n_0 = n_1$.

Define the homotopy by

$$\Gamma_s(t) = z_0 + [(1-s)r_0(t) + sr_1(t)] \exp(i(1-s)\theta_0(t) + is\theta_1(t)).$$

Clearly, $\Gamma_k = \gamma_k$ for $k = 0, 1$. The only thing to check is that Γ_s is a closed curve for each $s \in [0, 1]$, that is, to show $\Gamma_s(b) = \Gamma_s(a)$. As $r_k(a) = r_k(b)$, this is equivalent to

$$\exp(i(1-s)(\theta_1(b) - \theta_1(a)) + is(\theta_0(b) - \theta_0(a))) = 1.$$

This holds since $\theta_1(b) - \theta_1(a) = \theta_0(b) - \theta_0(a) = 2\pi n_0$ and

$$\exp(i(1-s)2\pi n_0 + is2\pi n_0) = \exp(2\pi i n_0) = 1.$$

□

Theorem 10.1 (Residue theorem). *Let $U \subset \mathbb{C}$ be an open set, $\gamma: [a, b] \rightarrow U$ a null-homotopic piecewise C^1 loop in U , and $z_1, \dots, z_n \in U \setminus \gamma([a, b])$ points not on the loop γ . If f is a holomorphic function on $U \setminus \{z_1, \dots, z_n\}$, then*

$$\frac{1}{2\pi i} \int_{\gamma} f(z) dz = \sum_{k=1}^n W_{z_k}(\gamma) \operatorname{Res}_{z_k}(f).$$

Proof.

For each $k = 1, \dots, n$ we find $r_k > 0$ and Laurent expansions

$$f(z) = \sum_{n=-\infty}^{\infty} a_n^{(k)} (z - z_k)^n \quad \text{for all } z \in D_{r_k}^{\times}(z_k).$$

Let $P_k(z) = \sum_{n=-\infty}^{-1} a_n^{(k)}(z - z_k)^n$ be the corresponding principal part. Hence $a_{-1}^{(k)} = \text{Res}_{z_k}(f)$. By Proposition 9.1, $P_k(z)$ is holomorphic on $D_{+\infty}^\times(z_0)$. Also, $f(z) - \sum_{k=1}^n P_k(z)$ defines a holomorphic function on U since its singularities at $\{z_1, \dots, z_n\}$ are removable. By Cauchy's Theorem 7.1,

$$\int_{\gamma} f(z) dz = \sum_{k=1}^b \int_{\gamma} P_k(z) dz.$$

By Proposition 10.2(b), the loop γ is homotopic in $\mathbb{C} \setminus \{z_k\}$ to $\partial D_{r_k}(z_k) * \dots * \partial D_{r_k}(z_k)$ (the same loop traversed $W_{z_k}(\gamma)$ -times). Combined with the fact that P_k is holomorphic on $\mathbb{C} \setminus \{z_k\}$, this implies by Theorem 7.2 that

$$\begin{aligned} \int_{\gamma} P_k(z) dz &= W_{z_k}(\gamma) \int_{\partial D_{r_k}(z_k)} P_k(z) dz \\ &= W_{z_k}(\gamma) \int_{\partial D_{r_k}(z_k)} \sum_{n=-\infty}^{-1} a_n^{(k)}(z - z_k)^n dz \end{aligned}$$

by uniform convergence,

$$W_{z_k}(\gamma) \int_{\partial D_{r_k}(z_k)} \sum_{n=-\infty}^{-1} a_n^{(k)}(z - z_k)^n dz = W_{z_k}(\gamma) \sum_{n=-\infty}^{-1} a_n^{(k)} \int_{\partial D_{r_k}(z_k)} (z - z_k)^n dz$$

using Equation 6.6 we have

$$W_{z_k}(\gamma) \sum_{n=-\infty}^{-1} a_n^{(k)} \int_{\partial D_{r_k}(z_k)} (z - z_k)^n dz = W_{z_k}(\gamma) a_{-1}^{(k)} 2\pi i$$

as required. □

10.1 Exercises

Exercise 10.1

Determine the winding numbers $W_{z_0}(\gamma)$ in the following cases.

- a. $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$, $\gamma(t) = e^{it}$, for $z_0 = 0, 2$.
- b. $\gamma: [2\pi, 10\pi] \rightarrow \mathbb{C}$, $\gamma(t) = \begin{cases} te^{it} & \text{if } t \in [2\pi, 6\pi], \\ 12\pi - t & \text{if } t \in [6\pi, 10\pi], \end{cases}$ for $z_0 = 0, 20$.
- c. $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$, $\gamma(t) = (2 + \cos(t))e^{2it}$, for $z_0 = 0, 4$.

Exercise 10.2

Use the residue theorem to compute

$$\int_{\partial D_r(0)} \frac{(z+2)^2}{z(z-1)^2} dz$$

for all $0 < r \neq 1$.

Hint: Consider the cases $r > 1$ and $r < 1$ separately.

Exercise 10.3

Apply the residue theorem to the contour $\partial D_1(0)$ and a suitable holomorphic function $f(z)$ to compute the integral


$$\int_0^{2\pi} \frac{dx}{2 + \cos(x)}.$$

Exercise 10.4

Compute the integral

$$\int_0^{+\infty} \frac{\cos(x)}{\sqrt{x}} dx$$

by considering $f(z) = \frac{e^{iz}}{\sqrt{z}}$ and the following contour $\gamma_{r,R}$ for $r \rightarrow 0$, $R \rightarrow +\infty$.




Hint: Recall that $\int_0^\infty e^{-x^2} dx = \sqrt{\pi}/2$.

Exercise 10.5

Compute the integral

$$\int_{-\infty}^{+\infty} \frac{dx}{1+x^6}$$

by applying the residue theorem to $f(z) = \frac{1}{1+z^6}$ and the following upper semi-circular contour γ_R for $R \rightarrow +\infty$.



Chapter 11


Applications to integrals

The residue theorem can be used to compute integrals of functions that do not have an obvious primitive. For this, we first need to find a suitable holomorphic function and contour to which we can apply the residue theorem, which is guesswork. The original integral is often a part of this contour and one shows that the rest of contour integral tends to zero.

We give three examples that illustrate the general technique.

Example 11.1.

Let $f(z) = \frac{1}{1+z^4}$, which is holomorphic except for poles of order one at all points ζ^{1+2k} , $k = 0, 1, 2, 3$, where $\zeta = e^{i\pi/4}$. For $R > 0$ introduce the semi-circular contour $\eta_R: [0, \pi] \rightarrow \mathbb{C}$, $\eta_R(t) = Re^{it}$ and let γ_R be the concatenation of η_R with the real interval $[-R, R]$, parametrized as in Example [6.1](#).



The residues are

$$\operatorname{Res}_{\zeta^{1+2k}}(f) = \frac{-\zeta^{1+2k}}{4}.$$

When $R > 1$, the only isolated singularities inside the contour are ζ, ζ^3 , hence $W_\zeta(\gamma_R) = W_{\zeta^3}(\gamma_R) = 1$ (more formally, this holds by Example 10.3 and Proposition 10.2) and $W_{\zeta^5}(\gamma_R) = W_{\zeta^7}(\gamma_R) = 0$ (more formally, by Cauchy's Theorem 7.1). By the residue Theorem 10.1,

$$\int_{\gamma_R} f(z)dz = 2\pi i[W_\zeta(\gamma_R) \operatorname{Res}_\zeta(f) + W_{\zeta^3}(\gamma_R) \operatorname{Res}_{\zeta^3}(f)] = \frac{\pi}{\sqrt{2}}$$

The contour integral decomposes as

$$\frac{\pi}{\sqrt{2}} = \int_{\gamma_R} f(z)dz = \int_{\eta_R} f(z)dz + \int_{[-R,R]} f(z)dz$$

and we have

$$\int_{-\infty}^{+\infty} f(z)dz = \lim_{R \rightarrow +\infty} \int_{[-R,R]} f(z)dz.$$

We claim that $\int_{\eta_R} f(z)dz \rightarrow 0$ as $R \rightarrow +\infty$. We have $L(\eta_R) = \pi R$ and

$$|f(z)| = \frac{1}{|z^4 + 1|} \leq \frac{1}{R^4 - 1}$$

where the inequality follows by Equation 1.12.

Hence Equation 6.7 implies $\left| \int_{\eta_R} f(z) dz \right| \leq \frac{\pi R}{R^4 - 1}$, which indeed tends to zero as $R \rightarrow +\infty$. Putting things together, we get

$$\int_{-\infty}^{+\infty} \frac{dx}{1+x^4} = \frac{\pi}{\sqrt{2}}.$$

More generally, we can use the approach above to **compute all integrals** $\int_{-\infty}^{+\infty} f(z) dz$ **of rational functions** $f(z) = \frac{p(z)}{q(z)}$ with $\deg(q) \geq \deg(p) + 2$ and $q(x) \neq 0$ for all $x \in \mathbb{R}$.

Integrals made only of trigonometric functions can often be evaluated using the residue theorem and the following method.

Example 11.2.

If $x \in \mathbb{R}$, then $\cos(x) = \frac{z+1/z}{2}$, $\sin(x) = \frac{z-1/z}{2i}$ for $z = e^{ix}$. Using this, we can change variables and use $dz = iz dx$ to rewrite

$$\int_0^{2\pi} \frac{\cos(x)^2}{\sin(x) + 2} dx = \int_{\partial D_1(0)} \frac{(1+z^2)^2}{2z^2(z^2 + 4iz - 1)} dz$$

as a contour integral, which is evaluated using the residue theorem. The singularities of $f(z) = \frac{(1+z^2)^2}{2z^2(z^2 + 4iz - 1)} = \frac{(1+z^2)^2}{2z^2(z-z_+)(z-z_-)}$ are at $z = 0$ (pole of order two) and $z_{\pm} = -i(2 \pm \sqrt{3})$ (poles of order one). Using Proposition 10.1(c), we find that the residues are

$$\begin{aligned} \text{Res}_0(f) &= \lim_{z \rightarrow 0} \frac{d}{dz} \frac{(1+z^2)^2}{2(z^2 + 4iz - 1)} = -2i, \\ \text{Res}_{z_-}(f) &= \lim_{z \rightarrow z_-} \frac{(1+z^2)^2}{2z^2(z-z_+)} = \frac{(1+z_-^2)^2}{2z_-^2(z_- - z_+)} = i\sqrt{3}. \end{aligned}$$

Now the residue theorem implies

$$\int_{\partial D_1(0)} \frac{(1+z^2)^2}{2z^2(z^2+4iz-1)} dz = 2\pi i(-2i+i\sqrt{3}) = 2\pi(\sqrt{3}-2).$$

This example can be generalized to compute integrals over $[0, 2\pi]$ of rational algebraic functions of $\sin(x)$ and $\cos(x)$.

The following integral was already computed in calculus using polar coordinates in the plane.

Example 11.3.

$$\int_{-\infty}^{+\infty} e^{-x^2} dx = \sqrt{\pi}$$

Let $a = (1+i)\sqrt{\frac{\pi}{2}}$. Then $a^2 = i\pi$. The function $f(z) = \frac{e^{-z^2}}{1+e^{-2az}}$ is holomorphic except for poles of order one at all points $a(k + \frac{1}{2})$, $k \in \mathbb{Z}$ (simple zeros of the denominator). Using the power series expansion of the exponential, we find

$$\begin{aligned} 1 + e^{-2az} &= 1 + e^{-2a(z-\frac{a}{2})}e^{a^2} = 1 - e^{-2a(z-\frac{a}{2})} \\ &= 2a\left(z - \frac{a}{2}\right) - \frac{1}{2!}4a^2\left(z - \frac{a}{2}\right)^2 + \dots \end{aligned}$$


and this gives the residue

$$\text{Res}_{a/2}(f) = \lim_{z \rightarrow a/2} \frac{e^{-z^2}(z - \frac{a}{2})}{1 + e^{-2az}} = \frac{e^{-a^2/4}}{2a} = \frac{-i}{2\sqrt{\pi}}.$$

From $a^2 = i\pi$ it is not hard to check

$$f(z) - f(z+a) = e^{-z^2}.$$

Hence the integral over $z \in [-\infty, +\infty]$ can be computed using the two horizontal parts of the following contour γ_R as $R \rightarrow +\infty$.



The only singularity inside the contour is $a/2$, so the residue theorem gives

$$\int_{\gamma_R} f(z)dz = 2\pi i \frac{-i}{2\sqrt{\pi}} = \sqrt{\pi}.$$

The contour integral over γ_R splits into four parts. We estimate the vertical parts $\eta_{\pm R}$ of the integral as follows. Let $z = x + iy$ with $x = \pm R$ and $y \in [0, \sqrt{\pi/2}]$. Then

$$|f(z)| \leq \frac{e^{-\Re(z^2)}}{1 - e^{-\Re(2az)}} = \frac{e^{y^2 - x^2}}{1 - e^{\sqrt{2\pi}(y-x)}} \leq \frac{e^{\pi/2 - x^2}}{1 - e^{-\sqrt{2\pi}x}}$$

tends to zero as $|x| \rightarrow +\infty$, uniformly in $y \in [0, \sqrt{\pi/2}]$. Hence Equation 6.7 implies

$$\int_{\eta_{\pm R}} f(z)dz \xrightarrow{R \rightarrow \infty} 0$$

and we conclude

$$\begin{aligned} \sqrt{\pi} &= \int_{\gamma_R} f(z)dz \\ &= \int_{-R}^{+R} f(z)dz + \int_R^{-R} f(z+a)dz + \int_{\eta_R} f(z)dz + \int_{\eta_{-R}} f(z)dz \\ &\xrightarrow{R \rightarrow +\infty} \int_{-\infty}^{+\infty} f(z)dz - \int_{-\infty}^{+\infty} f(z+a)dz = \int_{-\infty}^{+\infty} e^{-x^2} dx. \end{aligned}$$

Appendix A

All Solutions

A.1 Chapter 1 solutions

Solution 1.1

Verify

$$z = x + iy$$

and

$$i^2 = -1$$

straight from the definition Equation 1.2.

Let $z = (x, y) \in \mathbb{C}$. Identifying x with $(x, 0)$ and y with $(y, 0)$. We calculate

$$iy = (0, 1) \cdot (y, 0) = (0 - 0, 0 + y) = (0, y).$$

Consequently,

$$x + iy = (x, 0) + (0, y) = (x, y) = z.$$

To see that $i^2 = -1$, we calculate

$$(0, 1)(0, 1) = (0 - 1, 0 + 0) = (-1, 0).$$

This yields the desired conclusion under the identification $-1 = (-1, 0)$.

[Return to Exercise 1.1 on P22](#)

Solution 1.2

How many real solutions x does $x^2 + 1 = 0$ have? Show that the polynomial equation $z^2 + 1 = 0$ has exactly two solutions $z \in \mathbb{C}$.

This equation has no real solutions. To find all solutions in $\mathbb{C}$, we first rearrange to get $z^2 = -1$. This means that z must be one of the square roots of -1 . There are at least two ways to find all possible values of z .

The first approach uses Proposition 1.7.

Expressing -1 in polar form, we have $-1 = e^{i\pi}$.

By Proposition 1.7 the square roots of -1 , are

$$e^{\frac{i(\pi + 2k\pi)}{2}}$$

for $k = 0, 1$. This gives exactly two solutions:

$$i = e^{\frac{i(\pi)}{2}}, \text{ and } -i = e^{\frac{i(\pi + 2\pi)}{2}}.$$

The other approach is as follows. Write $z = x + iy$. Then,

$$z^2 = (x + iy)(x + iy) = (x^2 - y^2) + i(2xy) = -1.$$

This gives us two equations

$$x^2 - y^2 = -1 \tag{A.1}$$

$$2xy = 0. \tag{A.2}$$

The second equation implies one of x or y is 0.

If x is 0, then $-y^2 = -1$ and so $y = \pm 1$.

If y is 0, then $x^2 = -1$ and has no solutions (since x is a real number).

We conclude that $x = 0$ and $y = \pm 1$, which gives exactly two values for z , i and $-i$.

[Return to Exercise 1.2 on P22](#)

Exercise 1.3


Give examples of complex numbers $z, w \neq 0$ such that $z^2 + w^2 = 0$.

Take $z = 1, w = i$.

[Return to Exercise 1.3 on P22](#)

Solution 1.4

Sketch the position of the complex numbers i , $1 + i$, $\frac{3+2i}{4}$ in the plane.



[Return to Exercise 1.4 on P23](#)

Solution 1.5

Express the following complex numbers z in the form $x + iy$ with $x, y \in \mathbb{R}$.

$$(1 + i)^{20}, (5 + 3i)(1 + 2i), (1 - i)(2 + 3i), (1 - i)i(1 + i), \frac{2 + i}{1 - i}$$

$$(2i)^{10} = -1024, \quad -1 + 13i, \quad 5 + i, \quad 2i, \quad \frac{1 + 3i}{2}$$

[Return to Exercise 1.5 on P23](#)

Solution 1.6

Express the following complex numbers z in the form $x + iy$ with $x, y \in \mathbb{R}$.

$$1/i, \quad \frac{1}{1+i}, \quad \frac{3+i}{3-i}$$

$$-i, \quad \frac{1-i}{2}, \quad \frac{8+6i}{10}$$

[Return to Exercise 1.6 on P23](#)

Solution 1.7

Find the modulus and the conjugate of the following complex numbers.

$$2 + i, \quad i, \quad 5 - 3i, \quad \frac{1+i}{2+i}$$

$$|2 + i| = \sqrt{5}, \quad \overline{2 + i} = 2 - i \quad (\text{A.3})$$

$$|5 - 3i| = \sqrt{34}, \quad \overline{5 - 3i} = 5 + 3i \quad (\text{A.4})$$


$$\left| \frac{1+i}{2+i} \right| = \sqrt{\frac{2}{5}} = \frac{\sqrt{10}}{5}, \quad \frac{\overline{1+i}}{\overline{2+i}} = \frac{1-i}{2-i} \quad (\text{A.5})$$

[Return to Exercise 1.7 on P23](#)


Solution 1.8

Describe the sets $A = \{z \in \mathbb{C} \mid \Im(z) > 0\}$, $B = \{z \in \mathbb{C} \mid \Re(z) \leq 1\}$, $C = \{z \in \mathbb{C} \mid \Re((1+i)z) = 0\}$, **and** $A \cap B$ **geometrically.**


- The set A is the 'upper half-plane'




- The set B is the “left half-plane” shifted to the right



- The set C is the straight line $x = y$ since $\Re((1 + i)z) = x - y$.



- The intersection of A and B is pictured below




[Return to Exercise 1.8 on P23](#)

Solution 1.9

Describe the set $D = \{z \in \mathbb{C} \mid z \cdot \bar{z} = 1\}$ **geometrically.**

Hint: Write $z = re^{i\theta}$ in polar form.

Note that $z \cdot \bar{z} = |z|^2$. Hence points in D satisfy the equation $|z|^2 = 1$. This is the circle of radius one centred at the origin.



[Return to Exercise 1.9 on P24](#)


Solution 1.10

Draw all nine sets described by the following conditions on the complex number z .


$ z = 1,$	$ z < 1,$	$1 < z < 2,$
$ 1 + z > 1,$	$ 2 - z < 2,$	$3 < z + i < 4,$
$ z - 1 < z + 1 ,$	$ z = z + 1 ,$	$ z - 1 = z + i .$

- $|z| = 1$ is the unit circle.


- $|z| < 1$ is the open disk of radius 1 with center the origin.
- $1 < |z| < 2$ is the open annulus centered at the origin with radii 1 and 2.




- The complement of the set defined by $|1+z| > 1$ is the set of all $|1+z| \leq 1$, the closed disk of radius 1 centered at -1 . Hence $\{z \in \mathbb{C} \mid |1+z| > 1\}$ is the complement of this closed disk.




- $|2 - z| < 2$ is the open disk of radius 2 centered at 2.



- $3 < |z + i| < 4$ is an annulus centered at $-i$ with radii 3 and 4.
- $|z - 1| < |z + 1|$ is the set of points that are closer to 1 than to -1 . This is the 'right half-plane' $\{x + iy \mid x > 0\}$. In more detail, write $z = x + iy$. Then the inequality is equivalent to $(x - 1)^2 + y^2 < (x + 1)^2 + y^2$, so to $-2x < 2x$. This just means $x > 0$.



- $|z| = |z + 1|$ is the set of points equidistant from 0 and -1 . Write $z = x + iy$. Then the condition is equivalent to $x^2 + y^2 = (x + 1)^2 + 2(x + 1)y + y^2$, so equivalent to $2x + 1 + y(2x + 1) = 0$. This implies $x = -1/2$ and therefore $\{z \in \mathbb{C} \mid |z| = |z + 1|\}$ is the straight line $\{x + iy \mid x = -1/2\}$.



- $|z - 1| = |z + i|$ is the set of points equidistant from 1 and $-i$. This is again a straight line, namely $\{x - ix \mid x \in \mathbb{R}\}$. \end{itemize}

[Return to Exercise 1.10 on P24](#)

Solution 1.11

Let $S = \{x + iy \in \mathbb{C} \mid 0 \leq x, y \leq 1\}$. **Draw** S **and the sets**

$$A = \{2z \mid z \in S\},$$


$$B = \{\bar{z} \mid z \in S\},$$

$$C = \{-z \mid z \in S\},$$

$$D = \{z^2 \mid z \in S\}.$$

A is the square with corners $\{0, 2, 2 + 2i, 2i\}$. B is the square with corners $\{0, 1, 1 - i, -i\}$. C is the square with corners $\{0, -i, -1 - i, -1\}$.

D is much harder to work out. It is shown in the picture. The edge joining 0 to 1 remains as it was. The edge joining 1 to $1 + i$ can be parametrised as $\{1 + it \mid 0 \leq t \leq 1\}$ and, on squaring, becomes the curve parametrised as $\{(1 - t^2, 2t) \mid 0 \leq t \leq 1\}$ and this is the right-hand curved edge. In the same way the line joining $1 + i$ to i becomes the left-hand curved edge. Finally, the line joining i to 0 becomes the line joining -1 to 0.



[Return to Exercise 1.11 on P24](#)

Solution 1.12

Let $D = \{z \in \mathbb{C} \mid |z| < 1\}$ be the unit disk. Draw the sets

$$A = \{2z \mid z \in D\}, \quad B = \{z^2 \mid z \in D\}, \quad C = \{|z| \mid z \in D\}.$$

- Since $|2z| = 2|z| < 2 \iff |z| < 1$ the set A is the open disk of radius 2 at the origin.
- As $|z^2| < |z|^2 < 1$ for $z \in D$ we have $B \subset D$. Conversely, if $w \in D$, we know there exists a square root z with $z^2 = w$. As $|z| = |z^2|^{1/2} < |w|^{1/2} < 1$ it follows that $w \in B$. Hence $B = D$.
- C is the real interval $\{x + i0 \mid 0 \leq x < 1\} = [0, 1)$.

[Return to Exercise 1.12 on P25](#)

Solution 1.13

Show that $i = e^{i\pi/2}$ and $-1 = e^{i\pi}$.

We have

$$e^{i\pi/2} = \cos(\pi/2) + i \sin(\pi/2) = 0 + i = i$$

and

$$e^{i\pi} = \cos(\pi) + i \sin(\pi) = -1 + 0 = -1.$$

[Return to Exercise 1.13 on P25](#)

Solution 1.14

Express the following complex numbers z in the form $x + iy$ with $x, y \in \mathbb{R}$.

$$e^{i\pi/4}, \quad e^{i\pi}, \quad e^{i\frac{2\pi}{3}}$$

$$\frac{\sqrt{2}}{2}, -1, \frac{-1 + i\sqrt{3}}{2}$$

[Return to Exercise 1.14 on P25](#)

Solution 1.15

Write each of the following complex numbers in polar form $re^{i\theta}$ with $r > 0$ and $-\pi < \theta \leq \pi$.

$$i, -1, -i, 1+i, 1-i, i-1, \frac{1}{2} + i\frac{\sqrt{3}}{2}$$

Draw each of these numbers in the complex plane.

$$e^{i\pi/2}, e^{i\pi}, e^{-i\pi/2}, \sqrt{2}e^{i\pi/4}, \sqrt{2}e^{-i\pi/4}, \sqrt{2}e^{3i\pi/4}, e^{i\pi/3}.$$

Be careful to take the angle in the correct range $-\pi < \theta \leq \pi$.

Now we express $(e^{i\pi/2})^{2020}$ as

$$(e^{i\pi})^{1010} = (-1)^{1010} = 1.$$

We conclude that $i^{2021} = i$.

For the second expression, we write

$$(i + 1)^{20} = \left(\sqrt{2}e^{i\pi/4}\right)^{20} = 2^{10}e^{i5\pi} = -1024.$$

[Return to Exercise 1.16 on P25](#)

Solution 1.17

Solve the equation $(1 - i)^n - 2075 = 2021$ **and find** $n \in \mathbb{N}$.

Writing $1 - i = \sqrt{2}e^{-i\pi/4}$, the equation is equivalent to $2^{n/2}e^{-in\pi/4} = 4096$.

Taking the modulus gives $2^{n/2} = 4096$ so necessarily $n = 24$ and indeed we have $(1 - i)^{24} = 4096$.

[Return to Exercise 1.17 on P26](#)

Solution 1.18

Prove that for $z \in \mathbb{R}$ **we have**

$$\cos(z) = \frac{e^{iz} + e^{-iz}}{2}, \quad \sin(z) = \frac{e^{iz} - e^{-iz}}{2i}.$$

Use these equations to extend the definition of the functions $\cos(z), \sin(z)$ **to complex arguments** $z \in \mathbb{C}$.

Find $z \in \mathbb{C}$ **with** $\sin(z) = 2$.

Hint: Put $w = e^{iz}$ and reduce to a quadratic equation.

Expanding we have

$$\frac{e^{iz} + e^{-iz}}{2} = \frac{\cos z + i \sin z + \cos z - i \sin z}{2} = \frac{2 \cos z}{2} = \cos z \quad (\text{A.6})$$

$$\frac{e^{iz} - e^{-iz}}{2i} = \frac{\cos z + i \sin z - \cos z + i \sin z}{2i} = \frac{2i \sin z}{2i} = \sin z. \quad (\text{A.7})$$

We solve

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i} = 2$$

for $z \in \mathbb{C}$. Rearranging, we have

$$e^{iz} - e^{-iz} = 4i.$$

Multiplying through by e^{iz} gives,

$$e^{2iz} - 4ie^{iz} - 1 = 0$$

. Setting $t = e^{iz}$ gives the quadratic equation:

$$t^2 - 4it - 1 = 0$$

. Solving, we get

$$t = e^{i(x+iy)} = i(2 \pm \sqrt{3}).$$

Expanding out the term $e^{i(x+iy)}$, gives

$$e^{-y}e^{ix} = i(2 \pm \sqrt{3}).$$

We conclude that $x = \pi/2 + 2k\pi$ for $k \in \mathbb{Z}$ (since $e^{ix} = i$) and $y = -\ln(2 \pm \sqrt{3})$ (since $e^{-y} = 2 \pm \sqrt{3}$).

Therefore $z = (\pi/2 + 2k\pi) - \ln(2 \pm \sqrt{3})$ for $k \in \mathbb{Z}$.

[Return to Exercise 1.18 on P26](#)

Solution 1.19

Prove the following statements.

- a. $\overline{zw} = \overline{z} \cdot \overline{w}$ for all $z, w \in \mathbb{C}$
- b. $\overline{z_1 z_2 \cdots z_n} = \overline{z_1} \overline{z_2} \cdots \overline{z_n}$ for all $z_1, z_2, \dots, z_n \in \mathbb{C}$ (use induction)
- c. $\overline{(z^n)} = (\overline{z})^n$ for all $z \in \mathbb{C}$
- d. Let $p(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0$ be a polynomial with real coefficients $a_0, \dots, a_n \in \mathbb{R}$. Prove that $\overline{p(z)} = p(\overline{z})$. Deduce that all roots of $p(z)$ occur in complex conjugate pairs.

a. Let $z = x + iy$ and $w = u + iv$. Then

$$\begin{aligned} zw &= (xu - yv) + i(xv + yu) \implies \overline{zw} = (xu - yv) - i(xv + yu) \\ \overline{z} \cdot \overline{w} &= (x - iy)(u - iv) = xu - yv + (-ivx - iyu). \end{aligned}$$

b. The base case $n = 1$ is trivial. For the inductive step, we have

$$\begin{aligned} \overline{z_1 \cdots z_n z_{n+1}} &= \overline{z_1 \cdots z_n} \cdot \overline{z_{n+1}} && \text{by (a)} \\ &= (\overline{z_1} \cdots \overline{z_n}) \overline{z_{n+1}} && \text{by induction.} \end{aligned}$$

c. follows from (b) by setting $z_1 = \dots = z_n = z$.

d. We have

$$\begin{aligned}
 \overline{p(z)} &= \overline{a_n z^n + \dots + a_1 z + a_0} \\
 &= \overline{a_n z^n} + \dots + \overline{a_1 z} + \overline{a_0} && \text{since } \overline{\cdot} \text{ is additive} \\
 &= \overline{a_n} \overline{z^n} + \dots + \overline{a_1} \cdot \overline{z} + \overline{a_0} && \text{by a.} \\
 &= a_n \overline{z}^n + \dots + a_1 \overline{z} + a_0 && \text{by (c) and since } a_k \in \mathbb{R} \\
 &= p(\overline{z}).
 \end{aligned}$$

Finally, suppose $p(\zeta) = 0$. Then $p(\overline{\zeta}) = \overline{p(\zeta)}$.

[Return to Exercise 1.19 on P26](#)

Solution 1.20

Show that $|z| = |-z|$ **and** $|\overline{z}| = |z|$. **Prove also that** $|\lambda z| = \lambda|z|$ **for all** $\lambda \geq 0$.

Write $z = x + iy$, then for $\lambda \geq 0$,

$$\begin{aligned}
 |z| &= \sqrt{x^2 + y^2} = \sqrt{(-x)^2 + (-y)^2} = |-z| \\
 \lambda|z| &= \lambda\sqrt{x^2 + y^2} = \sqrt{\lambda^2(x^2 + y^2)} = \sqrt{(\lambda x)^2 + (\lambda y)^2} = |\lambda z|
 \end{aligned}$$

[Return to Exercise 1.20 on P27](#)

Solution 1.21

Prove that $\Re(z) = \frac{1}{2}(z + \overline{z})$, $\Im(z) = \frac{1}{2i}(z - \overline{z})$.

Let $z = x + iy$. Then

$$z + \bar{z} = x + iy + x - iy = 2x = 2\Re(z)$$

and

$$z - \bar{z} = x + iy - (x - iy) = 2iy = 2i\Im(z).$$

[Return to Exercise 1.21 on P27](#)

Solution 1.22

Prove that $\overline{e^z} = e^{\bar{z}}$. **Deduce that** $|e^z| = e^{\Re(z)}$.

Let $z = x + iy$, then

$$\begin{aligned}\overline{e^z} &= \overline{e^{x+iy}} = \overline{e^x(\cos(y) + i\sin(y))} \\ &= e^x(\cos(y) - i\sin(y)) = e^x(\cos(-y) + i\sin(-y)) = e^{\bar{z}}.\end{aligned}$$

For the second part,

$$|e^z| = \sqrt{e^z \cdot \overline{e^z}} = \sqrt{e^{z+\bar{z}}} = \sqrt{e^{2\Re(z)}} = e^{\Re(z)}.$$

[Return to Exercise 1.22 on P27](#)

Solution 1.23

Prove that $|z + w|^2 + |z - w|^2 = 2|z|^2 + 2|w|^2$.

Write $z = x + iy$ and $w = u + iv$, then

$$\begin{aligned}
|z + w|^2 + |z - w|^2 &= (x + u)^2 + (y + v)^2 + (x - u)^2 + (y - v)^2 \\
&= 2(x^2 + u^2) + 2(y^2 + v^2) = 2|z|^2 + 2|w|^2.
\end{aligned}$$

Return to Exercise 1.23 on P27

Solution 1.24

Show that $|z + w|^2 = |z|^2 + 2\Re(z\bar{w}) + |w|^2$. **Use this to determine the conditions on z, w for $|z + w| = |z| + |w|$ to hold.**

Write $z = x + iy$ and $w = u + iv$, then

$$|z + w|^2 = (x + u)^2 + (y + v)^2 = (x^2 + y^2) + 2xu + 2yv + (y^2 + v^2).$$

Noting that $\Re(z\bar{w}) = xu + yv$, we have

$$|z + w|^2 = |z|^2 + 2\Re(z\bar{w}) + |w|^2$$

as required.

For the second part write $w = \lambda z$ for some $\lambda \in \mathbb{C}$. We show that λ is a positive real number. Suppose $|z + w| = |z| + |w|$ then

$$|z + \lambda z| = |z| + |\lambda z|.$$

It follows, assuming $z \neq 0$ that

$$|1 + \lambda| = |1| + |\lambda|.$$

Using the preceding part $2\Re(\lambda) = 2|\lambda|$ and we conclude that $|\lambda| = \Re(\lambda)$ and so λ is a positive real number.

If $z = 0$, then we may take $\lambda = 0$.

[Return to Exercise 1.24 on P28](#)

Solution 1.25

Assuming we know the triangle inequality $|z + w| \leq |z| + |w|$ **for all** $z, w \in \mathbb{C}$, **prove the reverse triangle inequality**

$$|z - w| \geq ||z| - |w||.$$

Applying the triangle inequality to z and $w - z$ gives

$$|w| = |z + (w - z)| \leq |z| + |w - z|,$$

which we rearrange to

$$|w| - |z| \leq |w - z|. \quad (\text{A})$$

By exchanging the roles of z and w we get

$$|z| - |w| \leq |z - w|. \quad (\text{B})$$

The two inequalities (A) and (B) are equivalent to the single inequality $||z| - |w|| \leq |z - w|$.

[Return to Exercise 1.25 on P28](#)

Solution 1.26

Let K be a field with $\mathbb{R} \subset K \subset \mathbb{C}$. Prove that $K = \mathbb{R}$ or $K = \mathbb{C}$.

Suppose that $K \neq \mathbb{R}$. Then there exists $z = x + iy \in K$ with $y \neq 0$. We know that $x \in K$ since x is real. So, by the basic properties of a field, $iy = z - x \in K$. Now y is real (so in K) and non-zero. So $i = (iy)/y \in K$. That's all we need, since we can now deduce that $x + iy \in K$ for all $x, y \in \mathbb{R}$.

[Return to Exercise 1.26 on P28](#)

A.2 Chapter 2 solutions

Solution 2.1

Describe the image set of the complex function $f(z) = \frac{1+z}{1-z}$ with domain $D = \mathbb{C} \setminus \{1\}$. In other words, determine the set of all $w \in \mathbb{C}$ for which $w = \frac{1+z}{1-z}$ has a solution $z \in D$.

If $w \neq -1$, the equation $w = \frac{1+z}{1-z}$ can be rearranged as

$$z = \frac{w-1}{w+1}.$$

Hence $\mathbb{C} \setminus \{-1\}$ is contained in the image of f . Also, -1 is not in the image of f since $-1 = \frac{1+z}{1-z}$ would imply $z-1 = 1+z$, so $-1 = 1$, which is impossible.


[Return to Exercise 2.1 on P39](#)

Solution 2.2


Sketch the following curves $z(t)$ in the complex plane, where t is a real parameter.

- a. $z(t) = t(1 + i)$ for $0 \leq t \leq 1$
- b. $z(t) = \cos(t) + i \sin(t)$ for $0 \leq t \leq \pi$
- c. $z(t) = \cos(t) - i \sin(t)$ for $0 \leq t \leq \pi/2$
- d. $z(t) = \frac{1}{1+it}$ for $t \in \mathbb{R}$


a. is a straight line from the origin to $1 + i$.



b. Thinking in polar coordinates, this the upper semicircle from $+1$ to -1 .



- c. We have $z(t) = \cos(-t) + i \sin(-t) = \cos(s) + i \sin(s)$ where $s = -t \in [-\pi/2, 0]$. This is the quarter circle from $+1$ to $-i$.




- d. By plotting the points for $t = 0, \pm 1$ and $t \rightarrow \pm\infty$, we suspect that $z(t)$ lies entirely on the circle centered at $1/2$. We verify this as follows.

$$\begin{aligned} \left| z(t) - \frac{1}{2} \right| &= \left| \frac{1-it}{t^2+1} - \frac{1+t^2}{2+2t^2} \right| = \left| \frac{1+(it)^2-2it}{2+2t^2} \right| \\ &= \left| \frac{(1+it)^2}{2+2t^2} \right| = \frac{|1+it|^2}{|2+2t^2|} = \frac{1+t^2}{2+2t^2} = \frac{1}{2}. \end{aligned}$$

We know that $z(0) = 1$, $\lim_{t \rightarrow \pm\infty} z(t) = 0$, $z(\pm 1) = \frac{1 \mp i}{2}$ and that the image of z is a connected subset. Moreover, $z(t) \neq 0$ for all t . It follows that the image of $z(t)$ is a circle of radius $1/2$ centered at $1/2$ with the origin removed. Note, however, that this parameterization of the circle

has an unusual speed.



[Return to Exercise 2.2 on P39](#)

Solution 2.3

Let $\log(z)$ be the principal branch of the logarithm. Compute

$$\log(2i), \log(1+i), \log(-3i), \log(5).$$

Recall that the principal branch is defined by $\log(z) = \log(r) + i\theta$ if $z = re^{i\theta}$,

where $r > 0$, $\theta \in (-\pi, \pi)$. We obtain the values

$$\log(2) + \frac{i\pi}{2}, \quad \log(\sqrt{2}) + \frac{i\pi}{4}, \quad \log(3) - \frac{i\pi}{2}, \log(5).$$

[Return to Exercise 2.3 on P39](#)

Solution 2.4

Describe the following complex functions geometrically.

$$f(z) = 3z, f(z) = iz, f(z) = \frac{(1+i)}{\sqrt{2}}z$$

$f(z) = 3z$ scales every complex number (thought of as a vector in $\mathbb{R}^2$) by the factor 3.

$f(z) = iz$ rotates counterclockwise by 90 degrees.

As $\frac{1+i}{\sqrt{2}} = e^{i\pi/4}$, the map $f(z) = \frac{(1+i)}{\sqrt{2}}z$ is a counterclockwise rotation by 45 degrees.

[Return to Exercise 2.4 on P39](#)

Solution 2.5

Determine a domain of definition for the following complex functions.

$$f(z) = \frac{1}{z}, f(z) = \frac{1+z}{z-1}, f(z) = \frac{z^2-4}{z^2+2z}, f(z) = \frac{1}{\exp(z)}.$$

For $f(z) = 1/z$, $D = \mathbb{C} \setminus \{0\}$.

For $f(z) = \frac{1+z}{z-1}$, $D = \mathbb{C} \setminus \{1\}$.

For $f(z) = \frac{z^2-4}{z^2+2z}$, it seems at first that both $z = 0, -2$ are problematic, but we can rewrite $\frac{z^2-4}{z^2+2z} = \frac{z-2}{z}$ since $z^2 - 4 = (z-2)(z+2)$. We can therefore extend the definition of f to the domain $D = \mathbb{C} \setminus \{0\}$.

For $f(z) = \frac{1}{\exp(z)}$, $D = \mathbb{C}$ as $\exp(z) \neq 0$ for all z . This last fact holds since $|\exp(z)| = e^x \neq 0$ for all $z = x + iy$.

[Return to Exercise 2.5 on P40](#)

Solution 2.6

Determine the domain of definition for $f(z) = \frac{1}{\sin(z)}$.

We need to find the points at which $\sin(z) = 0$. Using the equality

$$\sin(z) = \frac{e^{iz} - e^{-iz}}{2i},$$

we need to find the points $z \in \mathbb{C}$ where $e^{iz} = e^{-iz}$, equivalently, where $e^{2iz} = 1$.

Writing $z = x + iy$, we find that $y = 0$, since

$$|e^{2iz}| = e^{-2y} = 1$$

and $x = k\pi$ since $2x = 0 + 2k\pi$, $k \in \mathbb{Z}$.

The domain of $1/\sin(z)$ is therefore $D = \mathbb{C} \setminus \{k\pi : k \in \mathbb{Z}\}$.

[Return to Exercise 2.6 on P40](#)

Solution 2.7

Find all solutions to the following equations:

- a. $e^z = -1$,
- b. $\sin(z) = -i$

a. As $-1 = e^{\pi i}$, the equation $e^z = -1$ implies $e^{z-\pi i} = 1$, hence the set of solutions is $\pi i + 2\pi i\mathbb{Z}$.

b. Recall that $\sin(z) = \frac{1}{2i}(e^{iz} - e^{-iz})$. Writing $q = e^{iz}$, the equation $\sin(z) = -i$ becomes $q - q^{-1} = 2$. This leads to the quadratic equation $q^2 - 2q - 1$ which has the roots $q_{\pm} = 1 \pm \sqrt{2}$. Hence $\sin(z) = -i \iff e^{iz} = q_+$ or $e^{iz} = q_-$. The first equation is equivalent to $iz \in \log(1 + \sqrt{2}) + 2\pi i\mathbb{Z}$ and by writing $q_- = (\sqrt{2} - 1)e^{\pi i}$ we find that the second equation is equivalent to $iz \in \log(\sqrt{2} - 1) + \pi i + 2\pi i\mathbb{Z}$. Summing up, the set of solutions is

$$\{-i \log(1 + \sqrt{2}) + 2\pi k \mid k \in \mathbb{Z}\} \cup \{-i \log(\sqrt{2} - 1) + \pi + 2\pi k \mid k \in \mathbb{Z}\}.$$

[Return to Exercise 2.7 on P40](#)

Solution 2.8

Prove that the composition $f \circ f'$ of two Möbius transformations is again a Möbius transformation. If we associate to f, f' the matrices

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, A' = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathbf{M}_{2 \times 2}(\mathbb{C}),$$

show that $f \circ f'$ is associated to the matrix product AA' .

Let $f(z) = \frac{az+b}{cz+d}$ and $g(z) = \frac{\alpha z+\beta}{\gamma z+\delta}$ be Möbius transformations. Then,

$$(f \circ g)(z) = \frac{ag(z) + b}{cg(z) + d}$$

. Using the definition of $g(z)$, this expands to

$$\frac{a\alpha z + a\beta + b\gamma z + b\delta}{\gamma z + \delta} \cdot \frac{\gamma z + \delta}{c\alpha z + c\beta + d\gamma z + d\delta}.$$

We consequently have,

$$f(g(z)) = \frac{(a\alpha + b\gamma)z + (a\beta + b\delta)}{(c\alpha + d\gamma)z + (c\beta + d\delta)}.$$

One can verify explicitly that $((a\alpha + b\gamma)(c\beta + d\delta)) - ((a\beta + b\delta)(c\alpha + d\gamma)) \neq 0$, however, this is neatly tackled by the second part of the question, since the product of two invertible matrices is again an invertible matrix.

Notice that,


$$AA' = \begin{pmatrix} a\alpha + b\gamma & a\beta + b\delta \\ c\alpha + d\gamma & c\beta + d\delta \end{pmatrix}$$

is precisely the matrix associated to $f \circ g$.

[Return to Exercise 2.8 on P40](#)

Solution 2.9

Draw the image grid for $\exp: \mathbb{C} \rightarrow \mathbb{C}^\times$.



[Return to Exercise 2.9 on P41](#)

A.3 Chapter 3 solutions

Solution 3.1

Recall the ratio and root test for series of real numbers.

Let $\sum_{k=0}^{\infty} a_n$ be a real series.

Ratio test Let $L = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$

- If $L > 1$, the the series is divergent;
- If $L < 1$ then the series is (absolutely) convergent;
- If $L = 1$ or does not exist, then the test is inconclusive.

Root test Let $C = \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}$

- if $C > 1$, then the series is divergent;
- if $C < 1$, then the series is (absolutely) convergent;
- if $C = 1$ and the limit approaches strictly from above then the series diverges;

- otherwise the test is inconclusive.

Return to Exercise 3.1 on P47

Solution 3.2

For which $z \in \mathbb{C}$ do the following limits exist?

$$\lim_{n \rightarrow \infty} n^{1/n} z, \lim_{n \rightarrow \infty} z^n, \lim_{n \rightarrow \infty} \frac{z^n}{n}, \lim_{n \rightarrow \infty} \frac{z^n}{n!}, \lim_{n \rightarrow \infty} \frac{z^n}{n^n}, \lim_{n \rightarrow \infty} n! z.$$

The sequence $n^{1/n}$ is convergent with limit equal to 1 and so the sequence $n^{1/n}|z|$ is convergent for all complex numbers z with limit equal to z .

If $|z| > 1$, then z^n is unbounded, hence does not converge. If $|z| < 1$, then $\lim_{n \rightarrow \infty} z^n = 0$.

Suppose $|z| = 1$ and that $\lim_{n \rightarrow \infty} z^n$ exists. As z^n is a Cauchy sequence, we have in particular that $|z - 1| = |z|^n |z - 1| = |z^{n+1} - z^n| \rightarrow 0$ for $n \rightarrow \infty$. Hence we must have $z = 1$ and in this case the sequence indeed converges.

As $\frac{z^{n+1}/(n+1)}{z^n/n} = \frac{n}{n+1} z \rightarrow z$ for $n \rightarrow \infty$, the series $\sum_{n=0}^{\infty} \frac{z^n}{n}$ converges for all $|z| < 1$ by the ratio test. Hence $\lim_{n \rightarrow \infty} \frac{z^n}{n} = 0$ for all $|z| < 1$. Alternatively, apply by l'Hôpital's rule to the real limit $\frac{x^n}{n}$, where $x = |z|$. If $|z| = 1$, then $|z^n/n| = 1/n \rightarrow 0$ as $n \rightarrow \infty$, so $\lim_{n \rightarrow \infty} \frac{z^n}{n} = 0$ in this case also. If $|z| > 1$, then $|\frac{z^n}{n}| = |z|^n/n$ is unbounded (for example, by l'Hôpital's rule) and hence the sequence is divergent.

As $\frac{z^{n+1}/(n+1)!}{z^n/n!} = \frac{z}{n+1} \rightarrow 0$ for $n \rightarrow \infty$ the series $\sum_{n=0}^{\infty} \frac{z^n}{n!}$ converges for all z by the ratio test. Hence $\lim_{n \rightarrow \infty} \frac{z^n}{n!} = 0$ for all $z \in \mathbb{C}$.

Notice that $n^n > n!$ and so $|z^n|/n! > |z^n|/n^n$ for all n . Since $z^n/n!$ converges

to 0, it follows that z^n/n^n converges to 0 for all z as well.

The final limit is divergent as the sequence $n!|z|$ is unbounded.

[Return to Exercise 3.2 on P47](#)

Solution 3.3

Show that every convergent sequence $(z_n)_{n \in \mathbb{N}}$ of complex numbers is bounded.

Take $\varepsilon = 1$. By assumption, there exists $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$ we have $|z - z_n| < \varepsilon = 1$. Apply the reverse triangle inequality from Sheet 1 to obtain from this

$$|z_n| - |z| \leq |z_n - z| < 1.$$

Hence $|z_n| \leq C = \max\{|z| + 1, |z_0|, \dots, |z_{n_0}|\}$.

[Return to Exercise 3.3 on P47](#)

Solution 3.4

Let $\sum_{k=0}^{\infty} z_k$ be a convergent series of complex numbers. Show that $\lim_{k \rightarrow \infty} z_k = 0$.

By definition, the sequence of partial sums $S_n = \sum_{k=0}^n z_k$ convergent, hence a Cauchy sequence. In particular, taking $m = n + 1$ in the definition of a Cauchy sequence, we have $|z_n| = |S_{n+1} - S_n| \rightarrow 0$ as $n \rightarrow \infty$.

[Return to Exercise 3.4 on P47](#)

Solution 3.5

For which $z \in \mathbb{C}$ do the following series converge?

$$\sum_{k=0}^{\infty} kz^k, \quad \sum_{k=0}^{\infty} (kz)^k$$

Suppose the series $\sum_{k=0}^{\infty} kz^k$ converges for z . Then by (a) and (b), $k|z|^k \leq C$ is bounded. Recall that $\lim_{k \rightarrow \infty} \sqrt[k]{x} = 1$ for all $x > 0$ and that $\lim_{k \rightarrow \infty} \sqrt[k]{k} = 1$. Hence

$$|z| \leq \frac{\sqrt[k]{C}}{\sqrt[k]{k}} \rightarrow 1 \quad \text{as } k \rightarrow \infty.$$

It follows that the series can only possibly converge for $|z| \leq 1$. We can alternatively use the ratio test to see that convergence fails for $|z| > 1$. If $|z| = 1$, then kz^k is not a null sequence (the modulus is always 1), hence the series does not converge by (b). If $|z| < 1$, the series converges by the ratio test:

$$\left| \frac{(k+1)z^{k+1}}{kz^k} \right| = \left(1 + \frac{1}{k} \right) |z| \rightarrow |z| < 1, \quad \text{as } k \rightarrow \infty.$$

For the next part, suppose that the series $\sum_{k=0}^{\infty} (kz)^k$ converges for z . Then by (a) and (b), $|kz|^k \leq C$ is bounded. Increasing C , we may suppose that $C \geq 1$ and then $\sqrt[k]{C} \leq C$. Now

$$|z| \leq \frac{\sqrt[k]{C}}{k} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

This implies $z = 0$ and the series indeed converges in this case.

[Return to Exercise 3.5 on P47](#)

Solution 3.6

Let $f: \mathbb{C} \rightarrow \mathbb{C}$ be a complex function. Show that

$$\lim_{z \rightarrow \infty} f(z) = w \iff \lim_{z \rightarrow 0} f\left(\frac{1}{z}\right) = w.$$

The left hand side means that for all $\varepsilon > 0$ there exists $M > 0$ such that $|f(z) - w| < \varepsilon$ for all $+\infty > |z| > M$. The right hand side means that for all $\varepsilon > 0$ there exists $\delta > 0$ such that $|f(1/u) - w| < \varepsilon$ for all $0 < |u| < \delta$.

Upon setting $z = 1/u$ the inequalities $+\infty > |z| > M$ and $0 < |u| < \delta$ are equivalent for $M = 1/\delta$.

[Return to Exercise 3.6 on P48](#)

Solution 3.7

The Riemann sphere is $\mathbb{S} = \{(a, b, c) \in \mathbb{R}^3 \mid a^2 + b^2 + c^2 = 1\}$. Show that the stereographic projection

$$F: \mathbb{S} \longrightarrow \mathbb{C} \cup \{\infty\}, F(a, b, c) = \begin{cases} \frac{a+ib}{1-c} & \text{if } c \neq 1, \\ \infty & \text{if } c = 1. \end{cases}$$

is a bijection between the Riemann sphere and the extended complex plane. Find a formula for the inverse function.




Figure A.1: Stereographic projection from the north pole

We use the sketch to find a putative inverse $G: \mathbb{C} \cup \{\infty\} \rightarrow \mathbb{S}$. The line from $(0, 0, 1)$ to $z = x + iy \in \mathbb{C}$ is parameterized as

$$\ell(t) = (1 - t)(0, 0, 1) + t(x, y, 0) = (tx, ty, 1 - t), \quad t \in [0, 1].$$

To find the intersection points with $\mathbb{S}$, we need to solve

$$1 = (tx)^2 + (ty)^2 + (1 - t)^2 = t^2(|z|^2 + 1) - 2t + 1$$

for t . The solutions to this quadratic equation are $t = 0$ and $t = \frac{2}{1 + |z|^2}$. Hence

$$G(z) = \ell\left(\frac{2}{1 + |z|^2}\right) = \left(\frac{2x}{1 + |z|^2}, \frac{2y}{1 + |z|^2}, \frac{|z|^2 - 1}{1 + |z|^2}\right)$$

for all $z \in \mathbb{C}$. Moreover, according to the sketch we should define $G(\infty) = (0, 0, 1)$. The verifications $F \circ G = \text{id}_{\mathbb{S}}$ and $G \circ F = \text{id}_{\mathbb{C} \cup \{\infty\}}$ are left to the reader.

[Return to Exercise 3.7 on P48](#)

A.4 Chapter 4 solutions

Solution 4.1

Write the following complex functions in the form $f = u + iv$:

$$f_1(z) = \sin(z), \quad f_2(z) = e^{z^2}, \quad f_3(z) = \cosh(z)$$

By definition, $f_1(z) = \frac{e^{iz} - e^{-iz}}{2i}$. Let $z = x + iy$. Then

$$\begin{aligned} u_1 = \Re(f_1(z)) &= \frac{f_1(z) + \overline{f_1(z)}}{2} = \frac{1}{2} \left(\frac{e^{iz} - e^{-iz}}{2i} - \frac{e^{-i\bar{z}} - e^{i\bar{z}}}{2i} \right) \\ &= \frac{e^{-y}e^{ix} - e^ye^{-ix} - e^{-y}e^{-ix} + e^ye^{ix}}{4i} \\ &= \frac{e^{ix} - e^{-ix}}{2i} \cdot \frac{e^y + e^{-y}}{2} = \sin(x) \cosh(y). \end{aligned}$$

Similarly,

$$v_1 = \cos(x) \sinh(y).$$

Next,

$$f_2(z) = e^{z^2} = e^{x^2 - y^2 + 2ixy} = \underbrace{\cos(x^2 - y^2)}_{=u_2} + i \underbrace{\sin(2xy)}_{=v_2}.$$

The function f_3 is treated in the same way as f_1 . The result is

$$u_3 = \cosh(x) \cos(y), \quad v_3 = \sinh(x) \sin(y).$$

[Return to Exercise 4.1 on P60](#)

Solution 4.2

At which points $z \in \mathbb{C}$ are the following functions complex differentiable? At which points are they real differentiable?

$$f_1(z) = z,$$

$$f_2(z) = \bar{z},$$

$$f_3(z) = z^3 + z,$$

$$f_4(z) = \frac{1}{2iz},$$

$$f_5(z) = |z|^2,$$

$$f_6(z) = \frac{|z|^2}{\bar{z}}$$

f_1, f_3 and $f_6(z) = z$ are complex differentiable at every $z \in \mathbb{C}$, which implies real differentiability at these points.

f_4 is holomorphic on $\mathbb{C} \setminus \{0\}$.

$f_2 = u_2 + iv_2$ is not complex differentiable at any $z \in \mathbb{C}$ since the Cauchy–Riemann equations ($\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$ and $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$) are violated at every $(x, y) \in \mathbb{C}$. Indeed, $u_2 = x$, $v_2 = -y$ and $\frac{\partial u_2}{\partial x} = 1 \neq -1 = \frac{\partial v_2}{\partial y}$. Of course, f_2 is real differentiable on all of $\mathbb{R}^2$ since both component functions $u_2(x, y) = x$, $v_2(x, y) = -y$ are evidently differentiable.

$f_5 = u_5 + iv_5$ is complex differentiable at the origin but not anywhere else. This follows by the Cauchy–Riemann equations, as $u_5 = x^2$, $v_5 = y^2$ and

$$\frac{\partial u_5}{\partial x} = 2x, \quad \frac{\partial u_5}{\partial y} = 0, \quad \frac{\partial v_5}{\partial x} = 0, \quad \frac{\partial v_5}{\partial y} = 2y.$$

f_5 is real differentiable on all of $\mathbb{R}^2$ since both component functions $u_5(x, y) = x^2$, $v_5(x, y) = y^2$ are evidently differentiable.

[Return to Exercise 4.2 on P60](#)

Solution 4.3

Let $f: V \rightarrow W$ be a bijection of open sets $V, W \subset \mathbb{C}$. Assume that f is complex differentiable at z_0 and that f^{-1} is complex differentiable at $w_0 = f(z_0)$. Show that $f'(z_0) \neq 0$. (The same result holds for real differentiable maps to show $\det J_f(z_0) \neq 0$.)

Let $g = f^{-1}$ so that $g(f(z)) = z$ for all $z \in V$. By the chain rule, $g \circ f$ is complex differentiable at z_0 with derivative

$$g'(f(z_0))f'(z_0) = (gf)'(z_0) = 1.$$

Hence $f'(z_0) \neq 0$.

[Return to Exercise 4.3 on P60](#)

Solution 4.4

Let $f: U \rightarrow \mathbb{C}$ be a holomorphic function with domain an open set $U \subset \mathbb{C}$. Suppose also that $f'(z)$ is holomorphic. Write $f(z) = u(x, y) + iv(x, y)$, where $z = x + iy$. As we will see, a consequence of f being holomorphic, is that u and v have continuous second order derivatives. Show that:

- $|f'(z)|^2 = \left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial v}{\partial x}\right)^2 = \left(\frac{\partial u}{\partial y}\right)^2 + \left(\frac{\partial v}{\partial y}\right)^2$
- Both u, v satisfy the Laplace equation $\Delta(u) = \Delta(v) = 0$, where the Laplace operator is defined as $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$.
- Fix $n \in \mathbb{N}$. Use b. to find a solution to $\Delta(u) = 0$ that satisfies $u(z) = \cos(n\theta)$ for all $|z| = 1$, $z = e^{i\theta}$ on the unit circle.
- Does the converse to b. hold?

Since f is holomorphic, the Cauchy–Riemann equations

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \quad (\star)$$

hold at every $z \in U$. For a., compute

$$\begin{aligned}
|f'(z)|^2 &= \left| \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \right|^2 && \text{by Prop. 4.5} \\
&= \left| \frac{\partial u}{\partial x} \right|^2 + \left| \frac{\partial v}{\partial x} \right|^2 \\
&= \left| \frac{\partial v}{\partial y} \right|^2 + \left| \frac{-\partial u}{\partial y} \right|^2 && \text{by } (\star) \\
&= \left| \frac{\partial u}{\partial y} \right|^2 + \left| \frac{\partial v}{\partial y} \right|^2.
\end{aligned}$$

For b., we take the partial derivative $\frac{\partial}{\partial x}$ of the first equation in $(\star)$ to get

$$\begin{aligned}
\frac{\partial^2 u}{\partial x^2} &= \frac{\partial^2 v}{\partial x \partial y} \\
&= \frac{\partial}{\partial y} \frac{\partial v}{\partial x} && \text{symmetry of 2nd derivatives} \\
&= -\frac{\partial}{\partial y} \frac{\partial u}{\partial y} && \text{by second equation in } (\star) \\
&= -\frac{\partial^2 u}{\partial y^2}.
\end{aligned}$$

Similarly, take the partial derivative $\frac{\partial}{\partial x}$ of the second equation in $(\star)$ to get

$$\frac{\partial^2 v}{\partial x^2} = -\frac{\partial^2 u}{\partial x \partial y} = -\frac{\partial^2 u}{\partial y \partial x} = -\frac{\partial^2 v}{\partial y^2}.$$

Note that the symmetry of the 2nd derivatives is a consequence of continuity of the second order derivatives.

For c., let $f(z) = z^n$. Then $u = \Re(z^n) = \frac{z^n + \bar{z}^n}{2} = r^n \cos(n\theta)$ solves Laplace's equation by part (b). Also, $r = |z| = 1$ implies $u(x, y) = \cos(n\theta)$, as required.

For d., No, the converse fails. Take for example $f(z) = \bar{z}$. Then $f = u + iv$ with $u = \bar{x}$ and $v = -\bar{y}$.

and v satisfy Laplace's equation but f is not holomorphic.

[Return to Exercise 4.4 on P60](#)

Solution 4.5

Find subdomains where $\sin, \cos, \sinh, \cosh, \tan, \tanh$ are injective. Compute the derivative of an inverse using the chain rule. Deduce an explicit formula for the derivative of an inverse.

$\sin z = \sin w$ implies $e^{iz} - e^{-iz} = e^{iw} - e^{-iw} = a$.

Set $t = e^{iz}$, and $s = e^{iw}$ we find that t and s are roots of the polynomial:

$$x^2 - xa - 1 = 0.$$

This polynomial has two roots, $x = \{r, \frac{1}{r}\}$.

We conclude that either $s = t$ or $st = 1$.

If $s = t$, then $e^{-y}e^{ix} = e^{-v}e^{iu}$. We conclude that $y = v$ and $x - u = 2k\pi$.

If $st = 1$ then, $e^{-y}e^{ix} = e^v e^{-iu}$. We conclude that $v = -y$ and $x + u = (2k + 1)\pi$. Where in both cases $k \in \mathbb{Z}$. In particular, $\sin z$ is injective on the domain $\{x + iy : x \in (0, \pi)\}$.

In a similar way $\cos(z)$ is also injective on the domain $\{x + iy : x \in (0, \pi)\}$

We can then deduce that $\sinh(z)$ and $\cosh(z)$ are injective on the domain $\{x + iy : y \in (0, \pi)\}$.

For $\tanh$: $\tanh z = \tanh w$ implies $e^{2(z-w)} = 1$. Writing $z = x + iy$ and $w = u + iv$, we have $2(x - u) = 0$ and $2(y - v) = 0 + 2k\pi$. We conclude that $\tanh$ is injective on the interval $\{x + iy : y \in (0, \pi)\} \setminus \{i(\pi/2 + k\pi) : k \in \mathbb{Z}\}$.

We can then deduce that $\tan(z) = \tanh(iz)$ is injective on $\{x + iy : x \in (0, \pi)\} \setminus \{(\pi/2 + k\pi) : k \in \mathbb{Z}\}$.

Using the chain rule, and the identities $\cos^2 + \sin^2 = 1$, $\cosh^2 - \sinh^2 = 1$, we have

$$\arcsin'(z) = \frac{1}{\cos(\arcsin(z))} = \frac{1}{\sqrt{1-z^2}}$$

$$\arccos'(z) = \frac{-1}{\sin(\arccos(z))} = \frac{-1}{\sqrt{1-z^2}}$$

$$\operatorname{arcsinh}'(z) = \frac{1}{\cosh(\operatorname{arcsinh}(z))} = \frac{1}{\sqrt{1+z^2}}$$

$$\operatorname{arccosh}'(z) = \frac{1}{\sinh(\operatorname{arccosh}(z))} = \frac{1}{\sqrt{z^2-1}}$$

[Return to Exercise 4.5 on P61](#)

Solution 4.6

Show that the set $\mathcal{O}(U)$ of holomorphic functions on U becomes a ring under the operations of point-wise addition and multiplication:

$$(f+g)(z) = f(z) + g(z),$$

$$(fg)(z) = f(z)g(z).$$

Since we already know that the set of all functions $U \rightarrow \mathbb{C}$ is a ring, it suffices to show that $\mathcal{O}(U)$ is a subring. As constant functions are entire, $\pm 1 \in \mathcal{O}(U)$. If

f, g are holomorphic U , then by the lecture notes $f + g$ and fg are holomorphic on U as well. Finally, $-f = -1 \cdot f$ is holomorphic on U .

[Return to Exercise 4.6 on P61](#)

Solution 4.7

Let $f: \mathbb{C} \rightarrow \mathbb{C}$ be an entire function with image $f(\mathbb{C}) \subset \mathbb{R}$. Prove that f is constant.

Writing $f = u + iv$, we have $v = 0$. Hence by the Cauchy–Riemann equations it follows that $\partial u / \partial x = 0$ and $\partial u / \partial y = 0$. This implies that u is constant: for every fixed $y_0 \in \mathbb{R}$ the real-valued differentiable function $\mathbb{R} \rightarrow \mathbb{R}$, $x \mapsto u(x, y_0)$ has an everywhere vanishing derivative and is therefore a constant $u(x, y_0) = c(y_0)$, $\forall x \in \mathbb{R}$, with the constant $c(y_0)$ possibly depending on $y_0 \in \mathbb{R}$. For every fixed $x_0 \in \mathbb{R}$ we similarly have that $u(x_0, y) = d(x_0)$ is a constant of y . Fix $(x_0, y_0) \in \mathbb{C}$. Then

$$u(x, y) = d(x) = u(x, y_0) = c(y_0) = u(x_0, y_0)$$

for all $(x, y) \in \mathbb{C}$, so $u(x, y)$ is constant.

[Return to Exercise 4.7 on P61](#)

Solution 4.8

Let U be an open set and assume that U is path-connected, meaning that for all $z_0, z_1 \in U$ there exists a continuously differentiable map $\gamma: [0, 1] \rightarrow U$ such that $\gamma(0) = z_0$, $\gamma(1) = z_1$.

- a. Let $f: U \rightarrow \mathbb{C}$ be a holomorphic function such that $f'(z) = 0$ for all $z \in U$. Prove that $f(z)$ is a constant function.**

Hint: Consider the real and imaginary parts of $f \circ \gamma$.

b. Show that the only entire functions $f(z)$ satisfying $f'(z) = Cf(z)$ for a constant $C \in \mathbb{C}$ are the functions $f(z) = De^{Cz}$, $D \in \mathbb{C}$.

c. Find all entire solutions $f(z)$ of $f''(z) = f(z)$.

Hint: Consider $g = f + f'$, $h = f - f'$

a. Let $f = u + iv$ and fix $z_0 \in U$. We will prove that $f(z) = f(z_0)$ for all $z \in U$. Pick a continuously differentiable map $\gamma: [0, 1] \rightarrow U$ with $\gamma(0) = z_0$, $\gamma(1) = z$. By Proposition 4.5 (the Cauchy–Riemann equations), u, v are real differentiable on U with vanishing derivatives. Hence $u \circ \gamma$ and $v \circ \gamma$ are real functions $[0, 1] \rightarrow \mathbb{R}$ with vanishing derivatives which we know are constant from calculus. Hence

$$f(z_0) = u(\gamma(0)) + iv(\gamma(0)) = u(\gamma(1)) + iv(\gamma(1)) = f(z).$$

b. Let $g(z) = f(z)e^{-Cz}$. Then $g'(z) = f'(z)e^{-Cz} + f(z)(-Ce^{-Cz}) = 0$. Hence $f(z)e^{-Cz} = g(z) = D$ is constant.

c. Since $g' = g$, $h' = -h$ we know from (b) that $g(z) = Ce^z$, $h(z) = De^{-z}$. Hence $f(z) = \frac{1}{2}(g(z) + h(z)) = \frac{Ce^z + De^{-z}}{2}$ is the general solution. Taking $(C, D) \in \{(1, 1), (1, -1)\}$ gives $\cosh(z)$, $\sinh(z)$ and we could alternatively say that the general solution is $A \cosh(z) + B \sinh(z)$.

[Return to Exercise 4.8 on P62](#)

A.5 Chapter 5 solutions

Solution 5.1

Find the radius of convergence for the following power series centered at the origin.

- i. $\sum_{n=0}^{\infty} \frac{z^n}{n^3},$
- ii. $\sum_{n=0}^{\infty} z^{3n},$
- iii. $\sum_{n=0}^{\infty} \frac{z^n}{n^n}.$

Hint: Recall the ratio and root tests for series of complex numbers

- i. We can apply the *ratio test* to determine the convergence of a series of complex numbers $\sum_{n=0}^{\infty} a_n$ provided $a_n \neq 0$ for n sufficiently large. Let $q = \limsup \left| \frac{a_{n+1}}{a_n} \right|$. If $q < 1$, then $\sum_{n=0}^{\infty} a_n$ converges absolutely. If $q > 1$, then $\sum_{n=0}^{\infty} a_n$ diverges. Apply this to $a_n = \frac{z^n}{n^3}$:

$$\left| \frac{z^{n+1}/(n+1)^3}{z^n/n^3} \right| = |z| \left(\frac{n}{n+1} \right)^3 \rightarrow |z| \quad \text{as } n \rightarrow \infty.$$

Hence, $q = |z|$ and (i) converges for $|z| < 1$ and diverges for $|z| > 1$, so $\rho = 1$.

- ii. By definition, the radius of convergence ρ is the supremum of all real numbers $r \geq 0$ such that the sequence $(a_n r^n)_{n \in \mathbb{N}}$ is bounded. Here $a_n = 1$ if 3 divides n and $a_n = 0$ otherwise. This is equivalent to the boundedness of the sequence $(r^{3n})_{n \in \mathbb{N}}$ which holds precisely for $r \leq 1$. Thus $\rho = 1$.

- iii. Recall the *root test* to determine the convergence of a series of complex numbers $\sum_{n=0}^{\infty} a_n$. Let $q = \limsup \sqrt[n]{|a_n|}$. If $q < 1$, then $\sum_{n=0}^{\infty} a_n$ converges absolutely. If $q > 1$, then $\sum_{n=0}^{\infty} a_n$ diverges. Apply this to $a_n = \frac{z^n}{n^n}$:

$$\sqrt[n]{|a_n|} = |z|/n \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Hence iii. converges for all $z \in \mathbb{C}$, and so $\rho = +\infty$.

[Return to Exercise 5.1 on P74](#)

Solution 5.2

Treating e^z , $\sin(z)$, $\cos(z)$ as formal power series with their usual Taylor expansion, find the terms of order ≤ 3 of the following power series:

- i. $e^z \sin(z)$,
- ii. $\sin(z) \cos(z)$,
- iii. $1/\cos(z)$

Using the definition of the product of formal power series,

$$\begin{aligned}e^z \sin(z) &= z + z^2 + \frac{z^3}{3} + O(z^4) \\ \sin(z) \cos(z) &= z - \frac{2z^3}{3} + O(z^4).\end{aligned}$$

To verify

$$1/\cos(z) = 1 + \frac{z^2}{2} + O(z^4)$$

we begin with

$$\cos(z) = 1 - z^2/2 + O(z^4)$$

and then using the geometric series,

$$1/\cos(z) = \frac{1}{1 - z^2/2} + O(z^4) = 1 + z^2/2 + O(z^4).$$

[Return to Exercise 5.2 on P74](#)

Solution 5.3

Let $P = \sum_{n=0}^{\infty} a_n z^n$, $Q = \sum_{n=0}^{\infty} b_n z^n$ be power series with positive radii of convergence $\rho_P, \rho_Q > 0$. Show that:

- a. $P + Q = \sum_{n=0}^{\infty} (a_n + b_n) z^n$ has radius of convergence $\rho \geq \min(\rho_P, \rho_Q)$.
 b. $PQ = \sum_{n=0}^{\infty} \left(\sum_{i+j=n} a_i b_j \right) z^n$ has radius of convergence $\rho \geq \min(\rho_P, \rho_Q)$.

a. Let $z \in D_{\min(\rho_P, \rho_Q)}(0) = D_{\rho_P}(0) \cap D_{\rho_Q}(0)$. Then the series of complex numbers $\sum_{n=0}^{\infty} a_n z^n$ and $\sum_{n=0}^{\infty} b_n z^n$ converge, hence the series $\sum_{n=0}^{\infty} (a_n + b_n) z^n$ converges as well. It follows that $D_{\min(\rho_P, \rho_Q)}(0) \subset D(P + Q, 0)$ is in the domain of $P + Q$, hence $\min(\rho_P, \rho_Q) \leq \rho$.

b. Recall that the Cauchy product of series of non-negative numbers converges: If $a_n, b_n \geq 0$ and $\sum_{n=0}^{\infty} a_n$, $\sum_{n=0}^{\infty} b_n$ converge, then $\sum_{n=0}^{\infty} \left(\sum_{i+j=n} a_i b_j \right)$ converges. To see this, let $A_N = \sum_{n=0}^N a_n$, $B_N = \sum_{n=0}^N b_n$ be the partial sums. Then the partial sums of the Cauchy product

$$\sum_{n=0}^N \sum_{i+j=n} a_i b_j \leq A_N B_N$$

are a bounded sequence (since $A_N B_N$ converges) that is increasing, hence convergent by the monotone convergence theorem of calculus.

Given this preliminary, the same argument as for a. proves $\min(\rho_P, \rho_Q) \leq \rho$.

[Return to Exercise 5.3 on P74](#)

Solution 5.4

Find a solution to the non-linear differential equation

$$f'(z) + f(z)^2 = 0, \quad f(0) = 1$$

on a disk centered at $z_0 = 0$ by making the ansatz $f(z) = \sum_{n=0}^{\infty} a_n z^n$, inductively determining the coefficients a_n , and finding the radius of convergence.

We substitute the power series into the differential equation. By Theorem 5.2, we have

$$f'(z) = \sum_{n=0}^{\infty} (n+1)a_{n+1}z^n$$

and for the product of power series (see Equation 5.3) we have

$$f(z)^2 = \sum_{n=0}^{\infty} \left(\sum_{i=0}^n a_i a_{n-i} \right) z^n.$$

Hence

$$\sum_{n=0}^{\infty} \left[(n+1)a_{n+1} + \sum_{i=0}^n a_i a_{n-i} \right] z^n = 0.$$

The identity theorem (Corollary 5.2) now implies

$$a_{n+1} = \frac{-1}{n+1} \sum_{i=0}^n a_i a_{n-i}, \quad \forall n \geq 0.$$

By assumption, $a_0 = f(0) = 1$. We solve by recursion,

$$\begin{aligned} a_0 &= 1 \\ a_1 &= -a_0^2 = -1 \\ a_2 &= \frac{-1}{2}(a_0 a_1 + a_1 a_0) = 1 \\ a_3 &= \frac{-1}{3}(a_0 a_2 + a_1^2 + a_2 a_0) = -1 \\ &\vdots \end{aligned}$$

This leads us to the conjecture $a_n = (-1)^n$ for all $n \in \mathbb{N}$. This holds in the

base case $n = 0$ and, assuming this result for all integers up to n , we have

$$a_{n+1} = \frac{-1}{n+1} \sum_{i=0}^n (-1)^i (-1)^{n-i} = \frac{-1}{n+1} \sum_{i=0}^n (-1)^n = (-1)^{n+1},$$

which proves the inductive step. In summary,

$$f(z) = \sum_{n=0}^{\infty} (-1)^n z^n,$$

which we recognize as a geometric series with $f(z) = 1/(1+z)$, whose radius of convergence we know is $\rho = 1$.

[Return to Exercise 5.4 on P75](#)

Solution 5.5

Let $P(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series centered at $z_0 = 0$ and assume that the radius of convergence $\rho > 0$ is positive. Suppose that $P(z) \in \mathbb{R}$ for all $z \in D_\rho(0) \cap \mathbb{R}$. Prove that all coefficients a_n , $n \in \mathbb{N}$, must be real numbers. Deduce that $\overline{P(z)} = P(\bar{z})$ for all $z \in D_\rho(0)$.

For all $z \in D_\rho(0) \cap \mathbb{R}$ we have by assumption the middle equality in

$$\sum_{n=0}^{\infty} a_n z^n = P(z) = \frac{P(z) + \overline{P(z)}}{2} = \sum_{n=0}^{\infty} \frac{a_n + \overline{a_n}}{2} z^n.$$

Pick $n_0 \in \mathbb{N}$ with $1/n_0 < \rho$. We can apply the identity theorem (Corollary 5.2) with $w_n = 1/(n + n_0)$, P as above, and

$$Q = \sum_{n=0}^{\infty} \frac{a_n + \overline{a_n}}{2} z^n$$

to conclude that $a_n = \frac{a_n + \overline{a_n}}{2}$ for all n . In other words, $a_n \in \mathbb{R}$.

We may then deduce that

$$\overline{P(z)} = \overline{\sum_{n=0}^{\infty} a_n z^n} \stackrel{(\S)}{=} \sum_{n=0}^{\infty} \overline{a_n z^n} = \sum_{n=0}^{\infty} a_n \bar{z}^n = P(\bar{z}).$$

Here (§) is easy to check from the definition of convergence of partial sums, as

$$\overline{\sum_{n=0}^N a_n z^n} = \sum_{n=0}^N \overline{a_n z^n}$$

holds for finite sums.

[Return to Exercise 5.5 on P75](#)

Solution 5.6

Prove that the ring of formal power series $\mathbb{C}[[T]]$ is an integral domain. In other words, show that $PQ = 0 \implies P = 0$ or $Q = 0$.

Suppose that $Q \neq 0$ and $PQ = 0$. We must show that $P = 0$. Write $Q = z^n R$ where R has non-vanishing constant term. According to Proposition 5.4(b), R is invertible, that is, we can find $S \in \mathbb{C}[[T]]$ such that

$$RS = 1, \quad SR = 1.$$

Then

$$0 = PQS = Pz^n RS = Pz^n.$$

This implies $P = 0$ (the coefficients of P and Pz^n are just shifted by n).

[Return to Exercise 5.6 on P75](#)

Solution 5.7

The binomial coefficient of a complex number $\alpha \neq 0$ and $k \in \mathbb{N}$ is defined as

$$\binom{\alpha}{k} = \begin{cases} \frac{\alpha(\alpha-1)\cdots(\alpha-k+1)}{k!} & \text{if } k > 0, \\ 1 & \text{if } k = 0. \end{cases}$$

The binomial series is the formal power series

$$B_\alpha = \sum_{k=0}^{\infty} \binom{\alpha}{k} T^k.$$

- a. Determine the radius of convergence of B_α .
- b. Prove that if $\alpha \in \mathbb{N}$, then $B_\alpha(z) = (z+1)^\alpha$.
- c. Show the generalized Vandermonde identity for $\alpha, \beta, \alpha + \beta \in \mathbb{C}^\times$,

$$\sum_{k=0}^n \binom{\alpha}{k} \binom{\beta}{n-k} = \binom{\alpha+\beta}{n}. \quad (\star)$$

- d. Prove that for $\alpha = 1/k$ the complex function $B_\alpha(z)$ satisfies $B_\alpha(z)^k = z+1$ on its domain. Hence $B_\alpha(z)$ is a k -th root of the function $z+1$.

- a. For $\alpha \in \mathbb{C} \setminus \mathbb{N}$ we have

$$\left| \frac{\binom{\alpha}{k+1} z^{k+1}}{\binom{\alpha}{k} z^k} \right| = \left| z \frac{\alpha - k}{k+1} \right| \rightarrow |z| \quad \text{as } k \rightarrow \infty.$$

Therefore, the ratio test implies $\rho = 1$ when $\alpha \notin \mathbb{N}$. When $\alpha \in \mathbb{N}$ only finitely many binomial coefficients $\binom{\alpha}{k}$ are non-zero, so B_α is a polynomial and $\rho = +\infty$ and indeed by the binomial theorem we have

$$B_\alpha(z) = (z+1)^\alpha, \quad \forall \alpha \in \mathbb{N}.$$

b. As $\binom{\alpha}{k} = 0$ if $k > \alpha$, this is just the binomial theorem:

$$\sum_{k=0}^{\alpha} \binom{\alpha}{k} z^k = (z + 1)^{\alpha}.$$

c. We can deduce from b. that $(\star)$ holds for all $\alpha, \beta \in \mathbb{N}$, since by the definition of the product of formal power series, we have

$$\begin{aligned} \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{\alpha}{k} \binom{\beta}{n-k} \right) z^n &= \sum_{k=0}^{\infty} \binom{\alpha}{k} z^k \cdot \sum_{\ell=0}^{\infty} \binom{\beta}{\ell} z^{\ell} \\ &= (1+z)^{\alpha} (1+z)^{\beta} = (1+z)^{\alpha+\beta} \\ &= \sum_{n=0}^{\infty} \binom{\alpha+\beta}{n} z^n. \end{aligned}$$

Fix $\beta \in \mathbb{N}$. Then both sides of $(\star)$ are polynomials in α , which agree on $\mathbb{N}$. Since a non-zero polynomial has only finitely many roots, this implies that both sides of $(\star)$ agree for all $\alpha \in \mathbb{C}$. Now fix $\alpha \in \mathbb{C}$. Then both sides of $(\star)$ are polynomials in β , which have just shown agree for all $\beta \in \mathbb{N}$. As before, this implies that $(\star)$ holds for all $\beta \in \mathbb{C}$. This proves the general case.

d. From c. and the definition of the product of power series, we have

$$B_{\alpha}(z)B_{\beta}(z) = B_{\alpha+\beta}(z).$$

By induction,

$$B_{\alpha}(z)^k = B_{k\alpha}(z) = B_1(z) = z + 1.$$

[Return to Exercise 5.7 on P75](#)

Solution 5.8

For $a > 0$ and $z \in \mathbb{C}$ define $a^z = \exp(\log(a)z)$. Show that:

- a.** $a^z b^z = (ab)^z$ for all $a, b > 0, z \in \mathbb{C}$
- b.** $a^z a^w = a^{z+w}$ for all $a > 0, z, w \in \mathbb{C}$
- c.** $|a^z| = a^{\Re(z)}$ for all $a > 0, z \in \mathbb{C}$

The Riemann ζ -function is defined as

$$\zeta(z) = \sum_{n=1}^{\infty} \frac{1}{n^z} \quad \text{for } \Re(z) > 1. \quad (*)$$

- d.** Prove that the series $(*)$ converges absolutely for all $z \in \mathbb{C}$ with $\Re(z) > 1$ and uniformly on every subset $S_\delta = \{z \in \mathbb{C} \mid \Re(z) > 1 + \delta\}$ with $\delta > 0$.

a. We have:

$$\begin{aligned} a^z b^z &= \exp(\log(a)z) \exp(\log(b)z) \\ &= \exp([\log(a) + \log(b)]z) \\ &= \exp(\log(ab)z) = (ab)^z \end{aligned}$$

b.

$$a^z a^w = \exp(\log(a)z) \exp(\log(a)w) = \exp(\log(a)(z + w)) = a^{z+w}$$

- c. Notice first that $|\exp(z)|^2 = \exp(z) \overline{\exp(z)} = \exp(z) \exp(\bar{z}) = \exp(z + \bar{z}) = \exp(2\Re(z))$. Hence $|\exp(z)| = \exp(2\Re(z))^{1/2} = \exp(\Re(z))$. Using

this,

$$|a^z| = |\exp(\log(a)z)| = \exp(\Re(\log(a)z)) = \exp(\log(a)\Re(z)) = a^{\Re(z)}.$$

- d. The series of absolute values $\sum_{n=1}^{\infty} |1/n^z| = \sum_{n=1}^{\infty} 1/n^x$ converges for $x > 1 + \delta$ by the comparison test: We have $1/n^x \leq 1/n^{1+\delta}$ and the series

$$\sum_{n=1}^{\infty} \frac{1}{n^{1+\delta}} \leq \int_1^{\infty} \frac{dt}{t^{1+\delta}} = \lim_{R \rightarrow \infty} \int_1^R \frac{dt}{t^{1+\delta}} = \lim_{R \rightarrow \infty} \left(1 - \frac{1}{R^\delta}\right) = 1 \quad (\dagger)$$

converges by the integral comparison test. It follows that $(\star)$ converges uniformly. Moreover, for fixed $\delta > 0$ the series $(\dagger)$ that we compare with is independent of $z \in S_\delta$. Thus the comparison test for series of functions applies and implies uniform convergence on each S_δ , $\delta > 0$.

[Return to Exercise 5.8 on P76](#)

A.6 Chapter 6 solutions

Solution 6.1





Recall from Example 4.6 that $\log: \mathbb{C}^- \rightarrow \mathbb{C}$ is a holomorphic primitive of $1/z$ on the slit plane $\mathbb{C}^-$. Combine Equation 6.9 and Example 6.3 to show that there is no holomorphic primitive of $1/z$ on the punctured plane $\mathbb{C}^\times$.

The punctured plane is an open subset of $\mathbb{C}$ on which $1/z$ is continuous. If $1/z$ has a holomorphic primitive F on $\mathbb{C}^\times$, then for any closed curve γ $\int_\gamma 1/z dz = 0$. However, by Example 6.3 there is a closed curve γ for which $\int_\gamma z^{-1} = 2\pi i \neq 0$. This gives the desired contradiction.

[Return to Exercise 6.1 on P88](#)

Solution 6.2

For the curves γ_k , $k = 1, 2, 3, 4$, as in the following sketch



- find piecewise C^1 parameterisations $\gamma_k: [a, b] \rightarrow \mathbb{C}$,
- compute the length $L(\gamma_k)$ (γ_3 may be omitted)
- evaluate $\int_{\gamma_k} z^2 dz$.

a. The parameterisations are

$$\gamma_1: [0, 1] \longrightarrow \mathbb{C}, \quad \gamma_1(t) = t(-2 + i),$$

$$\gamma_2: [0, 3] \longrightarrow \mathbb{C}, \quad \gamma_2(t) = \begin{cases} ti + (1-t)(1-i) & \text{if } t \in [0, 1], \\ (t-1)(-1-i) + (2-t)i & \text{if } t \in [1, 2], \\ (t-2)(1-i) - (3-t)(1+i) & \text{if } t \in [2, 3], \end{cases}$$

$$\gamma_3: [0, 2] \longrightarrow \mathbb{C}, \quad \gamma_3(t) = te^{-4\pi it},$$

$$\gamma_4: [-\pi, 0] \longrightarrow \mathbb{C}, \quad \gamma_4(t) = e^{-it}.$$

b. By the Pythagorean theorem, $L(\gamma_1) = \sqrt{2^2 + 1^2} = \sqrt{5}$. Similarly, $L(\gamma_2) = 2 + 2\sqrt{5}$. Compute

$$\begin{aligned} L(\gamma_3) &= \int_0^2 |\gamma_3'(t)| dt = \int_0^2 \sqrt{1 + 16\pi^2 t^2} dt \\ &= \left[\frac{4\pi\sqrt{16\pi^2 t^2 + 1}t + \sinh^{-1}(4\pi t)}{8\pi} \right]_0^2 \\ &= \sqrt{1 + 64\pi^2} + \frac{\sinh^{-1}(8\pi)}{8\pi} \end{aligned}$$

Finally, $L(\gamma_4) = \pi$, since a circle has circumference $2\pi r$.

c. Since $f(z) = z^2$ has the primitive $F(z) = z^3/3$, we can apply the complex FTC (Proposition~6.12) to evaluate the curve integrals by inserting the

endpoints into the primitive:

$$\begin{aligned}\int_{\gamma_1} z^2 dz &= F(-2+i) - F(0) = -\frac{2}{3} + \frac{11}{3}i \\ \int_{\gamma_2} z^2 dz &= F(1-i) - F(1-i) = 0 \\ \int_{\gamma_3} z^2 dz &= F(2) - F(0) = 8/3 \\ \int_{\gamma_4} z^2 dz &= F(1) - F(-1) = 2/3\end{aligned}$$

[Return to Exercise 6.2 on P88](#)

Solution 6.3

Define $\gamma_\epsilon: [-\pi + \epsilon, \pi - \epsilon] \rightarrow \mathbb{C}$, $\gamma_\epsilon(t) = e^{it}$ **for** $0 < \epsilon < \pi$. **Compute**

$$\lim_{\epsilon \rightarrow 0} \int_{\gamma_\epsilon} z^{1/2} dz,$$

where the square root $z^{1/2} = \exp(\log(z)/2)$ is defined as on Sheet 5 using the principal branch of the logarithm.

(Aside: $z^{1/2}$ is integrable in the sense of measure theory and using the more general integral defined there we can rewrite the above integral as $\int_{\partial D_1(0)} z^{1/2} dz$.)

Using the principal branch of the logarithm means that

$$z = re^{i\theta}, \theta \in (-\pi, \pi) \implies \sqrt{z} = \sqrt{r}e^{i\theta/2}.$$

Applying this to $z = e^{it}$, we have

$$\int_{\gamma_\epsilon} z^{1/2} dz = \int_{-\pi+\epsilon}^{\pi-\epsilon} e^{it/2} i e^{it} dt = \left[\frac{2}{3} e^{\frac{3}{2}it} \right]_{-\pi+\epsilon}^{\pi-\epsilon} \xrightarrow{\epsilon \rightarrow 0} \frac{-4i}{3}$$

Return to Exercise 6.3 on P89

Solution 6.4

Let γ be the straight line from $z = 1$ to $z = i$. Let $f(z) = 1/z^4$. Determine the maximum of $|f(z)|$ over all $z \in \mathfrak{I}(\gamma)$. Use this to estimate

$$\left| \int_{\gamma} f(z) dz \right| \leq 4\sqrt{2}.$$

Then compute $\int_{\gamma} f(z) dz$ directly and compare this to the estimate.

Parameterise the curve as $\gamma: [0, 1] \rightarrow \mathbb{C}$, $\gamma(t) = (1 - t) + ti$. Then

$$|f(\gamma(t))| = \frac{1}{|\gamma(t)|^4} = \frac{1}{((1 - t)^2 + t^2)^2}$$

which attains its maximum 4 at $t = 1/2$ (for example, set the t -derivative equal to zero). Since $L(\gamma) = \sqrt{2}$, we conclude from Proposition 6.9(d) that

$$\left| \int_{\gamma} f(z) dz \right| \leq 4\sqrt{2}.$$

To compute the integral directly, notice that $f(z)$ has the primitive $F(z) = -\frac{1}{3}z^{-3}$, so the complex FTC gives

$$\int_{\gamma} f(z) dz = F(i) - F(1) = \frac{1 - i}{3}.$$

Since $\left|\frac{1-i}{3}\right| = \frac{\sqrt{2}}{3}$, we see that our estimate was 12 times too large.

[Return to Exercise 6.4 on P90](#)

Solution 6.5

Let $f(z): \partial D_r(z_0) \rightarrow \mathbb{C}$ be a continuous function. Show that

$$\overline{\int_{\partial D_r(z_0)} f(z) dz} = -r^2 \int_{\partial D_r(z_0)} \overline{f(z)} (z - z_0)^{-2} dz.$$

By using the definition of the curve integral and using properties of complex conjugation, we find

$$\begin{aligned} \overline{\int_{\partial D_r(z_0)} f(z) dz} &= \overline{\int_0^{2\pi} f(z_0 + re^{it}) ire^{it} dt} \\ &= - \int_0^{2\pi} \overline{f(z_0 + re^{it})} ire^{-it} dt \\ &= -r^2 \int_0^{2\pi} \overline{f(z_0 + re^{it})} (re^{it})^{-2} ire^{it} dt \\ &= -r^2 \int_{\partial D_r(z_0)} \overline{f(z)} (z - z_0)^{-2} dz. \end{aligned}$$

[Return to Exercise 6.5 on P90](#)

A.7 Chapter 7 solutions

Solution 7.1

a. Sketch the curves

$$\gamma_0: [-1, 1] \rightarrow \mathbb{C},$$


$$\gamma_0(t) = t,$$

$$\gamma_1: [-1, 1] \rightarrow \mathbb{C},$$

$$\gamma_1(t) = e^{i\pi \frac{1-t}{2}}$$

and show that they are path-homotopic.

b. Let $0 < r_0 < r_1$ and $z_0 \in \mathbb{C}$. Prove that the loops $\partial D_{r_0}(z_0)$ and $\partial D_{r_1}(z_0)$ are freely homotopic in the closed annulus $\bar{A}_{r_0, r_1}(z_0)$.



The straight-line homotopy $\Gamma_s(t) = s\gamma_1(t) + (1-s)\gamma_0(t)$ is a path-homotopy, since using $\gamma_0(\pm 1) = \gamma_1(\pm 1)$ we find that

$$\Gamma_s(\pm 1) = s\gamma_1(\pm 1) + (1-s)\gamma_0(\pm 1) = \gamma_0(\pm 1)$$

is constant in $s \in [0, 1]$.

- b. Recall that $\gamma_0: [0, 2\pi] \rightarrow \mathbb{C}$, $\gamma_0(t) = r_0 e^{it}$ and $\gamma_1: [0, 2\pi] \rightarrow \mathbb{C}$, $\gamma_1(t) = r_1 e^{it}$. We will again use a straight-line homotopy $\Gamma_s(t) = s\gamma_1(t) + (1-s)\gamma_0(t)$. To show that Γ is a homotopy in the closed annulus, we must prove that $r_1 \leq |\Gamma_s(t)| \leq r_2$ for all s, t . This holds since

$$|\Gamma_s(t)| = |(tr_1 + (1-t)r_0)e^{it}| = tr_1 + (1-t)r_0$$

is the straight-line from r_0 to r_1 , hence has image in $[r_0, r_1]$.

[Return to Exercise 7.1 on P102](#)

Solution 7.2

Let $\alpha + i\beta \in \mathbb{C}$. **Determine**

$$\int_a^b e^{(\alpha+i\beta)t} dt$$

to compute

$$\int_a^b e^{\alpha t} \cos(\beta t) dt.$$

By the FTC, $\int_a^b e^{(\alpha+i\beta)t} dt = \left[\frac{e^{(\alpha+i\beta)t}}{\alpha+i\beta} \right]_a^b$.

As a preparation, notice that for $z = x + iy$ we have

$$\Re(e^z) = \frac{e^z + e^{\bar{z}}}{2} = \frac{e^x e^{iy} + e^x e^{-iy}}{2} = e^x \cos(y).$$

Similarly, $\Im(e^z) = e^x \sin(y)$. We apply this to $z = (\alpha + i\beta)t$ and compute

$$\begin{aligned} \int_a^b e^{\alpha t} \cos(\beta t) dt &= \int_a^b \Re e^{(\alpha + i\beta)t} dt \\ &= \Re \int_a^b e^{(\alpha + i\beta)t} dt \\ &= \Re \left[\frac{e^{(\alpha + i\beta)t}}{\alpha + i\beta} \right]_a^b \\ &= \left[e^{\alpha t} \frac{\alpha \cos(\beta t) + \beta \sin(\beta t)}{\alpha^2 + \beta^2} \right]_a^b. \end{aligned}$$

Return to Exercise 7.2 on P102

Solution 7.3

Let $\mathbb{C}^- = \mathbb{C} \setminus (-\infty, 0]$ be the slit plane.

- a. Show that any two points in $\mathbb{C}^-$ may be connected by a path in $\mathbb{C}^-$. Hence $\mathbb{C}^-$ is path-connected.
- b. Show that every closed curve $\gamma: [a, b] \rightarrow \mathbb{C}^-$ is null-homotopic in $\mathbb{C}^-$ by finding a homotopy


$$\Gamma: [0, 1] \times [a, b] \rightarrow \mathbb{C}^-, (s, t) \mapsto \Gamma_s(t)$$

satisfying $\Gamma_0(t) = \gamma(t)$ and $\Gamma_1(t) = 1$ for all $t \in [a, b]$. Hence $\mathbb{C}^-$ is simply-connected.

- c. Prove the analogues of a. and b. for a disk $D_r(z_0)$.
- d. Use Cauchy's Theorem to prove that the punctured plane $\mathbb{C}^\times = \mathbb{C} \setminus \{0\}$ is not simply-connected. That is, there exists a closed curve in $\mathbb{C}^\times$ that is not null-homotopic in $\mathbb{C}^\times$.

- a. Every point $z \in \mathbb{C}^-$ can be connected to 1 by a straight line $\gamma: [0, 1] \rightarrow \mathbb{C}$, $\gamma(s) = s + (1 - s)z$, which runs entirely in $\mathbb{C}^-$ since $\gamma(s) \in (-\infty, 0]$ would imply $z = \frac{\gamma(s) - s}{1 - s} \in (-\infty, 0]$, contradicting $z \in \mathbb{C}^-$.

To connect two points $z, w \in \mathbb{C}^-$, first connect z to 1 and then 1 to w .



b. Define

$$\Gamma_s(t) = s + (1 - s)\gamma(t)$$

by straight line interpolation (this method works for every star-shaped set; here 1 is a focal point). Of course, Γ is continuous, being an algebraic expression in continuous functions. We have already proven in (a) that the straight line from $\gamma(t)$ to 1 is contained in $\mathbb{C}^-$. Therefore the image of Γ is contained in $\mathbb{C}^-$. Clearly, $\Gamma_0(t) = \gamma(t)$ and $\Gamma_1(t) = 1$.

c. Same method as for a. and b. but with 1 replaced by the origin.

d. Let $\gamma = \partial D_1(0)$ be the contour $\gamma(t) = e^{it}$, $t \in [0, 2\pi]$. From Example 6.3 we know that

$$\int_{\gamma} \frac{dz}{z} = 2\pi i.$$

If $\mathbb{C}^\times$ were simply-connected, then since $f(z) = 1/z$ is holomorphic on $\mathbb{C}^\times$, Cauchy's Theorem would imply $\int_\gamma f(z)dz = 0$, a contradiction.

[Return to Exercise 7.3 on P102](#)

Solution 7.4

Let $0 < b < 1$.

a. Using the geometric series, find the power series expansion

$$\frac{1}{z - 1/b} = \sum_{n=0}^{\infty} a_n (z - b)^n$$

with center $z_0 = b$ and determine the radius of convergence ρ .

b. Use a. to show that for all $0 < r < \rho$

$$\int_{\partial D_r(b)} \frac{dz}{(z - b)(z - 1/b)} = \frac{2\pi i}{b - 1/b}.$$

c. Use b. to compute

$$\int_0^{2\pi} \frac{dt}{1 - 2b \cos(t) + b^2}.$$

(a) For $q = \frac{z-b}{b-1/b}$ we have

$$\frac{1}{z - 1/b} = \frac{\frac{-1}{1/b - b}}{1 - \frac{z-b}{1/b - b}} = - \sum_{n=0}^{\infty} \frac{(z - b)^n}{(1/b - b)^{n+1}}$$

which converges for $|q| < 1$ and diverges for $q \geq 1$. Equivalently, it converges for $|z - b| < |b - 1/b|$ and diverges for $|z - b| \geq |b - 1/b|$. Hence $\rho = |b - 1/b|$.

(b) We know that power series converge uniformly (Theorem 5.1) and that we may exchange integrals with uniform limits (Corollary 6.1).

Since $1/(z-b)$ is bounded on an annulus $A_{\epsilon, \rho-\epsilon}(b)$, the uniformly convergent series multiplied by $1/(z-b)$ still converges uniformly on $A_{\epsilon, \rho-\epsilon}(b)$ for any $0 < \epsilon < \rho/2$. Since the curve $\partial D_r(b)$ lies in this annulus,

$$\begin{aligned} \int_{\partial D_r(b)} \frac{dz}{(z-b)(z-1/b)} &= - \sum_{n=0}^{\infty} \int_{\partial D_r(b)} \frac{(z-b)^{n-1}}{(1/b-b)^{n+1}} dz \\ &= - \frac{2\pi i}{1/b-b} \end{aligned}$$

The last equality follows by Example 6.3.

- (c) Since $\frac{1}{(z-b)(z-1/b)}$ is holomorphic on $\mathbb{C} \setminus \{b, 1/b\}$ and the curves $\partial D_1(0)$ and $\partial D_r(b)$ are homotopic, we can apply Cauchy's Theorem to get

$$\int_{\partial D_r(b)} \frac{dz}{(z-b)(z-1/b)} = \int_{\partial D_1(0)} \frac{dz}{(z-b)(z-1/b)}.$$

Using the parameterization $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$, $\gamma(t) = e^{it}$, the right integral is

$$\int_0^{2\pi} \frac{ie^{it}}{(e^{it}-b)(e^{it}-1/b)} dt = \int_0^{2\pi} \frac{ib}{1-2b\cos(t)+b^2} dt.$$

Combined with b., we obtain

$$\int_0^{2\pi} \frac{dt}{1-2b\cos(t)+b^2} = \frac{2\pi}{b^2-1}.$$

[Return to Exercise 7.4 on P103](#)

Solution 7.5

Let $\gamma: [0, 1] \rightarrow D$ be a curve in $D \subset \mathbb{C}$ and let $-\gamma: [0, 1] \rightarrow D$, $(-\gamma)(t) = \gamma(1 - t)$ be the opposite curve. Prove that $\gamma * (-\gamma)$ is path-homotopic to a constant loop.

Define

$$\Gamma_s(t) = \begin{cases} \gamma(2t) & \text{if } t \in [0, \frac{1-s}{2}], \\ \gamma(1-s) & \text{if } t \in [\frac{1-s}{2}, \frac{1+s}{2}], \\ (-\gamma)(2(t - \frac{1}{2})) & \text{if } t \in [\frac{1+s}{2}, 1]. \end{cases}$$

Notice that $(-\gamma)(2(\frac{1+s-1}{2})) = \gamma(2(\frac{1-s}{2}))$, so the functions agree on the overlap and $\Gamma_s(t)$ is well-defined. The restriction of Γ to each of the closed sets

$$\begin{aligned} & \left\{ (s, t) \in [0, 1] \times [0, 1] \mid 0 \leq t \leq \frac{1-s}{2} \right\}, \\ & \left\{ (s, t) \in [0, 1] \times [0, 1] \mid \frac{1-s}{2} \leq t \leq \frac{1+s}{2} \right\}, \\ & \left\{ (s, t) \in [0, 1] \times [0, 1] \mid \frac{1+s}{2} \leq t \leq 1 \right\} \end{aligned}$$

is evidently continuous. It follows that Γ is continuous. Moreover, $\Gamma_0 = \gamma * (-\gamma)$ and $\Gamma_1 = \gamma(0)$.

[Return to Exercise 7.5 on P104](#)

A.8 Chapter 8 solutions

Solution 8.1

Compute the following integrals:

- i. $\int_{\partial D_1(0)} \frac{\sin(z)}{z^2} dz,$
- ii. $\int_{\partial D_{1/2}(0)} \frac{\cos(z)}{z-1} dz,$
- iii. $\int_{\partial D_2(1)} \frac{\sin(\cos(z))}{z-1} dz.$

For i., apply the Cauchy integral formula with center $z_0 = 0$,

$$f'(0) = \frac{1}{2\pi i} \int_{\partial D_1(0)} \frac{f(\zeta)}{\zeta^2} d\zeta,$$

to the holomorphic function $f(z) = \sin(z)$ to get $2\pi i \sin'(0) = 2\pi i$ for the integral in i..

Since $\cos(z)/(z-1)$ is holomorphic on $D_{1/2}(0)$, the integral ii. vanishes by the Cauchy theorem (Theorem 7.5).

For iii., apply the Cauchy integral formula to $f(z) = \sin(\cos(z))$. The result is $\sin(\cos(1))$.

[Return to Exercise 8.1 on P114](#)

Solution 8.2

Let $f: \mathbb{C} \rightarrow \mathbb{C}$ be entire. Suppose there is a constant $C > 0$ such that $|f(z)| \leq C|z|^d$ for all $z \in \mathbb{C}$. Prove that $f(z)$ is a polynomial of degree $\leq d$.

Hint: Generalize the proof of Liouville's theorem.

By the higher Cauchy integral formula (Theorem 8.2) we have

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n.$$

Moreover,

$$\begin{aligned} |f^{(n)}(0)| &= \left| \frac{n!}{2\pi i} \int_{\partial D_r(0)} \frac{f(\zeta)}{\zeta^{n+1}} d\zeta \right| \\ &\leq \frac{n!}{2\pi} L(\partial D_r(0)) \frac{Cr^d}{r^{n+1}} = \frac{n!C}{r^{n-d}} \end{aligned}$$

for all $r > 0$. Letting $r \rightarrow \infty$, this implies that $f^{(n)}(0) = 0$ for all $n > d$.

[Return to Exercise 8.2 on P114](#)

Solution 8.3

Let $f: U \rightarrow \mathbb{C}$ be holomorphic and $\overline{D}_r(z_0) \subset U$. Show that

$$f(z_0) = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{i\varphi}) d\varphi$$

and interpret this equation geometrically.

Using the parameterization $\gamma: [0, 2\pi] \rightarrow U$, $\gamma(\varphi) = z_0 + re^{i\varphi}$ to write out the

Cauchy integral formula explicitly, we get

$$\begin{aligned} f(z_0) &= \frac{1}{2\pi i} \int_{\partial D_r(z_0)} \frac{f(z)}{z - z_0} dz \\ &= \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z_0 + re^{i\varphi})}{re^{i\varphi}} ire^{i\varphi} d\varphi \\ &= \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{i\varphi}) d\varphi. \end{aligned}$$

This means that the value of a holomorphic function at the center of a disk is the average of its values over the boundary of the disk.

[Return to Exercise 8.3 on P114](#)

Solution 8.4

Let $f: U \rightarrow \mathbb{C}$ be holomorphic and $\overline{D}_r(z_0) \subset U$.

- a. Prove that $g(x) = f(z_0 + re^{ix})$ is a 2π -periodic function $\mathbb{R} \rightarrow \mathbb{C}$.
- b. Show that the Cauchy integral formula implies an absolutely and uniformly convergent Fourier expansion

$$g(x) = \sum_{n=0}^{\infty} \gamma_n e^{inx}, \quad \forall x \in \mathbb{R},$$

with only non-negative Fourier modes. Moreover, show that

$$\gamma_n = \frac{1}{2\pi} \int_0^{2\pi} g(x) e^{-inx} dx.$$

Note. More generally, a bi-infinite Fourier series $\sum_{n=-\infty}^{\infty} \gamma_n e^{inx}$ can be obtained as a superposition $g_1(x) + g_2(-x)$.

For a., use $e^{2\pi i} = 1$ to compute

$$g(x + 2\pi) = f(z_0 + re^{i(x+2\pi)}) = f(z_0 + re^{ix} e^{2\pi i}) = f(z_0 + re^{ix}).$$

For b., Theorem 8.2 implies that

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n, \quad (\star)$$

with radius of convergence $\rho \geq r$, where

$$a_n = \frac{1}{2\pi i} \int_{\partial D_r(z_0)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta. \quad (\dagger)$$

By Theorem 5.9, power series converge absolutely and uniformly on $D_\rho(z_0)$. Hence we can substitute $z = z_0 + re^{ix}$ into $(\star)$ and get an absolutely and uniformly convergent series

$$g(x) = f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n = \sum_{n=0}^{\infty} a_n r^n e^{inx}.$$

By expanding the contour integral $(\dagger)$ using the definition and the parameterization $\gamma(x) = z_0 + re^{ix}$, we find

$$\gamma_n = a_n r^n = \frac{r^n}{2\pi i} \int_0^{2\pi} \frac{g(x)}{r^{n+1} e^{i(n+1)x}} i r e^{ix} dx = \frac{1}{2\pi} \int_0^{2\pi} g(x) e^{-inx} dx.$$

[Return to Exercise 8.4 on P115](#)

Solution 8.5

Let $f: U \rightarrow \mathbb{C}$ be holomorphic and $\overline{D}_r(z_0) \subset U$. Using polar coordinates show that for the (surface) integral

over the unit disk $\overline{D}_r(x_0) = \{(x, y) \in \mathbb{R}^2 \mid (x - x_0)^2 + (y - y_0)^2 \leq r^2\}$ we have

$$f(z_0) = \frac{1}{\pi r^2} \int_{\overline{D}_r(z_0)} f(x + iy) dx dy$$

and interpret this equation geometrically.

Recall that $dx dy = r d\varphi dr$ for the surface integral in polar coordinates. Hence

$$\begin{aligned} \int_{\overline{D}_r(z_0)} f(x + iy) dx dy &= \int_0^r s \int_0^{2\pi} f(z_0 + s e^{i\varphi}) d\varphi ds \\ &= \int_0^r s \cdot 2\pi f(z_0) ds && \text{by Question 3} \\ &= \pi r^2 f(z_0) && \text{by FTC} \end{aligned}$$

Since the area of $\overline{D}_r(z_0)$ is πr^2 , this means that for a holomorphic function the value $f(z_0)$ at the center is the average value of $f(z)$ over the disk $\overline{D}_r(z_0)$.

[Return to Exercise 8.5 on P115](#)

A.9 Chapter 9 solutions

Solution 9.1

Determine whether the singularities of the following holomorphic functions $f(z)$ are isolated. If so, compute the residue. When applicable, find also the order of the pole or a holomorphic extension.

- i. $\frac{z+i}{z^2+1}$,
- ii. $\frac{1-\cos(z)}{z^2}$,
- iii. $\frac{1}{\cos(z)-1}$,
- iv. $e^{1/z}$

- (i) The singularities $z = \pm i$ are isolated. Since $\frac{z+i}{z^2+1} = \frac{1}{z-i}$, the singularity at $z = +i$ is removable, the holomorphic extension being $\frac{1}{z-i}$, while the

singularity at $z = -i$ is a pole of order one with residue $\text{Res}_{-i}(f) = 1$.

- (ii) The function has an isolated singularity at $z = 0$. The power series expansion of the cosine gives

$$1 - \cos(z) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n)!} z^{2n} = \frac{z^2}{2} - \frac{z^4}{24} + \dots \quad (\star)$$

Hence

$$\frac{1 - \cos(z)}{z^2} = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n)!} z^{2n-2} = \frac{1}{2} - \frac{z^2}{24} + \dots$$

is holomorphic. Therefore, the singularity is removable. The power series is a holomorphic extension. Alternatively, we can extend $f(z) = \frac{1 - \cos(z)}{z^2}$ on $\mathbb{C}^\times$ holomorphically to $\mathbb{C}$ by defining $f(0) = 1/2$.

- (iii) The function $f(z) = \frac{1}{\cos(z)-1}$ has a singularity whenever $\cos(z) = 1$, which is equivalent to $z \in 2\pi\mathbb{Z}$. Each of these singularities is isolated (take a punctured open ball around z of radius $r < 2\pi$). Consider the singularity at the origin.

$$\begin{aligned} \frac{1}{1 - \cos(z)} &= z^{-2} \frac{1}{\frac{1}{2} - \frac{z^2}{24} + \dots} && \text{by } (\star) \\ &= z^{-2} \left[2 + 4 \left(\frac{z^2}{24} + \dots \right) + \dots \right] && \text{as } \frac{1}{1/2 - q} = 2 + 4q + \dots \\ &= z^{-2} \left[2 + \frac{z^2}{6} + \dots \right]. \end{aligned}$$

The second factor in square brackets is a holomorphic function near the origin. Hence the singularity at $z = 0$ is a pole of order 2 with $\text{Res}_{z=0} f(z) = 0$. Since $f(z + 2\pi) = f(z)$, the same holds for the other singularities $z = 2\pi k$, $k \in \mathbb{Z}$.

- (iv) This function has an isolated singularity at $z = 0$, which is an essential singularity: By substituting $1/z$ into the power series expansion of the exponential function, we find

$$e^{1/z} = \sum_{n=0}^{\infty} \frac{z^{-n}}{n!} = 1 + \frac{1}{z} + \frac{1}{2z^2} + \frac{1}{6z^3} + \cdots,$$

which has infinitely many negative terms. The residue is $\text{Res}_{z=0}(e^{1/z}) = 1$.

[Return to Exercise 9.1 on P126](#)

Solution 9.2

Determine whether the domain $D(P)$ of the following formal bilateral series P is non-empty and, if so, find the radii of convergence. i. $\sum_{n=-\infty}^{\infty} \frac{z^n}{|n|!}$,

ii. $\sum_{n=-\infty}^{\infty} z^n$,

iii. $\sum_{n=-\infty}^{\infty} \frac{|n|!}{|n|} z^n$.

- (i) The power series $P_+ = \sum_{n=0}^{\infty} \frac{z^n}{n!}$ and $P_- = \sum_{n=1}^{\infty} \frac{z^n}{n!}$ both have the same radius of convergence $\rho_{\pm} = +\infty$. Hence by Proposition 9.5 the bilateral series has radii of convergence $r = 1/\rho_- = 0$, $R = \rho_+ = +\infty$. In other words, the bilateral series converges on the punctured plane $D(P) = \mathbb{C}^{\times}$.
- (ii) The series $P_+ = \sum_{n=0}^{\infty} z^n$ and $P_- = \sum_{n=1}^{\infty} z^n$ converge precisely for $|z| < 1$. Hence $P_+(z)$ and $P_-(1/z)$ converge simultaneously only if $|z| < 1$ and $|1/z| < 1$, so the intersection $D(P)$ is an empty set.
- (iii) Compute the radius of convergence of $P_+ = \sum_{n=0}^{\infty} \frac{n^n}{n!} z^n$ by the ratio

test:

$$\frac{\frac{(n+1)^{n+1}}{(n+1)!} |z|^{n+1}}{\frac{n^n}{n!} |z|^n} = \left(1 + \frac{1}{n}\right)^n |z| \rightarrow e|z| \quad \text{as } n \rightarrow \infty.$$

(Recall here from calculus that $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e$.) This implies $\rho_+ = e$. Similarly, $P_- = \sum_{n=1}^{\infty} \frac{n^n}{n!} z^n$ has $\rho_- = e$. Hence by Proposition 9.5 we have $r = 1/e$ and $R = e$ for the radii of convergence. In other words, the bilateral series converges on the annulus $A_{1/e, e}(0)$ contained in the domain $D(P)$, which in particular is non-empty.

[Return to Exercise 9.2 on P126](#)

Solution 9.3

Use the geometric series to find the Laurent expansion of

$$f(z) = \frac{1}{z(1-z)}$$

on the domains

- i. $0 < |z| < 1$,
- ii. $0 < |z-1| < 1$,
- iii. $1 < |z| < +\infty$.

(i) On $D_1(0)$ we have $\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n$, hence

$$f(z) = \sum_{n=-1}^{\infty} z^n \quad \text{on the punctured disk } 0 < |z| < 1.$$

(ii) We similarly express $1/z$ in powers of $z - 1$, namely

$$\frac{1}{z} = \frac{1}{1 - (1 - z)} = \sum_{n=0}^{\infty} (-1)^n (z - 1)^n.$$

Hence

$$f(z) = \sum_{n=-1}^{\infty} (-1)^n (z - 1)^n \quad \text{on the punctured disk } 0 < |z - 1| < 1.$$

(iii) Use the geometric series to write

$$\frac{1}{1 - z} = \frac{-1}{z} \frac{1}{1 - 1/z} = - \sum_{n=0}^{\infty} z^{-n-1}.$$

Hence

$$f(z) = - \sum_{n=0}^{\infty} z^{-n-2} \quad \text{on the annulus } A_{1,+\infty}(0).$$

[Return to Exercise 9.3 on P126](#)

Solution 9.4

Let $P = \sum_{n=-\infty}^{\infty} a_n T^n$, $Q = \sum_{n=-\infty}^{\infty} b_n T^n$ be formal bilateral series. Suppose that both domains $D(P, z_0)$, $D(Q, z_0)$ contain a non-empty annulus $A_{r,R}(z_0)$. Assume that for some radius s with $r < s < R$ we have

$$P(z) = Q(z), \quad \text{for all } |z - z_0| = s.$$

Prove the identity theorem, namely that $a_n = b_n$ for all $n \in \mathbb{Z}$.

Pick S with $r < s < S < R$. Applying Theorem 9.6 to $\overline{A}_{s,S}(z_0)$ gives

$$a_n = \frac{1}{2\pi i} \int_{\partial D_s(z_0)} \frac{P(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$$

and

$$b_n = \frac{1}{2\pi i} \int_{\partial D_s(z_0)} \frac{Q(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta.$$

By assumption, $P(\zeta) = Q(\zeta)$ for all $\zeta \in \partial D_s(z_0)$. Hence $a_n = b_n$ for all n .

[Return to Exercise 9.4 on P127](#)

Solution 9.5

Let U be open and $z_0 \in U$. Let $f: U \rightarrow \mathbb{C}$ and $g: U \setminus \{z_0\} \rightarrow \mathbb{C}$ be holomorphic functions.

i. If $f(z_0) \neq 0$ and g has a pole of order one at z_0 , prove that fg has a pole of order one at z_0 with

$$\text{Res}_{z_0}(fg) = f(z_0) \text{Res}_{z_0}(g).$$

ii. If $f(z_0) = 0$ and $f'(z_0) \neq 0$, prove that $1/f(z)$ has a pole of order one at z_0 with $\text{Res}_{z_0}(1/f) = 1/f'(z_0)$.

(i) By assumption,

$$f(z) = a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \dots$$

$$g(z) = \frac{b_{-1}}{z - z_0} + b_0 + b_1(z - z_0) + \dots$$

with $a_0 \neq 0$ and $b_{-1} = \text{Res}_{z_0}(g) \neq 0$. Therefore,

$$f(z)g(z) = \frac{a_0 b_{-1}}{z - z_0} + (a_0 b_0 + a_1 b_{-1}) + (a_1 b_0 + a_2 b_{-1})(z - z_0) + \dots.$$

We read off $\text{Res}_{z_0}(fg) = a_0 b_{-1} = f(z_0) \text{Res}_{z_0}(g) \neq 0$.

- (ii) As U is open, there is a disk $D_r(z_0) \subset U$. By Theorem 8.2, we can expand $f(z) = a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \cdots$ as a power series on $D_r(z_0)$ with $a_0 = f(z_0) = 0$ and $a_1 = f'(z_0) \neq 0$. The series $h(z) = \sum_{n=0}^{\infty} a_{n+1}(z - z_0)^n$ has the same radius of convergence and therefore defines a holomorphic function on $D_r(z_0)$. By construction,

$$f(z) = (z - z_0)h(z), \quad h(z_0) = a_1 = f'(z_0) \neq 0,$$

so

$$\frac{1}{f(z)} = \frac{1}{z - z_0} \frac{1}{h(z)}.$$

Since $h(z_0) \neq 0$, $1/h(z)$ is holomorphic near z_0 and we can apply Part (i) to $1/(z - z_0)$ and $1/h(z)$ to conclude that $\frac{1}{f(z)}$ has a pole of order one with

$$\text{Res}_{z_0} \left(\frac{1}{f(z)} \right) = \text{Res}_{z_0} \left(\frac{1}{z - z_0} \right) \frac{1}{h(z_0)} = \frac{1}{a_1} = \frac{1}{f'(z_0)}.$$

[Return to Exercise 9.5 on P127](#)

References

- Dowson, H. R. 1979. "Serge Lang, Complex Analysis (Addison-Wesley, 1977), Xi + 321 Pp., £1200." *Proceedings of the Edinburgh Mathematical Society* 22 (1): 65–65. <https://doi.org/10.1017/S0013091500027838>.
- Freitag, E., and R. Busam. 2009. *Complex Analysis*. Universitext. Springer Berlin Heidelberg. <https://books.google.co.uk/books?id=3xBpS-ZKlgsC>.
- Howie, J. M. 2012. *Complex Analysis*. Springer Undergraduate Mathematics Series. Springer London. <https://books.google.co.uk/books?id=0FZDBAAAQBAJ>.
- Jameson, G. J. O. 1987. "H. A. Priestley, Introduction to Complex Analysis (Oxford University Press, 1985), 197 Pp., £8.50." *Proceedings of the Edinburgh Mathematical Society* 30 (2): 325–26. <https://doi.org/10.1017/S0013091500028406>.
- Remmert, R. 1991. *Theory of Complex Functions*. Graduate Texts in Mathematics. Springer New York. <https://books.google.co.uk/books?id=uP8SF4jf7GEC>.
- Thomson, B. S., J. B. Bruckner, and A. M. Bruckner. 2001. *Elementary Real Analysis*. Prentice-Hall. https://books.google.co.uk/books?id=6l_E9OTFaK0C.