diffmrc Tutorial

A hands-on walkthrough of the differentiable multireference stack

Version 0.1

This tutorial follows a user from installation to a complete state-averaged calculation with a benchmark comparison. It is paired with the *Manual*, which carries the full reference: the API, the methods and implementation chapters, the verification records, and the development architecture. The package is an early-stage research code; the tutorial states the limitations where they bite.

Contents

1 Introduction **3**
1.1 Who this tutorial is for . . . 3
1.2 What this tutorial covers . . . 3
1.3 Project status . . . 3

2 Installing and running **4**
2.1 Requirements . . . 4
2.2 Runtime settings that are not optional . . . 4
2.3 Checking the installation . . . 4

3 A first calculation **5**
3.1 The script . . . 5
3.2 What each piece does . . . 6
3.3 Reading the result . . . 6

4 State-averaged workflows **6**
4.1 A state-averaged calculation . . . 6
4.2 Reading the root table . . . 7
4.3 State weights . . . 7

5 Choosing an active space **8**
5.1 The default window, and the cases that break it . . . 8
5.2 Auditing a window in one SCF . . . 8
5.3 The corrected windows . . . 8

6 Convergence and diagnostics **9**
6.1 The criterion is the gradient, not the energy . . . 9
6.2 Multiple stationary points . . . 10
6.3 The perturbation layer is the more sensitive one . . . 10

7 A benchmark walkthrough **10**
7.1 The protocol . . . 10
7.2 The runs . . . 11
7.3 Results . . . 11
7.4 Reading the table . . . 11

8 Limitations and how to check them **12**
8.1 Method-family limits 12
8.2 Implementation limits 13
8.3 Protocol limits 13

1 Introduction

`diffmrc` is an end-to-end differentiable multireference electronic structure stack. It spans configuration interaction (CASCI), complete active space self-consistent field (CASSCF), state-averaged CASSCF, and second-order N-electron valence perturbation theory (NEVPT2). Every energy, nuclear gradient, and second derivative the package produces comes out of the same code path by automatic differentiation; there is no hand-derived derivative equation anywhere in the source.

The stack is built on PySCF and pyscfad, with JAX providing the differentiable array operations. All arithmetic is double precision. The package does not replace PySCF: the Gaussian-basis integrals, the reference SCF, and the benchmark comparisons all come from PySCF and pyscfad, while the differentiable multireference chain is original to this codebase.

1.1 Who this tutorial is for

You should be comfortable with the chemistry of active spaces and with running a Python quantum-chemistry calculation from a script. You do not need prior exposure to JAX: the tutorial introduces the runtime settings that matter (`jax_enable_x64`, the CPU platform pin, and the thread budget) right where they are first needed.

1.2 What this tutorial covers

- Chapter 2: installing the package and setting up a working environment.
- Chapter 3: a complete CASSCF nuclear-gradient calculation for water, explained line by line.
- Chapter 4: state-averaged CASSCF with per-state NEVPT2, and how to read the root table it produces.
- Chapter 5: choosing an active space, with the frontier-window rules the benchmark work forced us to write down.
- Chapter 6: what the convergence diagnostics mean and how to act on them.
- Chapter 7: a full benchmark walkthrough on a real target, end to end.
- Chapter 8: where the method family and the present implementation stop being reliable.

The companion *Manual* carries what a tutorial should not: the API reference, protocol tables, the design of the differentiable chain and its second derivatives, the verification records, and the development architecture.

1.3 Project status

This is a research codebase, released to accompany a methods paper and to make its results reproducible. It is not a finished library: interfaces may change without notice, test coverage is partial, and the numerical behaviour of the orbital-optimization and perturbation layers depends on convergence settings that the code reports rather than enforces. Several limitations listed in Chapter 8 are properties of the method family and will not be removed by further development of this package. Use it for research, read the convergence diagnostics it returns, and do not treat it as a black box.

2 Installing and running

2.1 Requirements

Component	Version and role
Python	≥ 3.10
numpy, scipy	required; the analysis layer needs only these
pyscfad	$\geq 0.3.3$; provides the JAX-based `gto` and `scf`
pyscf	≥ 2.5; reference comparisons and integral drivers
jax	≥ 0.4; the differentiable array layer
matplotlib	only for the figure sources

`pyscfad` pins a specific JAX version, so install the package into a dedicated environment rather than into a shared one.

```
conda create -n diffmrc python=3.11
conda activate diffmrc
pip install -e ".[core,dev]"      # kernel, tests and lint tooling
```

For work on the analysis layer alone, `pip install -e ".[dev]"` is sufficient: that layer is plain NumPy and imports nothing from the numerical kernel.

2.2 Runtime settings that are not optional

The platform and thread settings must be in place before the first JAX import; the double-precision switch immediately after it.

```
import os
os.environ['JAX_PLATFORMS'] = 'cpu'      # before importing jax
for v in ('OMP_NUM_THREADS', 'MKL_NUM_THREADS',
          'OPENBLAS_NUM_THREADS', 'NUMEXPR_NUM_THREADS',
          'VECLIB_MAXIMUM_THREADS'):
    os.environ[v] = '8'

import jax
jax.config.update('jax_enable_x64', True)
```

`jax_enable_x64` must be set: the package is validated in double precision, and single-precision runs will produce silently wrong gradients. The CPU platform pin is deliberate. The stack has been developed and verified on CPU only; on a machine with a GPU the default JAX platform would select CUDA, where this build of pyscfad raises a solver error that has no bearing on the chemistry. Pin the platform for now; GPU support is not claimed.

The thread variables keep the BLAS layer inside the thread budget the driver assumes. They are not a performance suggestion: without them a single calculation can oversubscribe a shared node.

2.3 Checking the installation

```
pytest tests/ -v          # unit and integration tests
lint-imports              # verifies the layer boundaries
```

The layer contracts are worth a word even at install time. The package is organized in six layers (kernel, analysis, policy, driver, report, command line), and the import direction between them is enforced as a test. If `lint-imports` is unavailable in your environment, `tests/test_layer_contracts.py` performs the same check. A run that passes the suite but fails the contracts is a run the developers would want to see.

Two environment notes that cost us time, recorded so that they need not cost yours:

- pyscfad's molecule object cannot be traced inside `jax.jit`; the driver layer constructs plain-array callables for anything that is jitted. If you write your own jitted wrapper around a factory from `driver.nevpt2`, use the `theta`-only factories, which are built for that context.
- The first call into a heavy node of the chain compiles an XLA program and can take minutes; subsequent calls are fast. A slow first evaluation is expected, not a hang.

3 A first calculation

This chapter computes a CASSCF nuclear gradient for water at the 6-31G level, with an active space of two electrons in two orbitals (the highest occupied and lowest unoccupied orbital of the reference). It is the smallest complete run through the kernel: a reference SCF, an orbital-rotation optimization, and one nuclear gradient from automatic differentiation.

3.1 The script

```
import os
os.environ['JAX_PLATFORMS'] = 'cpu'
for v in ('OMP_NUM_THREADS', 'MKL_NUM_THREADS',
          'OPENBLAS_NUM_THREADS', 'NUMEXPR_NUM_THREADS',
          'VECLIB_MAXIMUM_THREADS'):
    os.environ[v] = '8'

import numpy as np
import jax
jax.config.update('jax_enable_x64', True)
from jax import numpy as jnp
from pyscfad import gto as pgto, scf as pscf

from diffmrc.core.rotation import block_rotation
from diffmrc.core.orblib import orbital_pairs
from diffmrc.core.optimizer import optimize_theta
from diffmrc.core.grad import (make_energy_fn, make_grad_fn,
                               solve_theta_star, nuclear_gradient)

mol = pgto.Mole()
mol.atom = 'O 0 0 0; H 0.7557 0 0.5880; H -0.7557 0 0.5880'
mol.basis = '6-31g'
mol.build()

nmo = mol.nao_nr()
ncore, nact, nele = 4, 2, 2
pairs = orbital_pairs(ncore, nact, nmo, include_cv=True)

E_cas = make_energy_fn(ncore, nact, nele)
mf = pscf.RHF(mol)
mf.kernel()
C_rhf = jnp.asarray(np.asarray(mf.mo_coeff))

h1e = mol.intor('int1e_kin') + mol.intor('int1e_nuc')
eri = mol.intor('int2e', aosym='s1')

E_theta = lambda t: E_cas(C_rhf @ block_rotation(t, nmo, pairs), h1e, eri,
                          mol.energy_nuc())
th_star, gnorm = solve_theta_star(mol, C_rhf, E_theta, len(pairs),
                                  optimize_theta, steps=2000,
                                  g_conv=1e-12)
```

```
g = nuclear_gradient(mol, make_grad_fn(nmo, pairs, th_star, E_cas, pscf))
print(g)          # shape (natm, 3), in Eh/bohr
```

3.2 What each piece does

The reference SCF is an ordinary spin-restricted Hartree–Fock from pyscfad, whose MO coefficients are the starting point of the chain. The rotation parameter vector `t` is turned into an orbital rotation by `block_rotation`; the pair list from `orbital_pairs` includes the core–virtual rotations, which are required for a correct stationary point — omitting them leaves the optimizer stopped at a point that is not stationary. `E_cas` is the CASCI energy of the rotated orbitals.

`solve_theta_star` runs the two-stage orbital optimization and returns the parameter vector *and* the gradient norm at the point it stopped at. Do not discard `gnorm`: it is the evidence of convergence, and the package reports stationarity from this quantity, not from an optimizer status code.

`nuclear_gradient` applies automatic differentiation to the whole converged chain with respect to the nuclear coordinates, with the orbital parameters held at their converged values. The mathematics of that construction is in the Manual (Chapter on the differentiable chain); for this tutorial the point is that no derivative expression was written anywhere in the package or in this script.

3.3 Reading the result

For this geometry the gradient is small but not zero — the atoms are close to, not at, the CASSCF/6-31G equilibrium. Two checks belong to every such run:

- `gnorm`, the orbital-rotation gradient norm, should sit at or below the threshold that `solve_theta_star` was asked for. If the refinement stalls above the threshold, the returned point carries the stall in its stopping reason; treat a stalled point as a candidate, not as a result.
- The energy is invariant under rotations within a degenerate orbital pair. If a molecule has exact degeneracies, small differences between two runs with different rotation orders are a property of the chart, not a failure of the code.

The next chapter moves from the ground state to several states at once — the setting in which the state-averaged machinery and the per-state perturbation theory come in.

4 State-averaged workflows

Excited states and vertical excitation energies require the orbitals to describe several electronic states at once. The driver layer provides a state-averaged CASSCF construction whose roots are grouped into physical states, with per-state NEVPT2 corrections, all in the same differentiable chain.

4.1 A state-averaged calculation

The smallest complete example is the hydrogen molecule, whose minimal active space holds one singlet ground state, one triplet, and higher singlets.

```
import numpy as np
from jax import numpy as jnp
from pyscfad import gto, scf

from diffmrc.analysis.states import select_groups
from diffmrc.core.nevpt2.impl import cas_dets_for2
from diffmrc.core.orblib import orbital_pairs
```

```
from diffmrc.driver.nevpt2 import (make_nevpt2_sa_theta_fn,
                                   sa_nevpt2_state_table)

mol = gto.Mole()
mol.atom = 'H 0 0 0; H 0 0 0.74'
mol.basis = '6-31g'
mol.build()

nmo = mol.nao_nr()
ncore, nact, nele, nroots = 0, 2, 2, 6
pairs = orbital_pairs(ncore, nact, nmo, include_cv=True)
builder = lambda m: scf.RHF(m)
dets = cas_dets_for2(ncore, nact, nele)

comp = make_nevpt2_sa_theta_fn(mol, ncore, nact, nele, nmo, pairs,
                               builder, nroots=nroots, e2_scale=1.0)
out = comp(jnp.zeros(len(pairs)))

rows, groups = sa_nevpt2_state_table(out, dets)
for g in groups:
    print(g['multiplicity'], g['dE_casscf'], g['dE_nevpt2'])
```

The theta-only factory is the right one whenever the geometry is fixed: it builds the reference SCF and the integrals once and returns a callable of the rotation parameters alone. This is also the factory that can be wrapped in `jax.jit` for orbital optimization, because it never touches the pyscfad molecule at call time.

4.2 Reading the root table

A state-averaged eigensolver returns *roots*, not states, and the difference matters twice over.

First, spin components are degenerate. A triplet contributes three degenerate roots (the $M = +1, 0, -1$ components); a call that asks for six roots of the hydrogen molecule receives three singlets and the three components of one triplet. `sa_nevpt2_state_table` collapses energy-degenerate roots into one group per physical state, labels each group with the multiplicity read from $\langle S^2 \rangle$ of its members, and flags a group whose members disagree on $\langle S^2 \rangle$ rather than guessing.

Second, a per-component contraction can be undefined. The NEVPT2 correction of a pure $|M| = 1$ component can come out as `nan`, and the representative of the group is then chosen as the lowest component whose correction is finite. Treat a group whose entries are all `nan` as a state this protocol cannot report, not as zero.

```
triplets = select_groups(groups, multiplicity=3)   # lowest triplet
singlets = select_groups(groups, multiplicity=1, n=None)
```

4.3 State weights

By default the roots enter the state average with equal weights. When the table is to be read per state, a second pass with between-group equal weights — the components of one state share that state's weight, states share equally — stabilizes the orbitals and usually tightens the final gradient. The two-pass pattern is what the benchmark scripts run; its effect on the reported excitation energies stayed within a few tenths of an eV for the systems in this tutorial's benchmark chapter, so a single equal-weight pass is a reasonable first look.

The weights are passed as `state_weights` to either factory. The vector must have one entry per root, in root order; the helper in the benchmark chapter shows how the group table from the first pass produces it.

5 Choosing an active space

The smallest useful active space for a vertical excitation is the frontier pair: the highest occupied orbital and the lowest unoccupied one, two electrons in two orbitals (or more, when the window must widen; see below). This chapter is about the cases where that default silently selects the wrong orbitals, how to detect them, and what to do instead. The detection rules live in `diffmrc.analysis.orbitals` and cost one SCF to apply.

5.1 The default window, and the cases that break it

The default window takes the frontier pair in energy order. It is correct for the common chromophores: the carbonyl $n \to \pi^*$ pair of aldehydes and ketones, and the $\pi \to \pi^*$ pair of a simple alkene. Widening it is unnecessary there, and this tutorial's benchmark sets it as the baseline.

Two patterns break it.

A linear molecule with two pi systems. In a nitrile, a diyne, or cyanogen, the occupied pi orbitals and the virtual pi orbitals each come as degenerate perpendicular components (x and y). An energy-ordered window can pair the x component of the occupied set with the y component of the virtual set. That excitation is symmetry-forbidden, its exchange integral vanishes, and the computed excitation energy is systematically off — measured at up to 2 eV for the molecules in this tutorial's benchmark set.

An amide. In formamide the conjugated π/π^* pair sits on top of the frontier, while the lone pair that gives the characteristic $n \to \pi^*$ transition lies one orbital lower. The default window computes a $\pi \to \pi^*$ state and compares it to an $n \to \pi^*$ reference: a mismatch of more than 2 eV in the state that was supposed to be reproduced.

5.2 Auditing a window in one SCF

The audit is a per-axis decomposition of the frontier orbitals' p weight. A delocalized single-axis orbital is a pi; an atom-centred single-axis orbital is a lone pair; a window whose HOMO and LUMO sit on different axes while the occupied frontier is degenerate is a cross-component pair.

```
import numpy as np
from diffmrc.analysis.orbitals import heavy_atoms, frontier_report

# mol and mf from a plain pyscfad SCF (see the first-calculation chapter)
labels = mol.ao_labels()
C = np.asarray(mf.mo_coeff)
e = np.asarray(mf.mo_energy)
nocc = mol.nelectron // 2

rep = frontier_report(C, labels, e, nocc, heavy_atoms(labels),
                      target='npi', nelectron=mol.nelectron)
print(rep['homo']['kind'], rep['cross_component'],
      rep['suggested_window'])
```

`frontier_report` returns everything that entered the suggestion — the HOMO-1/HOMO/LUMO identities, the degeneracy map of the occupied frontier, the index of a lone pair below the HOMO when the target needs one — so that the suggestion can be checked rather than trusted. For formamide it reports that the HOMO is a delocalized pi, finds the lone pair two orbitals below, and suggests a three-orbital window spanning lone pair, pi, and pi*.

5.3 The corrected windows

For the two failure modes the recipe is the same three-orbital frontier window, three orbitals instead of two:

- **Linear two-pi molecules**: start the window one orbital lower than the default, so that both components of each degenerate pair are inside and the CI can pick the in-component combinations.
- **Amides**: start the window at the lone pair, three orbitals wide, so that the lone pair, the pi system, and pi* are all active.

One bookkeeping rule travels with the wider window: the electron count must follow the window, not be hard-coded. The scripts in this project use

$$n_{\mathrm{ele}} = N_{\mathrm{ele}} - 2\,n_{\mathrm{core}}, \tag{1}$$

which is what the three-orbital windows above need (four active electrons for the linear two-pi molecules, not two).

The effect on the numbers, from this project's benchmark records:

System	State	default window	corrected window
Cyanogen, Σ_u^- singlet	$\pi\pi^*$	+1.05	+0.62
Cyanogen, Σ_u^+ triplet	$\pi\pi^*$	+1.95	+1.20
Formamide, A'' singlet	$n\pi^*$	+2.12	+0.11

The deviations are against the reference set's best estimates (eV). The corrected formamide window reproduces the reference SC-NEVPT2 value to 0.02 eV. The residual deviations of the linear two-pi systems are a separate story — they survive every window fix and are discussed in Chapter 8.

6 Convergence and diagnostics

A multireference calculation is only as good as the stationary point it was evaluated at, and the failure modes here are quiet ones. This chapter collects what the package reports and how to read it.

6.1 The criterion is the gradient, not the energy

The orbital optimization returns both a point and the gradient norm at that point, and the norm is the verdict. The energy alone cannot serve: this project measured a run in which the energy kept descending while the parameters walked to magnitudes of 10^{13} — the model had no stationary point in the direction it was sliding along, and an energy-only stopping rule called it a success. Treat a convergence claim without a gradient norm as no claim at all.

The stopping reason distinguishes the ways a run can end:

- `CONVERGED`: the gradient reached the refinement target. This is the only reason that is a stationarity claim.
- `STALLED_GRADIENT`: the iteration stopped because a step no longer reduced the gradient. The point may still be useful, but it is a candidate, not a certificate.
- `GUARD_STOPPED`: no damped step lowered the energy. On a converged point this is the normal way to finish a Newton pass.
- `ITERATION_CAP`: the budget ran out. Re-check the gradient before using the point.

6.2 Multiple stationary points

The CASSCF energy is not convex in the orbital parameters. Different starting points can reach different stationary points whose energies differ by more than a bond energy; the water molecule at 6-31G has two minima separated by 13 kcal/mol from the same molecule and basis. Nothing in the eigensolver signals this: both points are stationary.

Practical consequences:

- Report the stationary-point spread alongside any multireference number, or run several starts and keep the lowest.
- A point with zero gradient can still be a saddle. The kernel exposes a curvature check over the Hessian (`stationary.curvature_report`); when a number matters, run it. The orbitals obtained straight from the reference SCF ($\theta = 0$) are, for the water example in this tutorial, a saddle rather than a minimum.
- Treat the spread as protocol, not error: fixed settings can still select different stationary points on different machines, and the package reports the choice it made rather than normalizing it away.

6.3 The perturbation layer is the more sensitive one

NEVPT2 depends on the orbitals more sensitively than the CASSCF energy does. Convergence gates set for the energy can leave the correction unstable: this project measured orbital-gradient stalls that moved the correction by several kcal/mol between runs whose energies agreed to the microhartree. When the target quantity is the correction,

- tighten the orbital-gradient gate rather than the energy gate, and read the gradient norm the run reports;
- confirm the correction's stability across independent runs (different starting parameters, or the two state-averaging passes of Chapter 4), instead of trusting a single converged number.

7 A benchmark walkthrough

This chapter runs the package against an external reference set and shows how to read the comparison. The reference is the QUEST database of vertical excitation energies; the protocol follows its geometry convention, and the numbers below come from this project's recorded runs.

7.1 The protocol

- **Geometries**: the reference set's own optimized geometries (bohr), never re-optimized — a geometry difference is not a method difference, and mixing the two makes the comparison unreadable.
- **Basis**: 6-31G throughout. The reference set reports its values near the complete-basis limit, so each entry carries a basis-set component that this protocol does not remove. It is visible in the deviations below and is the first thing to check when interpreting them.
- **Active space**: the frontier window of Chapter 5, widened to three orbitals for the linear two-pi and amide patterns.
- **State averaging**: equal-weight, with a second between-group pass for most systems; the three window-corrected systems below are quoted from a single equal-weight pass. The pass difference stayed within a few tenths of an eV.

- **Comparison columns**: the reference set reports both its best estimate and its own SC-NEVPT2 value. Since the package implements SC-NEVPT2, the second column isolates method-to-method differences, while the first additionally carries the reference method's own error budget.

7.2 The runs

Each target is a script under `code/`: `t315` (first set), `t317` (second set), `t321` (window-corrected re-runs), with `t316` assembling the summary table from the recorded JSON. A single run is a state-averaged CASSCF with per-state NEVPT2, followed by `sa_nevpt2_state_table` and the state selection of Chapter 4.

7.3 Results

Deviations are in eV; a positive entry means the package's excitation energy lies above the reference.

System	State	package	vs best estimate	vs SC-NEVPT2
Acetaldehyde	$^1A''$	4.58	+0.28	+0.16
	$^3A''$	4.10	+0.15	+0.07
Glyoxal	^1A_u	2.57	−0.31	−0.42
	^3A_u	2.14	−0.35	−0.45
Cyclopropenone	$^1B_1$	4.90	+0.62	+0.82
	$^3B_1$	4.48	+0.55	+0.93
Formamide	$^1A''$	5.75	+0.11	+0.02
	$^3A''$	5.44	+0.08	+0.00
Cyanogen	$^1\Sigma_u^-$	6.99	+0.62	+0.62
	$^3\Sigma_u^+$	6.11	+1.20	+1.19
Propynal	$^1A''$	6.86	+1.35	+1.28
	$^3A'$	5.36	+0.93	+0.70
Cyanoacetylene	$^1\Sigma_u^-$	7.07	+1.28	+1.24
	$^3\Sigma_u^+$	5.92	+1.47	+1.43

The mean absolute deviations, by excitation family — the split is the point of the exercise:

Family	vs best estimate	vs SC-NEVPT2
$n \to \pi^*$ (8 states)	0.31	0.36
$\pi \to \pi^*$ (6 states)	1.14	1.08

7.4 Reading the table

The $n \to \pi^*$ states — the carbonyl and amide chromophores — land within about three tenths of an eV of both reference columns at this basis, which is the accuracy band the reference set itself assigns to SC-NEVPT2. The window-corrected formamide run reproduces the reference SC-NEVPT2 values to 0.02 eV.

The $\pi \to \pi^*$ states carry a uniform positive deviation of roughly 0.6 to 1.5 eV. Window corrections reduce it (cyanogen moves from +1.95 to +1.20 on its triplet) but do not remove it, and the residual has the same sign across these chemically different systems. A uniform, one-signed residual across systems is a property of the method or the protocol, not of the individual molecules; the candidate explanations — valence–Rydberg mixing at the 6-31G basis, and the multiconfigurational character of the $\pi\pi^*$ states beyond a two-orbital window — are discussed in




Figure 1: Deviations of the fourteen benchmark states, in eV: filled markers against the reference set's best estimate, open markers against its SC-NEVPT2 column. The bands group the two excitation families; the figure is the visual form of the two summary rows above.

Chapter 8. The honest summary of this benchmark is a capability map: the package's minimal-space protocol is quantitative for $n \to \pi^*$ vertical excitations and systematically high for $\pi \to \pi^*$ ones at this basis.

8 Limitations and how to check them

The limitations below are grouped by where they come from, because the grouping decides what to do about them: method-family limits will not be removed by further development of this package, implementation limits are measured and tracked, and protocol limits are the user's to manage. Each item comes with the check that detects it.

8.1 Method-family limits

The perturbation layer inherits the standard contracted-NEVPT2 approximations. Excitation energies are reproduced at the accuracy band the reference set assigns to SC-NEVPT2 (Chapter 7), and not better. Check: compare against a second implementation or the reference set's own NEVPT2 column before attributing a deviation to anything else.

Valence–Rydberg mixing is not resolved at small bases. $\pi \to \pi^*$ states that sit near Rydberg states — ethylene is the textbook case — shift by around an electron volt when diffuse functions enter the basis, and move again when the active window includes the diffuse pi orbitals. Check: run the target once with and once without diffuse functions; if the excitation moves by more than the claimed accuracy, the number is basis-limited, not converged.

Multiconfigurational $\pi\pi^*$ states exceed a two-orbital window. The uniform positive deviation of the $\pi \to \pi^*$ family in the benchmark chapter survives every window correction tried here. Check: widen the window by one occupied orbital and re-run; a deviation that shrinks is window-limited, one that does not is a method-family limit for this protocol.

No spin–orbit coupling and no relativistic Hamiltonian. The package works in a scalar-relativistic world with ECPs; fine structure and heavy-element spin physics are out of scope. Check: nothing to check — if the target needs SOC, this is the wrong tool.

8.2 Implementation limits

Dense diagonalization bounds the active space. The CI step builds and diagonalizes the full determinant space; spaces beyond a few thousand determinants become impractical, and iterative eigensolvers are future work.

Memory and compile time grow steeply with the parameter count. The differentiated chain broadcasts intermediates over the rotation parameters; at around five hundred parameters a single node of the Hessian path once requested a 402 GB allocation, which is why the second-order stages now run on Hessian-vector products. Expect the first evaluation of a large system to spend most of its time in XLA compilation. Check: watch the resident memory of the first optimization step, not the steady state.

CPU-only validation. The GPU path is not claimed; the platform must be pinned to CPU (Chapter 2).

8.3 Protocol limits

The convergence gate is the gradient. Review Chapter 6 before quoting any number; the stationary-point spread is part of the result.

The window rules cover the patterns met so far. The frontier audit of Chapter 5 catches the two failure modes this project hit; it is a rule set, not a guarantee, and a chemistry outside the covered patterns should be audited with the same tools rather than assumed correct.

Residual deviations carry protocol, not just method. The benchmark protocol fixes the basis at 6-31G and the geometry at the reference set's; a number quoted without those two facts travels badly. Check: every table in this documentation states basis, geometry source, window rule, and state-weighting pass — keep that habit when exporting numbers.