



Time series modelling in time domain

- Moving average
- Exponential smoothing
- Autocorrelation plot
- Autoregression model

1



Time series analysis

- Characteristics of time series:
 - Trends or other systematic variations
 - Periodic oscillations: with fixed or varying patterns
 - Irregularity: unexpected situations or events or scenarios, spikes in short time span
- Stationary time series (*stationary stochastic process*):
 - Mean, variance, covariances will not change with time
 - White noise: uniform uncorrelated normally distributed variations
- Modelling of time series
 - Reveal the characteristics or structure of the series
 - Facilitate prediction
 - Handle accompanying noises

2



Moving average

Original time series:

$$x_1, x_2, x_3, \dots, x_t, \dots, x_N$$

Smoothed time series:

$$y_2, y_3, \dots, y_t, \dots, y_N$$

Simple moving average:

$$y_t = \frac{1}{M} \sum_{i=1}^M x_{t-i}$$

Centered moving average:
(M is odd)

$$y_t = \frac{1}{M} \sum_{i=-M_2}^{M_2} x_{t-i} \quad M_2 = \frac{M-1}{2}$$

Centered moving average:
(M is even)

$$y_t = \frac{1}{M} \sum_{i=-M_2}^{M_2} x_{t-i}, \quad M_2 = \frac{M}{2}$$

3




Example of moving average

t	x(t)	Simple averaging			Simple moving averaging			Centered moving averaging (M=3)		
		smoothed	error	error squared	smoothed	error	error squared	smoothed	error	error squared
1	71	71	0	0						
2	70	71	-1	1				70,00	0,00	0,00
3	69	71	-2	4	70,00	-1,00	1,00	69,00	0,00	0,00
4	68	71	-3	9	69,00	-1,00	1,00	67,00	1,00	1,00
5	64	71	-7	49	67,00	-3,00	9,00	65,67	-1,67	2,78
6	65	71	-6	36	65,67	-0,67	0,44	67,00	-2,00	4,00
7	72	71	1	1	67,00	5,00	25,00	71,67	0,33	0,11
8	78	71	7	49	71,67	6,33	40,11	75,00	3,00	9,00
9	75	71	4	16	75,00	0,00	0,00	76,00	-1,00	1,00
10	75	71	4	16	76,00	-1,00	1,00	75,00	0,00	0,00
11	75	71	4	16	75,00	0,00	0,00	73,33	1,67	2,78
12	70	71	-1	1	73,33	-3,33	11,11			
SUM=		852		198	709,67		88,67	709,67		20,67
Average=		71	Root of mean square error =		70,967	MSE=		70,967	MSE=	
				4,06			2,72			1,31

4

Example of moving average




5

Example of simple moving average




Example of a moving average on a 12h window



Example of a moving average on a 24h window

6


Example of simple moving average



7

Example of simple moving average


$M = 6$ months



8

Example of simple moving average

M = 12 months



9

Exponential smoothing

Original time series $x_t, t = 1, 2, 3, \dots, N$

New smoothed time series: $y_t, t = 2, 3, 4, \dots, N$

$$y_2 = x_1$$

$$y_t = \alpha x_{t-1} + (1 - \alpha) y_{t-1} \quad 0 < \alpha < 1 \quad t \geq 3$$

$$y_3 = \alpha x_2 + (1 - \alpha) y_2$$

$$y_4 = \alpha x_3 + (1 - \alpha) y_3$$

$$y_5 = \alpha x_4 + (1 - \alpha) y_4$$

.....

10



Exponential smoothing

$$y_t = \alpha \sum_{i=1}^{t-2} \left[(1 - \alpha)^{i-1} x_{t-i} \right] + (1 - \alpha)^{t-2} x_1, \quad t \geq 2$$

Averaging/smoothing: sum of all coefficients is equal to **1**

$$\begin{aligned} & \alpha \sum_{i=1}^{t-2} (1 - \alpha)^{i-1} + (1 - \alpha)^{t-2} \\ &= \alpha \sum_{i=0}^{t-3} (1 - \alpha)^i + (1 - \alpha)^{t-2} \\ &= \alpha \frac{1 - (1 - \alpha)^{t-3+1}}{1 - (1 - \alpha)} + (1 - \alpha)^{t-2} = 1 \end{aligned}$$

$$\sum_{i=0}^n z^i = \frac{1 - z^{n+1}}{1 - z}$$

11



Examples of exponential smoothing

t	x(t)	Alpha = 0,1			Alpha=0,3			Alpha=0,5		
		y(t)	error	error squared	y(t)	error	error squared	y(t)	error	error squared
1	71									
2	70	71,00	-1,00	1,00	71,00	-1,00	1,00	71,00	-1,00	1,00
3	69	70,90	-1,90	3,61	70,70	-1,70	2,89	70,50	-1,50	2,25
4	68	70,71	-2,71	7,34	70,19	-2,19	4,80	69,75	-1,75	3,06
5	64	70,44	-6,44	41,46	69,53	-5,53	30,61	68,88	-4,88	23,77
6	65	69,80	-4,80	22,99	67,87	-2,87	8,25	66,44	-1,44	2,07
7	72	69,32	2,68	7,21	67,01	4,99	24,89	65,72	6,28	39,45
8	78	69,58	8,42	70,83	68,51	9,49	90,10	68,86	9,14	83,55
9	75	70,43	4,57	20,92	71,36	3,64	13,28	73,43	1,57	2,47
10	75	70,88	4,12	16,95	72,45	2,55	6,51	74,21	0,79	0,62
11	75	71,29	3,71	13,73	73,21	1,79	3,19	74,61	0,39	0,15
12	70	71,67	-1,67	2,77	73,75	-3,75	14,06	74,80	-4,80	23,08
SUM=		852		208,82			199,59			181,46
Average=		71	Root of mean square error =					MSE=		
				4,36			4,26		4,06	


12

Examples of exponential smoothing

t	x(t)	Alpha = 0,75			Alpha=0,95			Alpha=0,99		
		y(t)	error	error squared	y(t)	error	error squared	y(t)	error	error squared
1	71									
2	70	71,00	-1,00	1,00	71,00	-1,00	1,00	71,00	-1,00	1,00
3	69	70,25	-1,25	1,56	70,05	-1,05	1,10	70,01	-1,01	1,02
4	68	69,31	-1,31	1,72	69,05	-1,05	1,11	69,01	-1,01	1,02
5	64	68,33	-4,33	18,73	68,05	-4,05	16,42	68,01	-4,01	16,08
6	65	65,08	-0,08	0,01	64,20	0,80	0,64	64,04	0,96	0,92
7	72	65,02	6,98	48,71	64,96	7,04	49,56	64,99	7,01	49,13
8	78	70,26	7,74	59,98	71,65	6,35	40,35	71,93	6,07	36,85
9	75	76,06	-1,06	1,13	77,68	-2,68	7,20	77,94	-2,94	8,64
10	75	75,27	-0,27	0,07	75,13	-0,13	0,02	75,03	-0,03	0,00
11	75	75,07	-0,07	0,00	75,01	-0,01	0,00	75,00	0,00	0,00
12	70	75,02	-5,02	25,17	75,00	-5,00	25,00	75,00	-5,00	25,00
SUM=	852			158,09			142,39			139,66
Average=	71	Root of mean square error = 3,79			3,60			3,56		

13


Examples of exponential smoothing



14


Examples of exponential smoothing

Saudi Arabian oil production in million tones



15

Examples of exponential smoothing



16

Autocorrelation

Data points at different time epochs are correlated

$x_t, \quad t = 1, 2, 3, \dots, N$

Mean of a time series:

$$\bar{x} = \frac{1}{N} \sum_{t=1}^N x_t$$

Variance of a time series:

$$\gamma_0 = \frac{1}{N} \sum_{t=1}^N (x_t - \bar{x})^2$$

Autocovariance function for time lag τ :


$$\gamma_\tau = \frac{1}{N} \sum_{t=1}^{N-\tau} [(x_t - \bar{x}) (x_{t+\tau} - \bar{x})], \quad \tau = 1, 2, 3, \dots$$

Autocorrelation coefficient:

$$\rho(\tau) = \frac{\gamma_\tau}{\gamma_0}$$

17

Autocorrelation coefficients of relative sea level in Stockholm



18



Autoregression (AR) model

$$x_t, \quad t = 1, 2, 3, \dots, N \quad E(x_t) = \mu, \quad Var(x_t) = \sigma^2, \\ E(\varepsilon_t) = 0, \quad Var(\varepsilon_t) = \sigma_\varepsilon^2$$

$$x_t = \sum_{i=1}^p \alpha_i x_{t-i} + \varepsilon_t$$

Coefficients α_i can be determined via least squares adjustment,
or via Yule-Walker equation:

$$\begin{bmatrix} \gamma_1 \\ \gamma_2 \\ \dots \\ \gamma_p \end{bmatrix} = \begin{bmatrix} \gamma_0 & \gamma_{-1} & \gamma_{-2} & \dots \\ \gamma_1 & \gamma_0 & \gamma_{-1} & \dots \\ \gamma_2 & \gamma_1 & \gamma_0 & \dots \\ \dots & \dots & \dots & \dots \\ \gamma_{p-1} & \gamma_{p-2} & \gamma_{p-3} & \dots \end{bmatrix} \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \dots \\ \alpha_p \end{bmatrix}$$

19



AR model for $p=1, 2$

$p=1$

$$x_t = \alpha x_{t-1} + \varepsilon_t$$

$$\gamma_1 = \gamma_0 \alpha_1 \rightarrow \alpha_1 = \frac{\gamma_1}{\gamma_0} = \rho(1)$$

$$\sigma^2 = \alpha^2 \sigma^2 + \sigma_\varepsilon^2 \rightarrow \sigma^2 = \frac{\sigma_\varepsilon^2}{1 - \alpha^2}$$


$p=2$

$$x_t = \alpha_1 x_{t-1} + \alpha_2 x_{t-2} + \varepsilon_t$$

$$\begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} = \begin{bmatrix} \gamma_0 & \gamma_{-1} \\ \gamma_1 & \gamma_0 \end{bmatrix} \begin{bmatrix} \alpha_1 \\ \alpha_2 \end{bmatrix}$$

20

Examples of AR for 0, 1, 2



$p=0$, only white noise

$p=1$, if α is close to 1, mainly white noise, if α is close to 0 it is smoothing

$p=2$, if two α_s are positive, high frequency noise is reduced; if two α_s have opposite signs, output oscillates and change directions

21

Other types of time series models

- Autoregression moving average model (ARMA)

$$x_t = \sum_{i=1}^p \alpha_i x_{t-i} + \sum_{i=1}^q \beta_i \varepsilon_{t-i}$$

- Autoregression integrated moving average model (ARIMA)
- Double (triple) exponential smoothing
- Box-jenkins model

22