Cephalexin synthesis route extraction

Geteilt von Jara1991

Dies ist eine Kopie eines Chats zwischen Claude und Jara1991. Der Inhalt kann nicht verifizierte oder unsichere Inhalte enthalten, die nicht die Ansichten von Anthropic widerspiegeln. Der geteilte Snapshot kann Anhänge und Daten enthalten, die hier nicht angezeigt werden.

Melden

Files hidden in shared chats

Please use this publication for metadata extraction. Extract a single experiment representing the Cephalexin synthesis route.

Create_Experiment >

Experiment created successfully! Here's a summary of what was extracted from the paper:

Experiment ID: 30 — AEH-catalyzed cephalexin synthesis (WT)

Field	Value
Enzyme	WT AEH (*X. campestris pv. campestris*), 200 nM (≈0.0002 mM)
Substrates	PGME (20 mM, acyl donor), 7-ADCA (40 mM, nucleophile)
Products	Cephalexin, Phenylglycine (PG), Methanol
Temperature	25.0 °C
pH	7.0 (100 mM sodium phosphate buffer)

A few notes on the extraction choices:

- The **20 mM PGME / 40 mM 7-ADCA** concentrations represent the baseline condition from the time-course experiments (Fig. 7A starting point)

- The enzyme concentration of **0.0002 mM (200 nM)** was used consistently across all time-course experiments
- PGME concentration was varied from 20–160 mM across the full experiment set — if you'd like separate entries for each concentration condition, I can create those as well



Starten Sie Ihre eigene Konversation