accent '⃗' as m:acc: $\vec{a}$

accent '⃗' as m:acc: $\overrightarrow{AB}$

accent '̇' as m:acc: $\dot{x}$

accent '̈' as m:acc: $\ddot{x}$

accent '̃' as m:acc: $\tilde{f}$

accent '⏜' as m:acc: $\overset{\frown}{ABC}$

accent '̂' as m:acc: $\hat{x}$

accent '⃗' as m:acc: $\vec{F} = m\vec{a}$

accent '̇' as m:acc: $\dot{\omega} = \frac{M}{J}$

bar as m:bar: $\overline{x}$

bar as m:bar: $\overline{AB}$

bar as m:bar: $\overline{AB} \parallel \overline{CD}$

binomial = m:d around noBar m:f: $\binom{n}{k}$

binomial = m:d around noBar m:f: $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

binomial = m:d around noBar m:f: $C_n^k = \binom{n}{k}$

binomial = m:d around noBar m:f: $\binom{10}{3} = 120$

matrix with 2 rows as m:m: $\begin{Bmatrix} x \\ y \end{Bmatrix}$

matrix with 2 rows as m:m: $\begin{pmatrix} x \\ y \end{pmatrix}$

matrix with 3 rows as m:m: $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$

matrix with 2 rows as m:m: $\begin{Bmatrix} x \\ y \end{Bmatrix}$

matrix with 2 rows as m:m: $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$

matrix with 2 rows as m:m: $\begin{vmatrix} a & b \\ c & d \end{vmatrix}$

⏟ as m:groupChr: $\underbrace{a + a + \cdots + a}_{n} = na$

⏞ as m:groupChr: $\overbrace{x_1 + \cdots + x_n}^{n \text{ terms}}$

overset/stackrel as m:limUpp: $a \overset{def}{=} b$

overset/stackrel as m:limUpp: $a \overset{?}{=} b$

percent kept as %: $\Delta U = 5\% \cdot U_n$

percent kept as %: $\eta = \frac{P_2}{P_1} \cdot 100\%$

percent kept as %: $s = 4\%$

percent kept as %: $\% = 5$

percent kept as %: $\frac{5\%}{2}$

percent kept as %: $x_{5\%}$

radical containing a sum: $\sqrt{\frac{1}{N}\sum_{i=1}^{N}(x_i - \mu)^2}$

limit under lim: $\lim_{x \to 0} \frac{\sin x}{x} = 1$

limit under max: $\max_i x_i$

evaluation bar: $\frac{df}{dx}|_{x=0}$

primes and nested scripts: $f''(x) + x_{ij}^2$

delimiters around a fraction stretch (m:d): $\left(\frac{a}{b}\right)$

cases: stretching brace around the stack: $\begin{cases} x^2 & x \geq 0 \\ -x^2 & x < 0 \end{cases}$

plain fractions get no delimiters: $\frac{a}{b} + \frac{c}{d}$

f(x) is not turned into a delimiter: $f(x) = \frac{1}{x}$

lim is upright: $\lim_{n \to \infty} (1 + \frac{1}{n})^n = e$

system of equations: $\begin{matrix} x + 2y = 5 \\ 3x - y = 1 \end{matrix}$

aligned system (second model's style): $\begin{aligned} \dot{x}_1 &= a_{11}x_1 + b_1u \\ \dot{x}_2 &= a_{21}x_1 + b_2u \end{aligned}$

one-line aligned is just the line: $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$

integral as n-ary: $\int_0^{\infty} e^{-x^2}\,dx = \frac{\sqrt{\pi}}{2}$

a superscript minus is not an accent: $x^- + e^-$

bold vectors kept: $\mathbf{F} = m\mathbf{a}$

script L kept: $\mathcal{L}\{f(t)\}$

blackboard R kept: $x \in \mathbb{R}$

operator name rank upright: $\operatorname{rank} A = 2$

operator name adj upright: $\mathbf{A}^{-1} = \frac{1}{det\mathbf{A}} \operatorname{adj}(\mathbf{A})$

operator name sgn upright: $\operatorname{sgn}(x)$

slashed A: normal-size slash overlaid, none left after A: $\widehat{D}^2 = D^2 + i\gamma^5(\partial_5 \not{A}) + i\{\varphi, \partial_5\} - \partial_5^2$.

slashed partial: $(\not{\partial} - m)\psi = 0$

real division a/b is untouched: $v = s/t$

no stray space or empty base in scripts: $x^2 + \partial_5^2$

space before a unit is a preserved non-breaking space: $= 92.3\ \mathrm{N \cdot m}$

space before a unit is a preserved non-breaking space: $= 1.84\ s$

space before a unit is a preserved non-breaking space: $= 1.85\ \mathrm{kW}$

integral with no integrand: no placeholder box: $\int_0^t \neq \int_\alpha^t \neq \int_O^t$

subscript on x-dot, not on an empty base: $\begin{aligned} \dot{x}_1 &= -\frac{R}{L}x_1 \\ \dot{x}_2 &= \frac{1}{C}x_1 \end{aligned}$

\substack with \, keeps two rows: $\sum_{\substack{i=1 \\ i \neq j}}^{n} a_{ij}$

\substack two-line limit: $\lim_{\substack{x \to 0 \\ y \to 0}} f$

prescripts as m:sPre: ${}_{6}^{14}C$

normal integral unchanged: $\int_0^1 x dx$