## SSID: WWCode Password: password



## WOMEN WHO




## ML and Al Study Group

Twitter: @wwcodemanila FB: fb.com/wwcodemanila

#WWCodeManila #YourProgrammingLanguage #StudyGroup



## New Member's Introduction



#### **OUR MISSION**

Inspiring women to excel in technology careers.





#### **OUR VISION**

A world where women are representative as technical executives, founders, VCs, board members and software engineers.





#### STUDY GROUP

Study groups are events where women can come together and help each other learn and understand a specific programming language, technology, or anything related to coding or engineering.

#### **GUIDELINES**

- If you have a question, just ask
- If you have an idea, share it
- Make friends and learn from your study groupmates
- Do not recruit or promote your business

# SHOW & TELL

## LINEAR REGRESSION IN ONE VARIABLE

(a.k.a Univariate Linear Regression)

### Agenda

- 1. Review
- 2. Hypothesis Function
- 3. Cost Function
- 4. Gradient Descent Algorithm

#### Review

- Supervised vs. Unsupervised

#### Review


- Supervised vs. Unsupervised
- Supervised Learning:
  - Classification output variables take discrete class labels
  - **Regression** output variable takes continuous values


#### Review

- Supervised vs. Unsupervised
- Supervised Learning:
  - Classification output variables take discrete class labels
  - **Regression** output variable takes continuous values
- Univariate Linear Regression predicts a single output y from a single input value x


#### **Linear Regression in One Variable**

e.g. **Predicting Profit** of a City






We want to **fit** a line through the data.



We want to **fit** a line through the data.


And make it so that the line **generalizes** the data well.




We want to **fit** a line through the data.

And make it so that the line **generalizes** the data well.


e.g. If Imus City has a population of about 21k, then its predicted profit is...?




How do we come up with the "right" line?



How do we come up with the "right" line?



How do we come up with the "right" line?




#### **Equation of a Line**


- Recall the equation of a line?

#### **Equation of a Line**

- Recall the equation of a line?

$$y = mx + b$$





#### **Exercise:**

Draw the lines:

$$y = \frac{3}{2}x + 1$$
$$y = -2x - 5$$

#### **Equation of a Line**

- Recall the equation of a line?

$$y = mx + b$$

- More generally,

$$y = \theta_1 x + \theta_0$$

where  $\theta_0$ ,  $\theta_1$  are called "weights".  $\theta_0$  has a special name called the "bias"

- We need to find the right values for  $\theta_0$  and  $\theta_1$ .
- How do we get the right values?

- We need to find the right values for  $\theta_0$  and  $\theta_1$ .
- How do we get the right values?
- We start with a guess (or *hypothesis*) using random weights

- We need to find the right values for  $\theta_0$  and  $\theta_1$ .
- How do we get the right values?
- We start with a guess (or *hypothesis*) using random weights
- We define our **hypothesis function** as

$$h(x) = \theta_1 x + \theta_0$$

where  $\theta_0$  and  $\theta_1$  are set to some random values (e.g.  $\theta_0 = 0$ ,  $\theta_1 = 0$ )

- Let x be the input features and y be the true output values.
- Suppose we come up with two hypotheses:

Input x	Output y	$h_{\theta}(x)$ $\theta_0 = 2,  \theta_1 = 3$	$h_{\theta}(x)$ $\theta_0 = 4,  \theta_1 = 3$
0	4	?	?
1	8	?	?
2	9	?	?
3	13	?	?

- Let x be the input features and y be the true output values.
- Suppose we come up with two hypotheses:

Input x	Output y	$h_{\theta}(x)$ $\theta_0 = 2,  \theta_1 = 3$	$h_{\theta}(x)$ $\theta_0 = 4,  \theta_1 = 3$
0	4	2	?
1	8	5	?
2	9	8	?
3	13	11	?


#### How good is our hypothesis?

- We need a way to measure how close our hypothesis  $h_{\theta}(x)$  is to the true output values y.
- i.e. we want a function that measures how good or bad our hypothesis function performs

#### How good is our hypothesis?

- We need a way to measure how close our hypothesis  $h_{\theta}(x)$  is to the true output values y.
- i.e. we want a function that measures how good or bad our hypothesis function performs
- We will call such a function the **Cost Function**, which will measure the average error of the  $h_{\theta}(x)$ .

#### **Cost Function**



Note: The error is squared and the mean is halved as a convenience for computing the gradient descent later.

#### **Minimizing the Cost Function**

- We want to minimize the cost function, or the mean squared error (MSE).

#### **Minimizing the Cost Function**

- We want to minimize the cost function, or the mean squared error (MSE).
- In other words, we want to find the  $\theta_0$  and  $\theta_1$  that minimizes  $J(\theta_0,\theta_1)$

#### **Minimizing the Cost Function**

Input x	Output y
0	0
1	2
2	4
3	6

Suppose  $\theta_0 = 0$ .

- 1. Plot  $h_{\theta}(x)$  when:  $\theta_1 = 1$ 

  - $\theta_1 = 2$   $\theta_1 = 3$


2. Solve for:

• 
$$J(\theta_1 = 1)$$

• 
$$J(\theta_1 = 2)$$


• 
$$J(\theta_1 = 3)$$


#### $J(\theta)$ vs $\theta$ plot



- A parabola makes sense, since *J* is a quadratic function.
- The value of  $\theta_1$ that minimizes J is 2.

## **Minimizing J**





In general,


the values of the weights  $\theta_1$ ,  $\theta_0$  that minimize  $J(\theta_1, \theta_0)$ 



minimizes the distance between h(x) and every data point

- Have some function  $J(\theta_0, \theta_1)$
- Want to find  $\min_{\theta_0,\theta_1} J(\theta_0,\theta_1)$

- Have some function  $J(\theta_0, \theta_1)$
- Want to find  $\min_{\theta_0, \theta_1} J(\theta_0, \theta_1)$
- Outline of Gradient Descent Algorithm
  - Start with some  $\theta_0$ ,  $\theta_1$
  - Keep changing  $\theta_0, \theta_1$  to reduce  $J(\theta_0, \theta_1)$  until we hopefully end up at the minimum.




- We want to take steps down the cost function in the direction of the steepest descent
  - The direction is given by the derivative of the cost function:


$$\frac{\partial}{\partial \theta_j} J(\theta_j)$$
 (for  $j = 0$  and  $j = 1$ )

- The **size** of of each step is given by some learning rate  $\alpha$ .

repeat until convergence:




## Let's talk about derivatives for a sec.



To get to the minimum value, we take the **derivative** of our cost function.

$$\frac{\partial}{\partial \theta_1} J(\theta_1)$$


**Derivative** – the slope of the tangent line at a point




To get to the minimum value, we take the **derivative** of our cost function.

$$\frac{\partial}{\partial \theta_1} J(\theta_1)$$

**Derivative** – the slope of the tangent line at a point




What is the slope at the minimum of *J*?



What is the slope at the minimum of *J*?

The slope given by  $\frac{\partial}{\partial \theta_1} J(\theta_1)$  will always be zero!

#### Covergence




We say that our algorithm has converged when

$$\frac{\partial}{\partial \theta_1} J(\theta_1) = 0$$


$$\theta_1$$
: =  $\theta_1 - \alpha(0)$ 

 $\theta_1$  ceases to update to a new value.

#### **Derivative Intuition**




#### **Learning Rate Intuition**



If  $\alpha$  is too small, gradient descent can be slow.

If  $\alpha$  is too large, gradient descent can overshoot the minimum. It may fail to converge, or even diverge.



```
repeat until convergence { \theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1) \quad \text{(for } j = 0 \text{ and } j = 1) }
```

#### Correct: Simultaneous update

```
temp0 := \theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)
temp1 := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)
\theta_0 := temp0
\theta_1 := temp1
```

```
repeat until convergence { \theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1) \quad \text{(for } j = 0 \text{ and } j = 1) }
```

#### Correct: Simultaneous update

$$temp0 := \theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$$

$$temp1 := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$$

$$\theta_0 := temp0$$

$$\theta_1 := temp1$$

#### Incorrect:

$$\begin{array}{l} \operatorname{temp0} := \theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1) \\ \theta_0 := \operatorname{temp0} \\ \operatorname{temp1} := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1) \\ \theta_1 := \operatorname{temp1} \end{array}$$

```
repeat until convergence: {
                                        	heta_0 := 	heta_0 - lpha \, rac{1}{m} \sum_{i=1}^m (h_	heta(x_i) - y_i)
                                        	heta_1 := 	heta_1 - lpha \, rac{1}{m} \sum_{i=1}^m ((h_	heta(x_i) - y_i) x_i)
```

# Partner/Group/Individual Exercise

# Partner/Group/Individual Presentation

# References

## T.I.L.

#### SHARE IT! In front!

On Twitter: @wwcodemanila

Or FB: fb.com/wwcodemanila

Don't forget to tag WWCodeManila so we can retweet or share it.

#### Feedback Form

https://goo.gl/YzSqcS

Please don't rate the event on meetup.

Not helpful. It is best to just tell your concerns via the feedback form. We are a building a community not a Yelp restaurant.

# THANK YOU:)